

EVALUATION OF THE EFFECTIVENESS OF TWO LOCAL DRUG DELIVERY AGENTS IN THE MANAGEMENT OF CHRONIC PERIODONTITIS – A COMPARATIVE STUDY

Abstract

Background: The prime aim of periodontal therapy is the arrest of bacteria/host induced inflammation and to maintain healthy periodontium. Pathogenic organisms which are not mechanically accessible can be eliminated by antibacterial agents. Several local antimicrobial drug delivery agents have been developed with a view to maintain adequate levels of crevicular fluid at the target site and preventing systemic uptake. Aim: Tetracycline fibres (Periodontal plus AB) and curcumin gel (curenex) are two such drugs used as adjuncts to scaling and root planing in the treatment of chronic periodontitis. Comparative evaluation of the effectiveness of these two agents will be helpful in the selection of a particular type of treatment modality. Methodology: Eighty sites from 30 patients were selected for the study and were divided into two groups of 40 sites for tetracycline fibres and 40 sites for curcumin gel application as adjuncts to scaling and root planing in the treatment of chronic periodontitis. Clinical parameters such as gingival index, plaque index, oral hygiene index, probing pocket depth and gain in clinical attachment level were assessed in all the patients after 30 days and 90 days in both groups. Results and discussion: Study revealed that both tetracycline fibres and curcumin gel are effective agents as adjuncts to scaling and root planing in the treatment of chronic periodontitis. Conclusion: Though both tetracycline fibers and curcumin gel were found to be effective for the treatment of chronic periodontitis, tetracycline was found to be slightly more effective. This may be due to the property of substantivity exhibited by tetracycline.

Key words: Periodontitis, Local drug delivery, tetracycline fibers, curcumin gel

Introduction

Chronic periodontitis is a multifactorial disease resulting from the formation of periodontal pocket and progressive loss of attachment. Periodontal pocket formation represents the pathologic sequelae of microbial and inflammation mediated degradation of collagenous connective tissue and alveolar bone. Mechanical debridement may fail to eliminate the putative pathogens from the pockets completely because of the inaccessibility of the location of the organisms within the gingival tissue. Systemic administration of drugs leads to absorption into the blood stream and distribution throughout the body. Local drug delivery mode is considered more beneficial in the control of localized periodontal disease as adjunct to non-surgical periodontal therapy. Commonly used antimicrobial local drug delivery agents include Chlorhexidine- containing chip (Periochip), Doxycycline gel (Atridox), Minicycline microspheres

(Arestin), ethylene or vinyl acetate copolymer fiber containing tetracycline and bioresorbable tetracycline fiber (Periodontal Plus AB). Curcumin Longa has been used as a therapeutic agent in ayurvedic medicine for years. A comparative study on the clinical effectiveness of a synthetic drug tetracycline fiber (which may have side effects) and an ayurvedic drug curcumin gel (supposed to have no side effects) is considered worthwhile.

Review of Literature

Several authors have found that effective treatment of periodontitis is the mechanical debridement to remove biofilm and calculus from the affected root surfaces followed by adjunct therapy by local delivery of drugs (1 – 4) The advantages of local drug delivery are:

1. Achieving high concentration of the drug at localized site (100 times as compared to systemic therapy)
2. Suitable for agents which cannot be given systemically (such as chlorhexidine)
3. No systemic side effects
4. Low chances of super-infection or drug resistance
5. Reduction in total drug usage, drug accumulation and frequency

The adjunctive use of local delivery of drugs may enhance the results in isolated periodontal pocket sites that do not respond to conventional therapy (5). Purucker et al., (1994) compared the clinical response of local versus systemic treatment as adjuncts to scaling and root planing and observed that local delivery of tetracycline fiber and the systemic administration of amoxicillin/clavulinate given for 3 months after scaling and root planing produced similar clinical outcomes over the 9-month observation period. Newman et al., (1994) in their evaluation study to compare the efficacy of scaling and root planing (SRP) alone versus SRP plus adjunct therapy with tetracycline fiber observed that SRP plus adjunct therapy was more effective in the treatment of periodontitis. Kinane et al., (1999) compared the effectiveness of three periodontal local antimicrobial therapies: scaling and root planing alone, in conjunction with 25% tetracycline, 2% minacycline gel or 20% metronidazole gel and observed that application of three locally applied antimicrobial systems offered benefits and also that tetracycline gave the best effect.

Curcumin (diferuloylmethane), the main bioactive component of turmeric, is reported to have anti-inflammatory (8-10), antioxidant (11 - 14), anti-allergic, anticarcinogenic, antimutagenic, anticoagulant, antidiabetic, antibacterial, antiprotozoal, antifungal, antiviral, anti-ulcer, hypotensive and hypocholesterolemic activities (15,16). Kudva et al., (2012) in a comparative study on the adjunctive efficacy of turmeric, curcumin and traditional nonsurgical methods for treating periodontal pockets found that plaque index and gingival index showed significant

improvement in curcumin treated group. Several authors have recommended the local delivery of curcumin as an adjunct therapy in the treatment of chronic periodontitis (18-20).

Materials and Methods

Thirty patients in the age group 30 – 55 years attending the Department of Periodontics, PSM College of Dental Science & Research with chronic periodontitis having periodontal pocket depths of 5 – 8 mm after scaling and root planing were selected for the study. Ethical clearance was obtained for the study from the Institutional ethics committee (No.17/Ethics/PSMCDSR/2018 dt. 04/12/2018). A total of 80 sites (40 sites for each group) were included for evaluating the effectiveness of curcumin gel and tetracycline fibers. The sites were grouped as follows:

Group A: Application of curcumin gel – 40 sites

Group B: Application of tetracycline fibers – 40 sites

Tetracycline fibres were soaked and placed into the pocket with the help of a tweezer and periodontal probe was used to insert into periodontal pocket with gentle pressure. Curcumin gel was applied directly from a disposable syringe into the pocket without traumatizing or damaging the periodontal tissue. After insertion of the dug locally, the region was secured with Coe-pak™ (GC American Inc, USA.)

The clinical parameters: Plaque index, Oral hygiene index, Gingival index, Probing pocket depth, and Clinical attachment level was assessed using the techniques employed by Newman et al., (2002). Clinical parameters were assessed before starting the therapy and recorded as baseline values. All patients were refrained from chewing hard and sticky food, flossing on the treated side and not to disturb the area with tongue, finger or tooth pick. For the removal of coe-pak the participants were recalled after 1 week. Clinical parameters were reassessed at 30 days and 90 days and recorded.

Results

Gingival index: Mean gingival index showed significant decrease ($p < 0.001$) on treatment with curcumin and tetracycline observed at 30 days when compared to baseline level. No significant difference was seen at 90 days when compared to 30 days. The results obtained are represented in Table 1.

Table 1: Gingival index values in curcumin gel and tetracycline fibers treated patients with chronic periodontitis.

Group	No. of sites	Baseline	30 days	90 days
Curcumin	40	2.03 \pm 0.16	1.02 \pm 0.15*	1.02 \pm 0.15 ^{ns}
Tetracycline	40	2.15 \pm 0.18	1.04 \pm 0.14*	1.04 \pm 0.14 ^{ns}

Values are expressed as mean \pm SD. The values obtained at 30 days were compared with those obtained at baseline level. The values obtained at 90 days were compared with those obtained at 30 days. Independent 't' test was performed to calculate p value.

*Significant P <0.001., ns: Not significant

Oral hygiene index: Mean oral hygiene index values showed significant improvement ($p < 0.001$) on treatment with curcumin and tetracycline observed at 30 days when compared to baseline level. No significant difference was seen at 90 days when compared to 30 days in either curcumin or tetracycline treated groups. The results are depicted in Table 2.

Table 2: Oral hygiene index values in curcumin gel and tetracycline fibres treated patients with chronic periodontitis.

Group	No. of sites	Baseline	30 days	90 days
Curcumin	40	1.08 \pm 0.27	0.08 \pm 0.01*	0.08 \pm 0.01 ^{ns}
Tetracycline	40	1.05 \pm 0.22	0.05 \pm 0.01*	0.05 \pm 0.01 ^{ns}

Values are expressed as mean \pm SD. The values obtained at 30 days were compared with those obtained at baseline level. The values obtained at 90 days were compared with those obtained at 30 days. Independent 't' test was performed to calculate p value.

*Significant P <0.001., ns: Not significant

Plaque index: Mean plaque index showed significant decrease ($p < 0.001$) on treatment with curcumin and tetracycline observed at 30 days when compared to baseline level. No significant difference was seen at 90 days when compared to 30 days in either curcumin or tetracycline treated group. The results are given in Table 3.

Table 3: Plaque index values in curcumin gel and tetracycline fibres treated patients with chronic periodontitis.

Group	No. of sites	Baseline	30 days	90 days
Curcumin	40	1.98 \pm 0.36	0.98 \pm 0.15*	0.98 \pm 0.15 ^{ns}
Tetracycline	40	2.05 \pm 0.32	1.05 \pm 0.22*	1.05 \pm 0.22 ^{ns}

Values are expressed as mean \pm SD. The values obtained at 30 days were compared with those obtained at baseline level. The values obtained at 90 days were compared with those obtained at 30 days. Independent 't' test was performed to calculate p value.

* Significant P <0.001., ns: Not significant

Probing pocket length:

Mean probing pocket length showed significant improvement ($p < 0.001$) on treatment with curcumin and tetracycline observed at 30 days when compared to baseline level. No significant difference was seen at 90 days when compared to 30 days in either curcumin or tetracycline treated groups. The results are depicted in Table 4.

Table 4: Probing pocket depth values in curcumin gel and tetracycline fibers treated patients with chronic periodontitis.

Group	No. of sites	Baseline	30 days	90 days
Curcumin	40	5.10 \pm 0.30	4.28 \pm 0.51*	4.28 \pm 0.51 ^{ns}
Tetracycline	40	5.05 \pm 0.22	3.78 \pm 0.53*	3.73 \pm 0.55 ^{ns}

Values are expressed as mean \pm SD. The values obtained at 30 days were compared with those obtained at baseline level. The values obtained at 90 days were compared with those obtained at 30 days. Independent 't' test was performed to calculate p value.

*Significant $P < 0.001$., ns: Not significant

Clinical attachment level: Mean clinical attachment level showed significant improvement ($p < 0.001$) on treatment with curcumin and tetracycline observed at 30 days when compared to baseline level. No significant difference was seen at 90 days when compared to 30 days in either curcumin or tetracycline treated groups. The results are given in Table 5.

Table 5: Clinical attachment level values in curcumin gel and tetracycline fibres treated patients with chronic periodontitis.

Group	No. of sites	Baseline	30 days	90 days
Curcumin	40	2.03 \pm 0.16	1.02 \pm 0.15*	1.02 \pm 0.15 ^{ns}
Tetracycline	40	2.15 \pm 0.18	1.04 \pm 0.14*	1.04 \pm 0.14 ^{ns}

Values are expressed as mean \pm SD. The values obtained at 30 days were compared with those obtained at baseline level. The values obtained at 90 days were compared with those obtained at 30 days. Independent 't' test was performed to calculate p value.

*Significant $P < 0.001$., ns: Not significant

Discussion

Periodontitis is a biofilm associated disease and are difficult to treat with antibiotics unless biofilm is disrupted mechanically. Chronic periodontitis develops as a result of host-bacterial interaction in which the pathogens produce harmful byproducts that initiate host immune inflammatory responses leading to the breakdown of extracellular matrices and bone resorption, creating bone defects eventually leading to tooth loss (23). It is characterized by the

presence of gingival inflammation, periodontal pocket formation, clinical attachment loss and development of osseous deformities around the affected teeth. Scaling and root planing in conjunction with antibacterial agents were administered either systemically or locally (24). However, systemic administration has been found to be associated with side effects and local delivery of drugs was found more beneficial (24,25).

In this study, effort has been made to compare the effectiveness of two commercially available local delivery agents – curcumin gel (curenex - an ayurvedic drug) and tetracycline fibres (Periodontal plus AB - an allopathic drug) in treating chronic periodontitis. Statistically significant improvements were observed in all the parameters measured at 30 days compared to the values at baseline level, the response was slightly better in the case of tetracycline treated group. No difference was observed when the values at 90 days were compared with those at 30 days. The findings were in agreement with earlier studies (19,23,24).

In this study microbial evaluation was not carried out. Long term studies with a large sample size are required to further assess the clinical efficacy of both local delivery drugs.

Conclusion

It may be concluded that tetracycline fibers (Periodontal Plus AB) and curcumin gel (Curenex) can be used as adjuncts to scaling and root planing in the treatment of chronic periodontitis. Application of both agents showed significant beneficial effects, but tetracycline showed slightly better effect. The more beneficial effect of tetracycline may be due to the property of substantivity exhibited by this drug. Curcumin being an ayurvedic drug with little or no side effects can be an excellent alternative to tetracycline in treatment of chronic periodontitis.

References

1. Drisco C. Non-surgical periodontal therapy. *Periodontology*. 2001;25(1):77 – 78.
2. Drisco C. Periodontal debridement still the treatment of choice. *J Evidence based Dent Practice*. 2014;14: 33-41.
3. Pihlstrom B. Commentary: Treatment of periodontitis: Key Principles. *J Periodontology*. 2014;85(5):655-656.
4. Sanz I, Alonso B, Carasol M, Herrera D, Sanz M. Nonsurgical treatment of periodontitis. *J Evidence Based Dent Practice*.2012;12(3): 76-86.
5. Verma E, Belludi SA, Banthian R. Local drug delivery with chlorhexidine chip and tetracyclinefibers as an adjunct to mechanical therapy in isolated periodontal pockets – A case report. *Int J Clin Dentistry*.2011;4(4):383-90.

6. Purucker P, Mertes H, Goodson H, Bernimoulin JP. Local versus systemic adjunctive antibiotic therapy in 28 patients with generalized aggressive periodontitis. *J Periodontology*. 2001;72(9):1241-45.
7. Newman MG, Kornmann KS & Doherty FM. A six-month multicentre evaluation of adjunctive tetracycline fiber therapy used in conjunction with scaling and root planing in maintenance patients: Clinical results. *J Periodontology*. 1994;65(7): 685-91.
8. Sajithlal GB, Chithra P, Chandrakasan G. Effect of curcumin on the advanced glycation and cross linking of collagen in diabetic rats. *Biochemical Pharmacology*.1998;56(12):1607-14.
9. Chainani-Wu N. Safety and anti-inflammatory activity of curcumin: a component of turmeric (*Curcuma Longa*). *J Alternative & Complementary Medicine*. 2003;9(1):161-68.
10. Motterlini R, Foresti R, Bassi R, Green CJ. Curcumin, an antioxidant and anti-inflammatory agent, induces heme oxygenase-1 and protects endothelial cells against oxidative stress. *Free Radical Biology & Medicine*. 2000;28(8):1303-12.
11. Bhatia M, Urolagin SS, Pentyala KB, Urolagin SB, Menaka KB, Bhoi S. Novel therapeutic approach for the treatment of periodontitis by curcumin. *J Clinical & Diagnostic Res*.2014;8(12): ZC65-ZC69.
12. Ramsevak RS, DeWitth DL, Nair MG. Cytotoxicity, antioxidant and anti-inflammatory activities of curcumin I-III from *Curcuma Longa*. *Phytomedicine*.2000;7(4):303-08.
13. Osawa T, Sugiyama Y, Inayoshi M, Kawakishi S. Antioxidant activity of tetrahydrocurcuminoids. *Bioscience, Biotechnology & Biochemistry*. 1995;59(9):1609-12.
14. Elavarasu S, Suthanthiran T, Thangavelu A, Alex S, Palanisamt A, Kumar T S. Evaluation of super oxide dismutase levels in local drug delivery system containing 0.2% curcumin strip as an adjunct to scaling and root planing in chronic periodontitis: A clinical and biochemical study. *J Pharmacy Bioallied Sciences*. 2016;8(Suppl 1): S46.
15. Siddiqui AM, Cui X, Wu R, Dong W, Zhou M, Hu M, Simms HH, Wang P. The anti-inflammatory effect of curcumin in an experimental model of sepsis mediated by upregulation of peroxisome proliferator-activated receptor- γ . *Critical Care Medicine*. 2006;34(7):1874-82.
16. De R, Kundu P, Swarnakar S, Ramamurthy T, Chowdhury A, Nair GB, Mukhopadhyaya AK. Antimicrobial activity of curcumin against *Helicobacter pylori* isolates from India and during infections in mice. *Antimicrobial Agents and Chemotherapy*.2009;53(4):1592-97.

17. Kudva P, Tabassum ST, Gupta S. Comparative evaluation of the efficacy of turmeric and curcumin as a local drug delivery system – A Clinico-microbiological study. *General Dentistry*. 2012;60(5): e283-87.
18. Anitha V, Rajesh P, Shanmugham M, Priya R M, Prabhu S, Shivakumar V. Comparative evaluation of natural curcumin and synthetic chlorhexidine in the management of chronic periodontitis as a local drug delivery: A clinical and microbiological study. *Ind J Dental Res*. 2015;26(1): 57-60.
19. Siddharth M, Singh P, Gupta R, Sinha A, Shree S, Sharma K. A comparative evaluation of subgingivally delivered 2% curcumin and 0.2 % chlorhexidine gel adjunctive to scaling and root planing in chronic periodontitis. *J Contemp Dental Practice*. 2020;21(5):494-99.
20. Raghava KV, Sistla KP, Narayan SJ, Yadalam U, Bose A, Mitra K. Efficacy of curcumin as an adjunct to scaling and root planing in chronic periodontitis patients. – A randomized controlled clinical trial. *J Contemp Dent Practice*. 2019;20(7):842-46.
21. Newman M, Takei H, Caraza F. Caraza's Clinical Periodontology, Philadelphia, WB Sanders Co., 2002.
22. Gupta R, Pandit N, Aggarwal S, Verma A. Comparative evaluation of subgingivally delivered 10% doxycycline hyclate and xanthan-based chlorhexidine gel in the treatment of chronic periodontitis. *J Contemp Dent Practice*. 2008;9(7):25-32.
23. Kirkwood KL, Cirelli JA, Rogers JE, Giannobile WV. Novel host response therapeutic approaches to treat periodontal disease. *Periodontology*. 2000;43: 294 – 98.
24. Reynolds MA, Kao RT, Nares S, Camargo PM, Caton JG, Clem DS et al., Periodontal regeneration –intra-bony defects: Practical applications from the AAP regeneration workshop. *Clinical Advances in Periodontics*. 2015;5(1): 21-29.
25. Rupali K, Vandana KL, Shobha P. Effect of local drug delivery in chronic periodontitis patient: A meta-analysis. *J Ind Soc Periodontol*. 2011;15(4): 304-09.