

Histopathological and Oxidative Stress Response of *Oreochromis niloticus* Exposed to Varying Concentrations of Sawdust Extract

ABSTRACT

Indiscriminate disposal or dumping of wood wastes such as sawdust into water bodies has reached a menacing proportion with projected environmental and ecological consequences. This study therefore evaluated the effect of sawdust extract exposure on cultured water quality, histological and stress response mechanisms of *Oreochromis niloticus* under laboratory conditions. *O. niloticus* were divided into four groups (0%, 1/10, 1/100 and 1/1000 of LC₅₀) respectively in triplicates. Heavy metals composition was evaluated using Atomic Absorption Spectrophotometer. Phytochemical analysis of sawdust, physico-chemical studies of the cultured water, histopathological and oxidative enzyme activities were carried out using standard methods. The result showed presence of metals such as iron, cadmium, zinc, copper and magnesium. Presence of alkaloids, flavonoids, steroid, phenols, terpenoid, saponin and anthraquinone were detected. There was no significant difference in the water temperature and pH of the cultured water of *O. niloticus* exposed to the varying concentrations of the sawdust. These were observed to increase with increase in the concentrations of the sawdust. Dissolved oxygen was however highest in the control water and reduced with increase in the concentrations of sawdust extract exposure. Disintegrated lamella was observed in the intestine and gills, and graded necrosis, in the liver. There were inhibitions in the activities of superoxide dismutase (SOD) of both liver and kidney. Activity of malondialdehyde (MDA) however, increased with increased extract concentration. This study has therefore shown that sawdust extract could cause tissue - organ architectural distortions and stress exertion with consequent physiological imbalance of fish.

Keywords: Physiological, Histopathological, Inhibitions

1. INTRODUCTION

Sawmill industries has dotted water body banks in various parts of the country due to ease to access logs and increased demand. This huge sawmilling operation has consistently been generating wood wastes of different forms. Wood shavings and sawdust have been implicated as sources of water pollution by reducing light penetration of water and weakening the immune system of fish due to inert solids and toxic substances associated to these wastes [1].

Sawdust leaches degrade into some compounds that can be toxic to life. These are phenols and methylated phenols, benzoic acid, terpenes and triphenols [2]. Heavy metals being components of wood wastes at even low concentration can disrupt the ecological balance of the recipient

environment [3]. There is clear scientific evidence that if sawdust as wastes is improperly managed, wood residue can negatively impact the environment, contaminate and destroy fish habitat [4]. However, there is dearth of information on holistic physiological effects of sawdust on aquatic lives in Nigeria. The study was conceived to scientifically evaluate the level of impairment that may result on exposing fish to wood waste, particularly sawdust. This would help authorities concerned with waste management come up with policies to discourage direct dumping of sawdust into natural water bodies.

2. MATERIAL AND METHODS

2.1 SUBSTRATE COLLECTION AND ETHANOLIC EXTRACT PREPARATION

Sawdust was collected from Okobaba sawmill, Lagos State into clean plastic containers (20 x 12 x 10 cm³) with stoppers. Immediately after they were air dried and returned to the stoppered plastic containers from Okobaba sawmill, Lagos State. The collected sawdust was sieved using lab test shaker vibration (model GZ -200 with mesh size 7-600M) sieving machine to obtain fine homogenized particles and 0.3kg of the particles was weighed on an electrical weighing balance and soaked in 2.55L of 70% ethanol for 48 hours. The immersed sawdust was filtered using Whatmann's No. 1 filter paper to obtain a clear extract. The alcoholic sawdust extract was placed in a water bath at 80°C for 2 hours to evaporate the ethanol content completely. The stock solution was kept in a sterile plastic container and stored at 4°C. Thirty (30) ml of the sawdust extract was analyzed for its qualitative and quantitative phytochemical composition [5]. The work was carried out in the Biological Science laboratory of the Department of Biological Science, Yaba College of Technology.

2.2 PHYTOCHEMICAL SCREENING OF SAWDUST EXTRACT (QUALITATIVE)

Phytochemical tests were carried out on the alcoholic extract of the sawdust using a standard procedure to identify the constituents.

2.3 MEASUREMENT OF PHYSICO-CHEMICAL PARAMETERS OF WATER

Water was collected from the tap source and allowed to dechlorinate over night before use. The physico-chemical parameters of experimental water such as; temperature, pH, dissolved oxygen, electrical conductivity and nitrates, were measured every 4 days.

2.4 TRACE/HEAVY METALS DETERMINATION OF SAWDUST SAMPLE

A total of 20g of oven-dried sawdust was weighed using a sensitive weighing scale into separate porcelain crucibles and ashed at 950⁰c for 2hrs in a furnace and cooled. Out of the ashed samples, 2g of each was weighed into ten (10) 250ml separate beakers and digested using Aquaregia (a mixture of nitric acid and hydrochloric acid in the ratio 1:3). After which 75ml distilled water was added to each sample and boiled for 10 minutes. Each sample was then filtered into 100ml volumetric flask and allowed to cool. The volume then made up (Fe, Cd, Zn, Cu and Mg) using an Atomic Absorption Spectrophotometer [6].

2.5 FISH COLLECTION

One hundred and twenty (120) *Oreochromis niloticus* of average weights 3.0g ± 1.81 were purchased from Nigeria Institute of Oceanography and Marine Research (NIOMR), Lagos, for acute toxicity. The fingerlings were acclimated for 14days in 20L carrying capacity rectangular tanks.

2.6 ACUTE TOXICITY STUDIES

Square glass tanks (volume 4.5 litres; 15 × 15 × 15 cm³) were used as bioassay containers. In all bioassays for fish, test media was made up to two litres of water to hold 6 fingerlings per bioassay tanks in triplicates. *O. niloticus* fingerlings were placed in sawdust extract of varied concentrations (6ml/l, 7ml/l, 8ml/l and 9ml/l) [7]. The concentrations needed to kill all *O. niloticus* fish were found to be in the range of 2ml/l, 3ml/l, 4ml/l and 5ml/l. The difference in concentrations of Sawdust extract used for acute toxicity was based on range findings.

2.7 CHRONIC TOXICITY STUDIES (SUBLETHAL TOXICITY STUDIES)

Rectangular glass tanks (50cm by 36cm by 34cm) were used as bioassay containers. In all test media, 8 litres of water with 10 juveniles *O. niloticus* each were used. Different concentrations of the extract being constituents of the test media were obtained in methods adopted by [8] after calculating the 96hr LC₅₀ using the fractions 1/10, 1/100, and 1/1000. 96hr LC₅₀. Period of sublethal studies lasted 42 days, and the choice of 42 days was to establish suitable duration period for sublethal studies which is usually minimum of 21 days. All bioassays were in 3 replicates.

2.8 HISTOPATHOLOGICAL ANALYSIS OF THE TISSUES/ ORGANS (INTESTINE, GILLS AND LIVER OF *OREOCHROMIS NILOTICUS*)

Gill arch of the right side, liver, intestine and of the fishes were collected and accessioned.

Tissues and organs were then grossly examined and fixed in Bouin's fluid for 24 hours, washed in 70% ethanol and dehydrated in graded (10%, 30% and 60%) alcohol [9]. Organs were later cleared in xylene to remove excess alcohol from the tissue. The tissues and organs were thereafter impregnated and embedded in paraffin wax and allowed to solidify. Sections (5µm of thickness) were cut using rotary paraffin microtome and stained using haematoxylin and eosin. Light microscopy was later done by compound microscope and film photographed by Olympus DP – 10 Digital cameras attached to the microscope through a C-connector.

2.9 ANTIOXIDANT AND NON-ENZYMES ACTIVITY OF LIVER, GILLS AND KIDNEY OF *Oreochromis niloticus* Antioxidant enzymes activities were determined by standard methods:

2.9.1 Determination of Superoxide Dismutase (SOD) Activity

Superoxide Dismutase activity was determined by its ability to inhibit the autooxidation of epinephrine determined by the increase in absorbance at 480nm as described by [10]. The reaction mixture contained 2.95 ml 0.05 M sodium carbonate buffer pH 10.2, 0.02 ml of liver homogenate and 0.03 ml of epinephrine in 0.005 N HCl was used to initiate the reaction. The reference cuvette contained 2.95 ml buffer, 0.03 ml of substrate (epinephrine) and 0.02 ml of water.

Enzyme activity was calculated by measuring the change in absorbance at 480 nm for 5 min. $\Sigma = 4020M^{-1} cm^{-1}$ which is Molar extinction for SOD at

$$480nm. \frac{\Delta A \times V_T \times 10^6}{\Sigma \times V_s \times mg \text{ protein}}$$

where ΔA = Change in absorbance, V_T = Total volume (volume of sample reagent), Σ = Molar extinction, V_s = Volume Of sample alone.

2.9.2 Catalase Activity Determination

Catalase activity was determined according to [10]. It was assayed colorimetrically at 620nm and expressed as µmoles of H₂O₂ consumed/min/mg protein at 25°C. The reaction mixture (1.5ml) contained

1.0ml of 0.01M phosphate buffer (pH 7.0), 0.1ml of tissue homogenate and 0.4ml of 2M H₂O₂. The reaction was stopped by the addition of 2.0ml of dichromate-acetic acid reagent (5% potassium dichromate and glacial acetic acid

were mixed in 1:3 ratio). $\Sigma = 40M^{-1} \text{ cm}^{-1}$

$$\frac{\Delta_A \times V_T \times 10^6}{\Sigma \times V_s \times mg \text{ protein}}$$

2.9.3 Reduced Glutathione Determination

The reduced glutathione (GSH) content of liver, kidney and gill tissues as non-protein sulphhydryls was estimated. Ten percent TCA was added to the homogenate and centrifuged. One (1ml) of supernatant was treated with 0.5ml of Ellmans reagent (19.8mg of 5,5dithiobisnitro benzoic acid (DTNB) in 100ml of 0.1% sodium nitrate) and 3.0ml of phosphate buffer (0.2M, pH 8.0). The absorbance was read at 412nm and $\Sigma =$ Molar extinction co-efficient for GSH =

$$1.34 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}.$$

$$\frac{\Delta_A \times V_T \times 10^6}{\Sigma \times V_s \times mg \text{ protein}}$$

2.9.4 Determination of Malondialdehyde (MDA)

Malondialdehyde (MDA), an index of lipid peroxidation was determined using the method of [11]. One ml of the supernatant was added to 2 ml of (1:1:1 ratio) TCA-TBA-HCl reagent (thiobarbituric acid 0.37%, 0.24N HCl and 15% TCA) tricarboxylic acid- thiobarbituric acidhydrochloric acid reagent boiled at 100°C for 15 minutes and allowed to cool. Flocculent materials were removed by centrifuging at 3000 rpm (revolutions per minute) for 10 minutes. The supernatant was removed and the absorbance read at 532 nm against a blank. MDA was calculated using the molar extinction coefficient for MDATBA- complex of $1.56 \times 10^5 \text{ M}^{-1} \text{ CM}^{-1}$ [11].

2.10 STATISTICAL ANALYSIS

Data were analyzed using the Statistical Package for Social Sciences (SPSS) version 20 [12]. Mean values were compared using Analysis of Variance (ANOVA). Post hoc test was done using the student-Newman-Keuls (SNK). Means were presented as Mean \pm Standard deviation while P-value was set at 0.05 level of significance.

3. RESULTS

3.1 PHYTOCHEMICAL COMPOSITION

The result of qualitative phytochemical screening of the ethanol extract of sawdust revealed the presence of alkaloids, flavonoid, steroid, phlobatanin, phenols, terpenoid, anthraquinone and saponin (Table 1).

Table 1: Qualitative phytochemical composition of the sawdust extract obtained from Okobaba Sawmill Industry, Ebute Metta Mainland, Lagos State

Phytochemical constituents	Status
Alkaloid	Positive
Flavanoid	Positive
Steroid	Positive
Philobatanin	Positive
Terpenoid	Positive

Anthraquinone	Positive
Saponin	Positive

3.2 Trace Metals composition

Five metals were detected in the sawdust extract used (Table 2). These metals include iron, cadmium, zinc, copper and magnesium. Of all these, iron was considerably highest in concentration. This was followed by zinc respectively. However, copper and cadmium had the lowest concentrations in the sawdust extract.

Table 2: Metals composition of the sawdust extract from Okobaba Sawmill Industry, Ebute Metta Mainland, Lagos State

METALS	QUANTITY (mg/g)
Iron (Fe)	6.0492
Cadmium (Cd)	0.0136
Zinc (Zn)	2.2303
Copper (Cu)	0.010
Magnesium (Mg)	0.3349

Table 3: Physicochemical parameters of culture water for *Oreochromis niloticus* exposed to different concentrations of sawdust extract

	Control	0.003 ml/l	0.03 ml/l	0.3 ml/l
Temperature (°C)	28.60±0.24 ^a	28.40±0.65 ^a	28.00±0.22 ^a	27.80±0.40 ^a
Conductivity (S/m)	600±3.25 ^d	800±0.82 ^c	970±0.14 ^b	1032±0.6 ^a
TSS (ppm)	86±0.83 ^d	128±1.03 ^c	382±0.02 ^b	420±1.25 ^a
BOD (ppm)	30±1.18 ^d	36±0.01 ^c	42±1.36 ^b	58±0.34 ^a
DO (ppm)	5.80±0.71 ^a	5.20±1.60 ^a	4.00±1.28 ^a	3.60±1.10 ^a
Nitrate (ppm)	10.50±0.13 ^d	18.20±1.87 ^c	20.50±1.28 ^b	23.40±0.18 ^a
Phosphate (ppm)	10.20±0.15 ^d	16.33±1.88 ^c	18.35±0.99 ^b	20.30±0.60 ^a
Turbidity (Ftu)	70±1.35 ^d	150±0.35 ^c	171±1.03 ^b	243±1.70 ^a

pH 7.80±0.96^a 7.30±0.50^a 6.90±1.40^a 6.10±0.61^a

^{abcd}Means (±Standard error of mean) in the same row having similar superscripts were not significantly different at P >0.05

3.3 PHYSICO-CHEMICAL PARAMETERS OF CULTURE WATER FOR *OREOCHROMIS NILOTICUS*

The physico-chemical parameters of the experimental treatment water are shown in Table 3. Temperature was relatively stable in all treatment water with the highest degree Celsius of 28 recorded at 0.3% extract concentration. P^H (6.1) and D.O (3.6ppm) were recorded at 30% extract concentration compared to values obtained at 3.0% and 0.3%. The highest conductivity was recorded at 0.3 ml/l (1032 ± 0.6 S/m) and this was significantly higher (p<0.05) than conductivity recorded at 0.003 ml/l and 0.03 ml/l extract concentrations. The Biological Oxygen Demand increased with increased concentration. The highest BOD (58 ± 0.34 ppm) was recorded at 0.3 ml/l extract concentration compared to BOD recorded at 0.003 ml/l and 0.03 ml/l extract concentrations. Turbidity was significantly higher (243 ± 1.7 ftu) compared to those recorded at 0.003 ml/l (150 ± 0.35 ftu) and 0.03 ml/l (171 ± 1.03 ftu).

3.4 MEAN WEIGHT OF *Oreochromis niloticus* EXPOSED TO DIFFERENT CONCENTRATIONS OF SAWDUST EXTRACT

Weight increase was also recorded in *O. niloticus* administered with the varying concentrations of sawdust extracts (0.003 ml/l, 0.03 ml/l and 0.3 ml/l) and the control over the six weeks study period (Figure 3). A gradual weight increase was observed between initial and week 2 in all fish exposed to varying concentrations of the extract. However, between week 2 and 3, there was weight gain in fish exposed to 0.03ml/l and 0.3ml/l. There was no significant difference (p>0.05) recorded in the mean weight gain of the control and those administered with 0.3ml/l of the saw dust extract (Table 4). The weight gain was significantly higher (p < 0.05) than those of the *O. niloticus* groups administered with 0.003 ml/l and 0.03 ml/l sawdust extracts respectively.




Figure 1: Mean weekly weight of *Oreochromis niloticus* on ethanolic extract of sawdust extract

Table 4: Mean weight gain of *Oreochromis niloticus* administered with varying concentrations of sawdust extract

Treatment	Live Weight		
	Initial	Final	Gain
Control	6.41 ± 1.30 ^a	19.40 ± 0.63 ^a	12.99±0.58 ^a
0.003 ml/l	6.39 ± 1.84 ^a	18.20 ± 0.60 ^b	11.81±0.01 ^b
0.03 ml/l	6.45 ± 0.58 ^a	16.53 ± 0.19 ^c	10.08±0.01 ^c
0.3 ml/l	6.40 ± 0.08 ^a	19.69 ± 0.02 ^a	13.29±0.06 ^a

^{abcd}Means (±Standard error of mean) in the same column having similar superscripts were not significantly different at P > 0.05

3.5 HISTOPATHOLOGY OF SOME BODY ORGANS EXPOSED TO DIFFERENT CONCENTRATIONS OF SAWDUST EXTRACT

3.5.1 Section of the Intestine of *Oreochromis niloticus* exposed to different concentrations of sawdust extract

The intestinal tissues of the control *O. niloticus* showed normal architecture with abundant mucosal folds, underlying submucosa folds and muscularis devoid of inflammatory cell infiltrates (Plate 1). On the other hand, the intestine of the *O. niloticus* groups administered with the varying concentrations of the sawdust extract showed disintegrating columnar epithelium with reduced mucosa folds thickness and gastric intestinal atrophy.




Plate 1: Section of experimental *Oreochromis niloticus* intestine; A = Control; B = 0.3 ml/l sawdust extract; C = 0.03 ml/l sawdust extract; D = 0.003 ml/l sawdust extract; N = Normal architecture; GIA = Gastric intestinal atrophy. (Mg = X40).

3.5.2 Section of the Gills of *Oreochromis niloticus* exposed to different concentrations of sawdust extract

The Histology of gills of *O. niloticus* administered with the varying concentrations of the sawdust extract is shown in Plate 2. Disintegrated columnar epithelium, increased lumen, increase in length of longitudinal muscle layer and increased mucosa were observed in the groups administered with the different concentrations of the sawdust extract. However, normal gill architecture consisting of the dense columnar epithelium, mucosa and submucosa lining, epithelial cells, longitudinal muscle layer and thin columnar cells was observed in the control group.




Plate 2: Section of experimental *Oreochromis niloticus* gill; A = Control; B = 0.3 ml/l sawdust extract; C = 0.03 ml/l sawdust extract; D = 0.003 ml/l sawdust extract; N = Normal architecture; MLN and LN = Lamellar necrosis; FP = Focal epithelial proliferations. (Mg = X40).

3.5.3 Section of the Liver of *Oreochromis niloticus* exposed to different concentrations of sawdust extract

The liver of the control *O. niloticus* showed normal architecture (Plate 3). Extensive proliferation of fibrous tissue (Hepathic fibrosis) with mild inflammation was however observed in the liver of those administered with the varying concentrations of the sawdust extract.




Plate 3: Section of experimental *Oreochromis niloticus* liver; A = Control; B = 0.3 ml/l sawdust extract; C = 0.03 ml/l sawdust extract; D = 0.003 ml/l sawdust extract; NO = Normal architecture; HF = Hepatic fibrosis. (Mg = X40).

3.6 Activities of Antioxidant enzymes level of the liver, kidney and gills of *Oreochromis niloticus* exposed to varying concentrations of sawdust extract

The activities of superoxide dismutase (SOD), reduced glutathione (GSH) and catalase in the liver, kidney and gills of the experimental *Oreochromis niloticus* is presented in table 5. In the liver, GSH activity was recorded highest at 0.003ml/l among the exposed concentrations with a value of 13.50nm/mg pro. GSH activity in the kidney among exposed concentrations of sawdust was highest at 0.003ml/l with a value of 10.12nm/mg pro. In the gills, the highest value (11.30nm/mg pro) of GSH activity among exposed concentrations was at 0.003ml/l. The liver recorded the highest value of GSH activity (13.50nm/mg pro) among the three organs exposed to the sawdust extract.

SOD activity in the liver was recorded highest at 0.003ml/l among the concentrations exposed to sawdust extract with a value of 110.11nm/mg pro. In the kidney, the highest SOD activity was recorded at the lowest concentration with a value of 98.15nm/mg pro. The gills recorded the highest value (122nm/mg pro) of SOD activity at the lowest concentration among the concentrations exposed to the extract. Among these three organs exposed to the extract, the highest value (122nm/mg pro) of SOD activity was in the gills at 0.003ml/l concentration.

In the liver, CAT activity was recorded highest at 0.003ml/l among the exposed concentrations with a value of 604.88nm/mg pro. CAT activity in the kidney among exposed concentrations of sawdust was highest at 0.003ml/l with a value of 466.55nm/mg pro. In the gills, the highest value (495.96nm/mg pro) of CAT among exposed concentrations was at 0.003ml/l. The liver recorded the highest value of CAT activity (604.88nm/mg pro) among the three organs exposed to the sawdust extract. There was no significant difference ($p > 0.05$) in the activities of reduced glutathione (GSH) and superoxide dismutase (SOD) recorded in the liver of the control *O. niloticus* and those administered with 0.003 ml/l of the sawdust extract (Table 5). These were significantly higher ($p < 0.05$) than those administered with 0.03 ml/l and 0.3 ml/l concentrations of the sawdust extract. Catalase activity of the liver was also significantly

higher in the control group and observed to significantly reduce with increasing concentration of the sawdust extract administration.

In the kidney of the experimental *O. niloticus*, GSH and catalase activities were significantly higher in the control group. These were observed to significantly reduce in the experimental *O. niloticus* with increase in the concentration of the sawdust extract administration. Kidney SOD activity was not significantly different ($p > 0.05$) between the control group and those administered with 0.003 ml/l and 0.03 ml/l of the sawdust extract. Kidney SOD activity was however significantly lower ($p < 0.05$) in the *O. niloticus* group administered with 0.3 ml/l of sawdust extract. Similarly, GSH and catalase activities of the gills were significantly higher ($p < 0.05$) in the control group those administered with the varying concentrations of the sawdust extract. These were however observed to reduce with increase in the concentration of the sawdust extract administration. On the other hand, SOD activity of the gill was significantly lower in the *O. niloticus* group administered with 0.3 ml/l of the sawdust extract. SOD activity was however not significantly different ($p > 0.05$) between the control group and those administered with 0.003 ml/l and 0.03 ml/l of the sawdust extract.

Table 5: Antioxidant enzymes activity of *O. niloticus* administered with varying concentrations of sawdust extract

	Conc. (ml/l)	GSH (nm/mg pro)	SOD(nm/mg pro)	Catalase(nm/mg pro)
Liver	Control	16.14±0.75 ^a	114.03±1.44 ^a	615.02±0.31 ^a
	0.003	13.50±0.43 ^a	110.11±0.42 ^a	604.88±0.29 ^b
	0.03	6.82±0.91 ^b	104.00±1.70 ^b	594.06±0.01 ^c
	0.3	3.81±0.94 ^b	99.60±1.06 ^b	573.17±0.25 ^d
Kidney	Control	16.80±0.68 ^a	103.41±1.86 ^a	501.70±1.85 ^a
	0.003	10.12±1.42 ^b	98.15±1.15 ^a	466.55±0.31 ^b
	0.03	4.01±0.34 ^c	96.71±1.93 ^a	457.62±0.17 ^c
	0.3	3.21±0.07 ^c	82.31±1.19 ^b	432.11±0.1 ^d
Gill	Control	18.32±1.08 ^a	125.00±1.36 ^a	506.33±3.18 ^a
	0.003	11.30±0.22 ^b	122.00±0.50 ^a	495.96±0.24 ^b
	0.03	3.86±0.13 ^c	120.63±0.04 ^a	455.01±2.83 ^c
	0.3	3.01±0.55 ^c	116.3±0.48 ^b	461.63±0.93 ^c

^{abcd}Means (±Standard error of mean) in the same column for each of the organs having similar superscripts were not significantly different at $P > 0.05$

3.6.1 Lipid peroxidation in the liver, kidney and gills of *Oreochromis niloticus* cultured on varying concentrations of sawdust extract

The level of lipid peroxidation recorded in the liver, kidney and gills of the *O. niloticus* were not significantly different ($p > 0.05$) between the control group and those administered with the varying concentrations of the sawdust extract (Figure 2). Lipid peroxidation was also observed to be lowest in the control group and increase with increase in the concentration of the sawdust extract. Lipid peroxidation

was therefore highest in kidney, followed by liver and gills of the *O. niloticus* at 0.3 ml/l concentration of the sawdust extract.




Figure 2: Levels of lipid peroxidation (Malondialdehyde – MDA) in *O. niloticus* administered with varying concentrations of sawdust extract; ^{abcd}Means in the same cluster having similar superscripts are not significantly different at $p < 0.05$; Error bars represents Standard error of mean.

3.7 DISCUSSION

The study has demonstrated decreased dissolved oxygen with increased sawdust extract concentration compared to the control. Lower level of D.O in the culture water of *O. niloticus* exposed to the varying concentrations of the sawdust extract could result in suffocation and mortality of fish species. Heavy metals inherent in sawdust could have contributed to morphological, physiological and histopathological aberrations and disruptions of fish as evident in this study. Histological responses have been pointedly reported as valuable markers of toxicology. Histopathological changes such as increased cytoplasmic vacuolation, definitive necrosis and hepatocellular alteration were observed in the liver of *O. niloticus*. This agreed with the findings of [13] [14] [15].

Increased vacuolation of the hepatocytes is a signal of degenerative process that suggests metabolic damage. This compromised integrity status of the liver of *O. niloticus* reflects the level of degeneration. The intestine of the exposed fish showed many changes and the pathological changes are gradual abnormality and elongation of the intestinal wall with disintegrating columnar epithelium and increased human with increased mucosa and submucosa.

Disintegrated lamella was equally observed in the intestine, this agreed with [16] [17]. The severity of gill damage as observed depends on the concentration of extract on time of exposure. Due to the peculiar morphology of the gill filaments and lamella *O. niloticus* the rate and degree of degeneration was well defined. Impairment of gills causes destruction of oxygen - carbon dioxide exchange [18]. Damage of the gills can also be associated to the presence of phenolic compounds in the sawdust.

The activity of antioxidant enzymes may be enhanced or inhibited under chemical stress depending on the intensity and the duration of the stress applied, as well as, the susceptibility of the exposed

species. Under undisturbed conditions, a balance exists between the production of Reduction Oxygen Species (ROS) and oxidation processes. The inhibition in the activities of GSH, SOD and CAT in the livers and kidneys of the fish (*Oreochromis niloticus*) reported in the laboratory studies after 42 days exposure may be a response to oxidative stress caused by some phytochemicals inherent in the extract and the heavy metals in the superoxide radicals. Phytochemicals such as alkaloids have been implicated with stress in fish [19]. The inhibition of the enzyme SOD by the test chemicals will therefore lead to increased oxidative stress in the liver and the kidney tissues. There is a nexus between the activities of SOD and CAT. Inhibition of the enzyme of SOD will result in decline in the mechanism of CAT due to reduced H₂O₂ production from SOD. This was proven in this study as there was a significant reduction in CAT activity in species examined. GSH is a massive biomarker of oxidative stress. It has the ability to bind with electrophilic xenobiotics and converting them into lose excretable product. Decreased level of GSH was observed in the liver, gill and kidney of fish species examined in this study after exposure to sawdust extract. The anthraquinone content of the extract could also be responsible for the inhibition of SOD, CAT and GSH [20].

The inhibition of antioxidant defense in this study correlates with an increase in oxidative damage. This was evident with increased MDA levels, particularly in the liver of exposed fish. Significant increase in MDA in the liver, kidney and gills of *Oreochromis niloticus* agreed with the findings of [21] that reported MDA increase in the tissues of fish exposed to varied concentrations of hydrocarbons. The increased MDA indicates that ROS formed may be associated with the metabolism of sawdust leading to peroxidation of membrane lipids of the respective organs.

4. CONCLUSION

The results from this investigation showed that sawdust extract of different wood origins exert stress inducing effect on the general physiology of the animal

(*Oreochromis niloticus*) though at varied degree during exposure period. This assertion was by induction and inhibition of enzymes and antioxidants among other parameters investigated.

The high mortality, particularly during acute exposure over period of both acute and sublethal exposure expounded the toxic nature of the extract. Physiological aberrations depended on the physicochemical features of test water, extract phytochemicals, hydrocarbon load (in-tool lubricant), rate and period of exposure. Organs such as liver, intestine and gills were severally impaired. From the findings reported, it can be concluded that exposure of *O. niloticus* to varied concentrations of sawdust extract exerts enormous stress due to physiological imbalance.

REFERENCES

1. FAO/WHO (1991). Protein Quality Evaluation. Report of Joint FAO/WHO Expert Consultation. *FAO Food and Nutrition Paper* 51, FAO/WHO, Rome, Italy. 247pp.
2. Peters, G.B., H.J. Dawson, B.F. Hruttfjord, and R.R. Whitney. (1976). Aqueous leachate from western red cedar: Effects on some aquatic organisms. *Journal of the Fisheries Research Board of Canada* 33:2703-2709.
3. Farombi, E.O. O.A. Adelowo, Y.R. Ajimoko., (2007). Biomarkers of oxidative stress and heavy metal levels as indicators of environmental pollution in African Cat fish (*Clarias gariepinus*) from Nigeria Ogun river. *Int. J. Environ. Res. Public Health*. 4(2): 158165.
4. Arimoro, F.O., Ikomi, R.B. and OsalorC.E. (2006). The Impact of Sawmill Wood Wastes on the Water Quality and Fish Communities of Benin River, Niger Delta Area, Nigeria. *World Journal of Zoology*, 1(2): 94-102.

5. Anita, P. and Shalini, T. (2014). Concept of standardization, extraction and prephytochemical screen strategies for herbal drug. *Journal of Pharmacognosy and Phytochemistry*, 2(5): 115-119.
6. Radulescu, C., Dulama, I.D., Stih, C., Ionita, I., Chilian, A., and Necula, C. (2014). Determination of heavy metal levels in water and therapeutic mud by atomic absorption spectrometry. *Romania Journal of Physics*, 59(9-10): 1057-1066
7. Solbe J. F. (1995); Fresh water in Handbook of Ecotoxicology (Edited by Peter Collius) Blackwell Science. 683pp.
8. Otitolaju, A.A. (2000). Joint action toxicity of Heavy metals and their bioaccumulation by benthic animals of the Lagos lagoon. *Phd thesis*. University of Lagos. 229pp.
9. Egonwan, R.I. (2007). An ultrastructural study of the seminal vesicle of the hermaphadite dust of the land snail *Limicolaria flammea* (muller) (Pulmonata: Acatinidae). *Pakistan Journal of Biological Sciences*. 10(11): 1835–1839.
10. Zhang, J., Chen, R., Yu, Z. and Xue, L. (2017). Superoxide Dismutase (SOD) and Catalase (CAT) Activity Assay Protocols for *Caenorhabditis elegans*. *Bio-protocol* 7(16): e2505.
11. Mateos, R., Lecumberri, E., Ramos, S., Goya, L. and Bravo L.(2005). Determination of malondialdehyde (MDA) by high-performance liquid chromatography in serum and liver as a biomarker for oxidative stress; Application to a rat model for hypercholesterolemia and evaluation of the effect of diets rich in phenolic antioxidants from fruits. *Journal of Chromatography B Analytical Technology and Biomedical Life Science*, 827(1):76-82.
12. IBM Corporation (2011) IBM SPSS statistics for Windows, version 20.0. Armonk, NY: IBM Corp.
13. Camargo, M. M. P. and C. B. R. Martinez (2007). Histopathology of gills, kidney and liver of a Neotropical fish caged in an urban stream. *Neotropical Ichthyology*, 5(3): 327-336.
14. Eisenhut, M. (2006). Changes in ion transport in inflammatory disease. *Journal of inflammation (London, England)*, 3:5.
15. Wolf, J.C. and Wheel, J.R.(2018).A critical review of histopathological findings associated with endocrine and nonendocrine hepatic toxicity in fish models. *Aquatic Toxicology*, 197:60-78.
16. Singh, S. and Mehrotra, A. (1999). Histopathological changes induced by carbaryl in the intestine of fresh water fish *Nandus nandus*.*Journal of Ecotoxicology and Environment Monitoring*, 11(2):129-132.
17. Kaoud, H. A., Zaki, M.M., EL-Dahshan, A.R., Saeid, S. and El Zorba, H.Y., (2011). Amelioration the toxic effects of cadmium-exposure in nile tilapia (*Oreochromis niloticus*) by using Lemnagibba L. *Life Science Journal*, 8: 185-195.
18. Dash, S., Das, S.K., Samal, J., and Thatoi, H.N. (2018). Epidermal mucus, a major determinant in fish health: a review. *Iranian journal of veterinary research*, 19(2), 72–81.
19. Omitoyin, B.O., Ogunsami, A.O. and Adesina, B.T. (2006). Studies on acute toxicity of Piscicidal plant extracts (*Tetrapleura tetraptera*) on tilapia (*Sarotherodon gailiaeus*) fingerlings. *Tropical Journal of Animal Science*, 2(2):191-197.
20. Liu, B., Ge, X., Xie, J., Xu, P., He, Y., Cui, Y., Ming, J., Zhou, O. and Pan, L. (2012). Effects of Anthraquinone extract from *Rheum officinale* Bail on the physiological responses and HSP 70 gene expression of *Megalobrama amblycephala* under *Aeromona hydrophilia* infection, *Fish Shell Immunology*, 1: 1-7.
21. Otitolaju A and Olagoke, O. (2011). Lipid peroxidation and antioxidant defense enzymes in *Clarias gariepinus* as useful biomarkers for monitoring exposure to polycyclic aromatic hydrocarbons. *Environment Monitoring Assessment*, 182: 205-213