

Species richness, diversity and distribution of phytoplankton in fertilised ponds of the western highlands agro-ecological zone of Cameroon

ABSTRACT

In developing countries, advanced aquaculture techniques are still at the embryonic stage. Semi-intensive fish farming system, based on the use of fertilisers is the most currently practiced methods in ponds. Thus, this study was conducted from January to December 2021 in the ponds of the West Cameroon Region and at the Ichthyology and Applied Hydrobiology Research Unit of the University of Dschang with the aim of contributing to a better understanding of species richness and distribution of phytoplankton populations. For this purpose, a total of fifteen ponds including 03 in Bamendou fertilised with cow dung, four in Batié including two fertilised with pig manure and two with wheat bran, one in Dschang fertilised with chicken manure, four in Fokoué including two fertilised with pig manure and two unfertilised, and three in Foubot, fertilised with cow dung, were assessed. These ponds were chosen based on the availability of fish farmers and the type of fertiliser administered. Together with water physicochemical properties measurement, phytoplankton sampling was done between 6 a.m. and 8 a.m. on a monthly basis. The results showed that phytoplankton species and genera richness was higher in unfertilised ponds in Fokoué ponds and in animal feed enriched ponds in Batié, respectively. However, the highest family richness was obtained in Dschang receiving chicken manure with a percentage of 88.8% of the total family richness. The species *Microcystis aeruginosa* exhibited the highest species frequency (57%) out of all the species recorded in the ponds of Batié fertilised with pig manure. The lowest Shannon-Weaver diversity indices and Pielou evenness indices were recorded in the Fokoué ponds independent of the type of fertiliser, and in the Dschang ponds receiving chicken manure. The outputs of this work are better proposals for the production of phytophagous or bulldozer fish, such as carp, in relation to the site and the type of fertilisers in the Western highlands agroecological zone of Cameroon in particular and the world in general.

Keywords: Phytoplankton, Fertilised, Ponds, Species Richness, Fish, Diversity Indices

1. INTRODUCTION

In many sub-Saharan African countries, food shortage remain a reality [1, 2]. According to the Ministry of Agriculture and Rural Development (MINADER), in Cameroon, 2 865 906 people representing 11% of the population suffer from acute food insecurity. The chief causes of malnutrition comprise poverty, lack of access to drinking water, sanitation, medical care, education and food [3]. Undernutrition is mainly due to a lack of animal protein, which is under-consumed by the most helpless populations. Among fishery resources, fish offers 51% of protein to the human diet [4]. Under such circumstances,

man must cultivate fish to unravel the problem of food insecurity, imports and capture fishing, which is declining over the years. Fish farming in ponds could thus constitute a way of enlightening production given the complexity of the pond food webs. Phytoplankton forms the support of the pelagic food chain [5] and is responsible for an essential part of the primary production in aquatic ecosystems, especially fish ponds. Phytoplankton refers to all the plant micro-organisms suspended in water, capable of producing their own organic substance via photosynthesis, from solar energy, water, oxygen and nutrient salts [6]. Apart from direct feeding of the fish with exogenous food, fish production in ponds depends on the planktonic richness which is itself a function of the influences of mineral and organic fertilizers. Knowledge of the production proficiency of a fish pond is essential to avoid the appearance of trophic deadlocks. Several studies have been carried out on the biogeography of phytoplankton in Europe and America [7-11] but phytoplankton remains poorly documented in Africa [12]. In Cameroon, some work has been done on faunal surveys and the dynamics of phytoplankton in the Municipal Lake of Yaoundé [13] as well as phytoplankton characterization in a floodplain and in a river in relation to nutrient status [14, 15]. At the current state of knowledge, only the work of Nana et al. [16] focused on the distribution and diversity of phytoplankton in fertilised ponds. Yet, the nature of phytoplankton species must allow their proper integration into the food chain as fish rely on them as food source. Therefore, it is a necessity to intensify the study of planktonic algae in order to access the biodiversity of lentic ecosystems and to clarify the ecology of the species present there. This study was aimed at contributing to a better knowledge of the bio-ecology of fish ponds through a characterization of phytoplankton population in the ponds of the western highlands agro-ecological zone of Cameroon. More specifically, this research evaluated the effect of the sampling site and the type of fertiliser on the species richness and distribution of phyla, families, genera and species of phytoplankton as well as on species diversity.

2. MATERIAL AND METHODS

2.1. Study sites

The work was carried out from January to December 2021 in the ponds of the West Cameroon and at the Ichthyology and Applied Hydrobiology Research Unit of the University of Dschang. The geoclimatic features of the study area are as follows. North Latitude: 5° and 6°; East Longitude: 10° and 11°, average altitude: 1400 m, average annual rainfall: 1500 mm, temperature: 14-25°C, dry season: mid-November to mid-March, rainy season: mid-March to mid- November.

2.2. Characteristics of phytoplankton collection ponds

Three sites were chosen, far enough apart to be considered sufficiently representative of the spatial variation of the ecosystem. Using a questionnaire and direct observations, information on the characteristics of the ponds and their organisation by fish farmers was collected. For this purpose, a total of 15 ponds were selected depending on the type of fertilizer administered. The choice of these ponds was also based on approachability. The characteristics of the study ponds are summarized in Table 1.

Table 1: Chief characteristics of the study ponds

Sites	Geographical Coordinates	Altitude (m)	Types de Fertilisers	Composition of fertilisers
Batié	LN : 5°17'-5°18' and LE : 10°17'- 10°19'	1 500	Wheat bran	N : 0,01 ; P : 0,02 ; N/P : 0,5
			Wheat bran Pig manure Pig manure	N : 0,59 ;P : 0,60 ; N/P : 0,93
Bamendou	LN : 5°22'-5°28' and LE : 10°52'-10°54'	1 400	CBV	N : 0,45 ; P : 0,25 ; N/P :1,8
			CBV	
			CBV	

Fokoue	LN : 5°34'-5°36' and LE : 10°14'-10°17'	1 276	Pig manure Pig manure Not fertilized	N : 0,5 ; P : 0,45 ; N/P : 1,11
Dschang	LN : 5°44'-5°48' and LE : 9°85'-10°05'	1 391	Not fertilized Chicken droppings	N : 3 ; P : 2 ; N/P : 1,5
Foumbot	LN : 5° 20' to 5° 22' and LE : 10° 17' to 10° 21'	1 120	CBV CBV CBV	N : 0,3 ; P : 0,15 ; N/P : 2

CBV: Cow dung compost box; N: Nitrogen; P: Phosphorous; N/P: Nitrogen-phosphorous ratio; LN: Latitude (North); LE: Longitude (East)

2.3. Conduct of the study and data gathering

2.3.1. Determination of water physicochemical characteristics

Together with the collection of phytoplankton, the physicochemical characteristics of the water were measured monthly according to the techniques advocated by Agadjihouédé et al. [17]. Transparency, depth, pH, dissolved oxygen, temperature and conductivity of the water were analysed *in situ* respectively using a Secchi disk, a water level gauge, a pH- meter, a multimeter and a conductivity meter. A 350 ml water sample was also taken in each pond and transferred to the laboratory in a refrigerated enclosure for the determination of nutrient (nitrates, ammoniacal nitrogen, nitrites and phosphates) using a HACH brand spectrophotometer DR/2800™ following APHA [18]. The non-ionized ammonia concentration was inferred from that of the ammoniacal nitrogen according to Equation 1 [19].

$$N - NH_3 = \frac{N - NH_4^+}{1 + 10^{10 - pH - 0.03T^{\circ}C}} \quad (1)$$

2.3.2. Phytoplankton sampling

Phytoplankton sampling was done monthly between 6 and 8 am according to the techniques recommended by Agadjihouédé et al. [17]. For this point, 1 litre of water was sampled at 20 points scattered on the entire area of the pond and at a depth of 30 cm. The total volume of water collected (i.e. 20 L) was filtered through plankton net with a 40 µm mesh opening according to Grogga [6]. After obtaining the filtrate, a volume of 250mL of phytoplankton concentrate was introduced into well-labelled containers and fixed with 5% formalin in the proportions of ¾ of the sample and ¼ of formalin as recommended by Nguetsop et al. [20] for quantitative and qualitative examinations.

2.3.3. Qualitative and quantitative analysis of phytoplankton

After homogenization of the phytoplankton concentrate in the laboratory, a 10 µL sample was taken using a micropipette and mounted between slide and cover slip for qualitative examination of phytoplankton species. For each sample, three slides were prepared to ensure the reproducibility [20]. The identification of the species was made with a 40X objective of an optical microscope following vertical transects chosen at random using plates and identification keys [21-25].

Quantitative analysis was executed using a ZEISS 47 12 02 inverted microscope with 40X objective. For this purpose, after homogenization of the filtrate, three sub-samples of 10 mL were taken using a pipette and each dropped into a settling dish. The set-up was allowed for ten to fifteen minutes for sedimentation of the phytoplankton organisms [26]. Counting was done in six fields taken at random

from the counting chamber. Finally, the average number of phytoplankton organisms was evaluated from the total number of fields per dish and reported as number of cells per unit volume of water.

2.4. Data processing and analysis

Following a normality test, two-way ANOVA was used to test the influences of sites and types of fertilisers on phytoplankton community. The Pearson correlation test was used to determine the relationship between water physicochemical characteristics and phytoplankton density of groups. For these analyses, the statistical software SPSS Version 20.0 was used.

Phytoplankton density was calculated using the following Equation 2:

$$D = (n \times v) / V \quad (2)$$

Where:

- D is the density (expressed in individuals per litre)
- v is the total volume of the sample analysed (mL)
- V is the volume of water filtered in the field (L)

The results of phytoplankton densities gotten were used in the calculations of various indices to characterize the composition and the evolution of phytoplankton communities, namely the Shannon Diversity Index and the Evenness Index.

Shannon Diversity Index

The Shannon Diversity Index [27] was computed according to Equation 3.

$$H' = - \sum_{i=1}^S \frac{n_i}{N} \text{Log}_2 \frac{n_i}{N} \quad (3)$$

With:

- H': Shannon Diversity Index
- n_i : Number of individuals of species i
- S: total number of species or taxonomic richness
- N: the total number of individuals of all species found (N)

The Shannon index has the ability to take into account species abundance and it is sensitive to species with low frequencies [28]; this index can be interpreted as follows [29]:

- $H' > 4$: clean water
- $3 \leq H' \leq 4$: mildly polluted water
- $2 \leq H' < 3$: moderately polluted water
- $H' < 2$: heavily polluted water

The Pielou Evenness Index

The Pielou Evenness Index [30] was calculated according to Equation 4. The Pielou Index varies between 0 (one species dominates) and 1 (all the species tend to have the same abundance).

$$E = \frac{H'}{\text{Log}_2 S} \quad (4)$$

With:

- E is the Pielou Index
- H' is the Shannon diversity index

- S is the total number of species or taxonomic richness

Species frequency was calculated using the formula:

$$F = (Pa/P) \times 100 \quad (5)$$

With:

- Pa = total number of samples containing the species considered
- P = total number of samples assessed

3. RESULTS

3.1. Richness and distribution of phyla, families, genus and species of phytoplankton depending on the site and type of fertilizer

The richness and distribution of phytoplankton species, genera and families according to site and type of fertilizer is summarised in Table 2.

Table 2: Effect of site and type of fertiliser on the richness and distribution of phyla, families, and genus and phytoplankton species

Phytoplankton species	Sites and types of fertilisers						
	Bamendou Cow dung	Pig manure	Batié Wheat bran	Dschang Chicken dung	Fokoué Pig manure	Not fertilised	Foumbot Cow dung
Cyanophyta							
Merismopediaceae							
Merismopedia							
<i>Merismopedia convoluta</i>	+	-	-	-	-	-	+
<i>Merismopedia glauca</i>	-	+	+	-	-	-	-
<i>Merismopedia tunnissina</i>	-	+	-	-	-	-	-
Chroococcaceae							
Microcystis							
<i>Microcystis aeruginosa</i>	-	+	+	+	+	+	+
<i>Microcystis densa</i>	-	+	+	+	-	-	+
<i>Microcystis holsatica</i>	-	+	+	+	-	-	+
<i>Microcystis incerta</i>	-	+	+	+	+	+	+
<i>Microcystis robusta</i>	-	+	+	+	+	+	+
Nostocaceae							
Nostoc							
<i>Nostoc parmelooides</i>	+	-	-	-	-	-	-
Oscillatoriaceae							
Oscillatoria							
<i>Oscillatoria amphibia</i>	-	+	+	-	-	-	-
<i>Oscillatoria chlorina</i>	-	+	+	-	-	-	-
<i>Oscillatoria granulata</i>	+	+	+	-	-	-	-
<i>Oscillatoria laxa</i>	-	+	+	-	-	-	-
<i>Oscillatoria teribriformis</i>	+	-	-	-	-	-	-
Gomphosphaeriaceae							
Gomphosphaeria							
<i>Gomphosphaeria naegeliana</i>	-	-	+	-	-	-	+
Chlorophyta							
Chlorophyceae							
Actinastrum							
<i>Actinastrum aciculare</i>	+	+	+	-	-	-	+
Ankistrodesmus							
<i>Ankistrodesmus fusiformis</i>	-	-	-	-	+	+	-
Chlamydomonas							
<i>Chlamydomonas conica</i>	-	+	+	-	-	-	-

<i>Chlamydomonas epiphytica</i>	+	-	-	-	-	-	+
<i>Chlamydomonas globosa</i>	+	+	+	+	+	+	+
<i>Chlamydomonas muriella</i>	+	-	-	-	-	-	-
Monoraphidium							
<i>Monoraphidium braunii</i>	-	-	-	+	-	-	+
<i>Monoraphidium convolutum</i>	-	-	-	-	-	-	-
<i>Monoraphidium contortum</i>	-	+	+	-	-	-	-
Tetraedon							
<i>Tetraedon minimum</i>	-	+	+	-	-	-	-
Hydrodictyaceae							
Kirchneriella							
<i>Kirchneriella lunaris</i>	+	+	-	-	-	-	-
<i>Kirchneriella obesa</i>	+	+	+	+	+	+	+
Pediastrum							
<i>Pediastrum borganum var longicorne</i>	-	-	-	+	-	-	-
<i>Pediastrum duplex</i>	+	+	+	+	+	+	+
Oocystaceae							
Eremosphaera							
<i>Eremosphaera gigas</i>	+	-	-	-	-	-	+
Micractinium							
<i>Micractinium pusillum</i>	-	+	+	-	-	-	-
Scenedesmaceae							
Coelastrum							
<i>Coelastrum cambricum</i>	+	+	+	-	-	-	-
<i>Coelastrum cambricum var intermedium</i>	+	-	-	+	-	-	-
Scenedesmus							
<i>Scenedesmus abundans</i>	-	+	+	-	-	-	-
<i>Scenedesmus acuminatus</i>	-	+	+	-	-	-	-
<i>Scenedesmus armatus var bicaudatus</i>	+	+	+	-	-	-	-
<i>Scenedesmus carinatus</i>	+	+	+	-	-	-	-
<i>Scenedesmus dispa</i>	-	+	+	-	-	-	-
<i>Scenedesmus javanensis</i>	-	+	+	-	-	-	+
<i>Scenedesmus nygocadi</i>	-	+	+	-	-	-	+
<i>Scenedesmus obliquus</i>	-	+	+	-	-	-	+
<i>Scenedesmus obtisus f ecornis</i>	-	+	+	-	-	-	+
<i>Scenedesmus opoliensis var mononesis</i>	-	+	+	+	-	-	-
<i>Scenedesmus perforatus f bicauda</i>	-	+	+	-	-	-	-
<i>Scenedesmus protuberans</i>	-	-	-	-	+	+	-
<i>Scenedesmus quadricauda var longispina</i>	-	+	+	+	+	+	-
<i>Scenedesmus sp</i>	+	-	-	-	-	-	-
Tetrastrum							
<i>Tetrastrum hétéracumthum</i>	-	+	+	-	-	-	-
<i>Tetrastrum staurogeniaforme</i>	-	+	+	-	-	-	+
Ulothrichaceae							
Ulothrix							
<i>Ulothrix bipyrenii</i>	-	-	-	+	-	-	-
<i>Ulothrix bipyronoidosa</i>	+	-	-	-	+	+	-
<i>Ulothrix subtilissima</i>	+	+	+	+	-	-	-
<i>Ulothrix tenerrina</i>	+	+	+	+	+	+	+
<i>Ulothrix tenuissima</i>	+	+	+	+	-	-	-
Closterium							
<i>Closterium abruptum</i>	+	-	-	-	-	-	-

<i>Closterium abruptum</i> var <i>brevius</i>	-	-	-	-	+	+	+
<i>Closterium calosparum</i>	-	+	+	-	-	-	-
<i>Closterium cherenbergii</i>	-	+	+	-	-	-	-
<i>Closterium diana</i>	+	-	-	-	-	-	-
<i>Closterium lincatul</i>	-	-	-	-	+	+	+
<i>Closterium lineatum</i> var <i>africanum</i>	+	-	-	-	-	-	+
<i>Closterium macilentum</i>	-	+	+	-	-	-	-
<i>Closterium moniliferum</i> f <i>giganteum</i>	+	-	-	-	-	-	-
<i>Closterium turgidum</i>	+	-	-	-	-	-	-
<i>Closterium jenneri</i> var <i>robustum</i>	+	-	-	-	-	-	-
Cosmarium							
<i>Cosmarium aversiforme</i>	-	+	+	-	-	-	-
<i>Cosmarium decoratum</i>	+	-	-	-	-	-	+
<i>Cosmarium dispersa</i>	+	-	-	-	-	-	+
<i>Cosmarium mononiforme</i>	-	+	+	-	-	-	-
<i>Cosmarium</i> sp	-	-	+	-	-	-	-
<i>Cosmarium undulatum</i> var <i>minutum</i>	-	+	+	-	-	-	+
<i>Cosmarum pusillum</i>	-	+	+	-	-	-	-
Euastrum							
<i>Euastrum spinulosum</i> var <i>lindae</i>	+	-	-	-	-	-	-
Staurastrum							
<i>Staurastrum biencanum</i>	-	+	+	-	-	-	-
<i>Staurastrum brevispina</i>	-	+	+	+	-	-	-
<i>Staurastrum caledonense</i>	+	+	+	+	-	-	-
<i>Staurastrum gladiusum</i>	-	+	+	-	-	-	+
<i>Staurastrum hexacerum</i>	+	+	+	+	-	-	+
<i>Staurastrum hystix</i>	-	+	+	+	-	-	-
<i>Staurastrum inflexum</i>	-	+	+	-	-	-	+
<i>Staurastrum intelliferum</i>	-	-	-	-	-	-	+
<i>Staurastrum mucranatus</i>	-	-	-	+	-	-	+
<i>Staurastrum setigerum</i> var <i>tristichum</i>	-	-	-	+	-	-	-
Zygnemataceae							
Spirogyra							
<i>Spirogyra irregularis</i>	-	+	+	-	-	-	-
Chrysophyta							
Chrysophyceae							
Mallomonas							
<i>Mallomonas bronchartiana</i>	+	+	+	+	+	+	-
Bacillariophyta							
Naviculaceae							
Amphora							
<i>Amphora coffaeiformis</i>	+	-	-	-	-	-	+
Cymbella							
<i>Cymbella affinis</i>	-	+	+	-	-	-	-
<i>Cymbella amphicephala</i>	+	-	-	-	-	-	-
<i>Cymbella caespitosa</i>	+	-	-	-	-	-	-
<i>Cymbella hybrida</i>	+	-	-	-	-	-	-
Gomphonema							
<i>Gomphonema affine</i>	+	-	-	-	-	-	-
<i>Gomphonema angustum</i>	-	+	+	+	-	-	-
<i>Gomphonema augur</i>	-	+	-	-	+	+	-
<i>Gomphonema gracile</i>	+	+	+	-	+	+	-
<i>Gomphonema parvulum</i>	-	+	+	+	-	-	-
<i>Gomphonema pusilla</i>	-	+	-	-	-	-	-

Navicula							
<i>Navicula americana</i>	+	-	-	-	-	-	-
<i>Navicula brasiliana</i>	-	+	+	-	-	-	-
<i>Navicula laevissima</i>	+	-	-	-	-	-	-
<i>Navicula pseudotuscula</i>	-	+	+	-	-	-	-
<i>Navicula pupula</i>	-	+	+	-	-	-	-
<i>Navicula stroemii</i>	-	+	+	-	+	+	-
Pinnularia							
<i>Pinnularia dactylus</i>	+	-	-	-	-	-	-
<i>Pinnularia gibba</i>	+	+	+	-	-	-	-
<i>Pinnularia lundii</i>	+	-	-	-	-	-	-
<i>Pinnularia macilenta</i>	+	+	+	+	-	-	-
<i>Pinnularia microstauron</i>	+	-	-	-	-	-	+
<i>Pinnularia subgibba</i>	-	-	-	+	-	-	+
Frustulia							
<i>Frustulia rhomboides</i>	-	+	+	-	-	-	-
<i>Frustulia scalpelliformis</i>	-	+	-	-	-	-	-
Fragilariaceae							
Fragilaria							
<i>Fragilaria capucina</i>	+	+	+	+	-	-	+
<i>fragilaria construens</i>	+	-	+	-	-	-	+
<i>Fragilaria SP</i>	-	+	+	-	-	-	+
<i>Fragilaria ulna</i>	+	-	-	-	-	-	-
Surirellaceae							
Surirella							
<i>Surirella elegans</i>	-	+	+	-	-	-	-
<i>Surirella ovalis</i>	-	+	+	-	-	-	-
<i>Surirella robusta</i>	+	-	-	+	-	-	-
Aulacaseiraceae							
Aulacaseira							
<i>Aulacaseira islandica</i>	+	-	-	+	-	-	+
Nitzschiaceae							
Orthoscira							
<i>Orthoscira roeseana</i>	-	-	-	+	-	-	+
<i>Nitzschia sp</i>	-	+	+	-	-	-	-
Euglenophyta							
Euglenaceae							
Euglena							
<i>Euglena acus</i>	-	+	+	-	-	-	-
<i>Euglena anabaena var minima</i>	+	-	-	-	-	-	+
<i>Euglena chrenbergii</i>	+	-	-	-	-	-	-
<i>Euglena chrenbergii var africana</i>	+	-	-	-	-	-	-
<i>Euglena limnophila</i>	-	+	+	+	-	-	-
<i>Euglena oxymis var charkowiensis</i>	+	-	-	-	-	-	-
<i>Euglena oxyuris f minima</i>	-	+	+	-	-	-	-
<i>Euglena pisciforme</i>	+	-	-	-	+	+	-
<i>Euglena rostrifera</i>	-	+	+	-	-	-	-
<i>Euglena sanguinea</i>	+	-	-	-	-	-	-
<i>Euglena sp</i>	-	+	-	-	-	-	-
<i>Euglena spirogyra</i>	+	-	-	-	+	+	-
<i>Euglena texta</i>	+	+	+	-	-	-	+
<i>Euglena variabilis</i>	-	-	+	-	-	-	-
Lepocinclis							
<i>Lepocinclis ovum</i>	-	+	+	-	+	+	+
<i>Lepocinclis ovum var butschu</i>	+	-	-	-	-	-	-
<i>Lepocinclis ovum var gracilicauda</i>	+	-	-	-	-	-	-
<i>Leptocinclis acuminatua</i>	-	-	-	-	+	+	+

Phacus							
<i>Phacus acuminatus</i>	+	+	+	-	-	-	+
<i>Phacus applanatus</i>	-	-	-	-	+	+	-
<i>Phacus brachykentron</i>	+	-	-	-	-	-	-
<i>Phacus caudatus</i>	+	-	-	-	-	-	-
<i>Phacus curvicauda</i>	+	-	-	-	-	-	+
<i>Phacus gamsii</i>	-	-	-	-	+	+	-
<i>phacus glaber</i>	+	+	+	-	-	-	-
<i>Phacus hamatus</i>	+	-	-	-	-	-	-
<i>Phacus horidus</i>	+	-	-	-	-	-	-
<i>Phacus lismorensis</i>	+	-	-	-	-	-	-
<i>Phacus longicauda</i>	+	-	-	-	-	-	-
<i>Phacus platalea</i>	+	-	-	-	-	-	-
<i>Phacus pleuroncetes</i>	+	-	-	-	-	-	-
<i>Phacus ranula</i>	-	+	+	-	+	+	-
<i>Phacus tartus</i>	+	+	+	-	-	-	-
Trachelomonas							
<i>Trachelomonas armata f longicollis</i>	+	-	-	-	-	-	+
<i>Trachelomonas conica</i>	+	+	+	-	-	-	-
<i>Trachelomonas conica var granulata</i>	+	-	-	-	-	-	-
<i>Trachelomonas conica var punctata</i>	+	-	-	-	-	-	-
<i>Trachelomonas cylindrica</i>	-	-	-	-	+	+	-
<i>Trachelomonas hispida var coranata</i>	+	+	+	-	+	+	-
<i>Trachelomonas hispida</i>	+	-	-	-	-	-	-
<i>Trachelomonas hispida var duplex</i>	-	+	+	-	-	-	-
<i>Trachelomonas klebsii</i>	+	-	-	-	-	-	-
<i>Trachelomonas molesta</i>	+	-	-	-	-	-	+
<i>Trachelomonas nigerica</i>	+	-	-	-	-	-	-
<i>Trachelomonas oblonga</i>	+	-	-	-	-	-	-
<i>Trachelomonas planctonica var oblonga</i>	+	-	-	-	-	-	+
<i>Trachelomonas verrucosa</i>	-	+	+	-	-	-	-
<i>Trachelomonas volcascinopsis</i>	-	+	+	-	-	-	+
<i>Trachelomonas volvocina</i>	+	+	+	+	-	-	-
<i>Trachelomonas volvocina var punctata</i>	+	-	-	-	-	-	-
Colaciaceae							
Colacium							
<i>Colacium cyclopicola</i>	+	-	-	-	-	-	-

+ = present - = absent

3.2. Species richness and distribution in the ponds in the western highlands of Cameroon

Table 2 shows that of the 220 species identified, only 3 species (*Chlamydomonas globosa*, *Kirchneriella obesa* and *Pediastrum duplex*) were represented in all the ponds. With the exception of the Foubot ponds fertilised with cow dung, *Mallomonas bronchartiana* has been identified in all environments. However, apart from the ponds of Bamendou fertilized with cow dung, *Microcystis aeruginosa*, *Microcystis incerta* and *Microcystis robusta* were identified in all types of ponds. Independent of the type of fertiliser, the species *Micractinium pusillum* was exclusively exemplified in Batié. *Pediastrum borganum var longicorne* and *Ulothrix bipyrenii* were restricted to the Dschang fertilized with chicken manures.

Of the 220 species, 37 (*Nostoc parmeloides*, *Oscillatoria teribriformis*, *Chlamydomonas muriella* etc) were represented only in Bamendou. Merely five species (*Ankistrodesmus fusiformis*, *Scenedesmus protuberans*, *Phacus applanatus*, *Phacus gamsii* *Trachelomonas cylindrica*) were recognised singularly in the ponds of Fokoué. *Staurastrum intelliferum* was the only species unique to Fombot municipality ponds.

For the similar type of fertilizer, 13 species (*Merismopedia convoluta*, *Chlamydomonas epiphytica*, *Eremosphaera gigas*, etc.) were identified only in ponds fertilised with cow dung. On the other hand, no species was singularly observed in the ponds fertilised with pig manure.

3.3. Richness and distribution of genera in the ponds of West Cameroon

The distribution of phytoplankton genera is summarised in table 2. Of the 38 genera listed, only 8 or 21.05% (*Chlamydomonas*, *Kirchneriella*, *Pediastrum*, *Scenedesmus*, *Gomphonema*, *Navicula*, *Euglena* and *Trachelomonas*) were recognised in all the categories of pond.

The genera represented only in the ponds fertilized with the Batié feed (31 genera; *Tetrastrum*, *spirogira*, *Cosmarium*, etc.) were more abundant than in all the categories of pond.

When contrasting the sites with the same type of fertilizer, the genera listed specifically in Batié (17 genera; *Merismopedopedia*, *Oscillatoria*, *Staurastrum*, etc.) were more abundant than those listed only in Fokoué (0 genus) fertilised with pig manure. Relative to cow dung, the genera listed only in Bamendou (09 genera; *Euastrum*, *Colastrum*, *Oscillatoria*, etc.) were more than the three genera (*Gomphosphaeria*, *Monoraphidium*, *Tetrastum*) listed in Fombot.

3.4. Richness and distribution of families in the ponds of West Cameroon

The distribution of phytoplankton families as summarised in Table 2 shows that, of the 19 families, merely 6 or 31.58% (Chlorophyceae, Hydrodictyaceae, Scenedesmaceae, Ulothrichaceae, Naviculaceae and Euglenophyceae) were exemplified in all types of ponds.

The Nastocaceae was exclusively identified in the ponds of Bamendou fertilized with cow dung. However, the Oocystaceae was specific to ponds fertilised with the Batié animal feed. The Zygnemataceae was only listed in Batié irrespective of the type of fertilizer used. The Surirellaceae was only represented in the ponds of Batié fertilized with pig manure and of Dschang fertilized with chicken manures.


When comparing sites with the identical type of fertiliser, the families Merimopediaceae, Oscillatoriaceae, Gomphosphaeriaceae, Zynemataceae, Surirellaceae and Nitzscheiaceae represented in the Batié ponds fertilized with pig manure were absent in Fokoué. However, all the families (Chlorophyceae, Chrocococaceae, Hydrodictyaceae, Scenedesmaceae, Ulothricheaceae, Chrysophyceae, Surirellaceae, Naviculaceae and Euglenophyceae) listed in the ponds of Fokoué fertilized with pig manure were also identified in the ponds fertilized with pig manure in Batié. Relative to cow dung, the Merimopediaceae, Naviculaceae, Fragilariaceae and Aulacoseinaceae were listed in Bamendou and Fombot. However, the families Colaciaceae, Chrysophyceae, Oscillatoriaceae and Nastocaceae identified in Bamendou were absent in Fombot. Additionally, only the Chrocococaceae present in Fombot was absent in Bamendou. When comparing sites with equal type of fertilizer, the Merimopediaceae, Chlorophyceae, Hydrodictyaceae, Scenedesmaceae, Ulothricheaceae, Naviculaceae, Fragilariaceae, Euglenophyceae and Aulacoseinaceae were recognised in the ponds of Bamendou and Fombot fertilised with cow dung.

3.5. Phytoplankton density according to site and type of fertilizer

The average phytoplankton densities according to sampling site and the type of fertiliser are illustrated in Figure 1. The ponds of Dschang fertilised with chicken manure had significantly higher densities ($p < 0.05$) as compared to Foubot and Fokoué regardless of the type of fertiliser.

The phytoplankton mass was significantly ($p < 0.05$) higher in Bamendou as matched with Foubot (cow dung). In pond fertilised with pig manure, phytoplankton density was significantly higher in Batié compared to Fokoué.

At the same site, at Batié as well as at Fokoué, densities were comparable regardless of the type of fertilizer administered.



a, b, c: bars carrying the same letters are not pas significantly different ($p > 0,05$)

Figure 1: Phytoplankton density according to sampling site and type of fertiliser

3.6. Effect of site and type of fertiliser on phytoplankton species frequency

The influence of the site and the type of fertiliser on the phytoplankton species frequency is illustrated in Figure 2-7. None of the three phytoplankton species common to all ponds was dominant in all the ponds. Nevertheless, *Microcystis robusta* presented the highest frequency, i.e. 53.36% of the total percentage of species in Foubot ponds (fertilised with cow dung) as compared to the frequency of all the species listed in all the types of ponds. *Kirchneriela obesa* was more foremost in the unfertilized ponds in Fokoué.

In the same sampling site, in Batié, *Microcystis aeruginosa* was more recurrent in ponds fertilized with pig manure while *Microcystis incerta* was more abundant in ponds fertilised with animal feed. In Fokoué, the species *Ulothrix bypirenoida* was more frequent in ponds fertilised with pig manure, yet *Kirchneriela obesa* was more prevailing in unfertilized ponds.

When comparing the sites with equal type of fertilizer, *M. aeruginosa* was more frequent in Batié whereas *U. bypirenoida* was more represented in Fokoué (pig manure). Relative to cow dung, *Nostoc parmeloides* was more represented in Bamendou while *Microcystis robusta* was more frequent in Foubot.




Figure 2: Phytoplankton relative frequency in Bamendou (cow dung)




Figure 3: Phytoplankton relative frequency in Batié (pig manure)




Figure 4: Phytoplankton relative frequency in Batié (Wheat bran)




Figure 5: Phytoplankton relative frequency in Fokoué (pig manure)




Figure 6: Phytoplankton relative frequency in Fokoué (Not fertilised)




Figure 7: Phytoplankton relative frequency in Foubot (cow dung)

3.7. Phytoplankton diversity indices according to sampling site and type of fertiliser

The impact of site and type of fertiliser on diversity indices is summarized in Table 3.

Table 3: Phytoplankton diversity index according to sampling site and type of fertiliser

Diversity index (Bit/ ind)	Sampling sites and types of fertilisers						
	Bamendou	Batié		Dschang	Fokoué		Foumbot
	Cow dung	Pig manure	Wheat bran	Chicken manure	Pig manure	Not fertilised	Cow dung
Shannon - Weaver (H')	2.27	2.37	2.15	1.34	1.09	1.14	1.99
Piélou Evenness (J)	0.42	0.57	0.44	0.22	0.12	0.10	0.31

Table 3 displays that independently of the site and the type of fertiliser, the highest values of the Shannon-Weaver, and the Piélou evenness indices were observed in the Batié ponds enriched with pig manure while the lowest value occurred in unfertilized ponds of Fokoué.

Sampling site being equal, in Batié, the values of these diversity indices were higher in the ponds fertilized with pig manure and lower in the ponds enriched with animal feed. In Fokoué, the highest Shannon-Weaver index was recorded in unfertilized ponds as compared to ponds fertilised with pig manure. However, the highest Piélou evenness index value was observed in ponds enriched with pig manure compared to those receiving no fertiliser.

In relation to sites with the same type of fertiliser, the highest Shannon-Weaver and Piélou equitability indices were recorded in Bamendou compared to Foumbot for cow dung. For pig manure, these values were higher in the ponds of Batié compared to those of Fokoué.

3.8. Correlation between water physicochemical characteristics and phytoplankton density according to sampling site and type of fertiliser

The correlations between the physicochemical characteristics of the water and the phytoplankton densities (Table 4) showed that, with the exception of the relationship between the density of phytoplankton and phosphates, on the one hand and the transparency on the other hand, which were not significant, they were significant ($p < 0.01$), negative and strong for all the other physicochemical characteristics of the water in the unfertilised ponds of Fokoué. Likewise, phytoplankton density was negatively and strongly correlated with dissolved oxygen in Dschang ponds fertilized with chicken droppings. Conversely, the density of phytoplankton was positively and very strongly correlated ($p < 0.01$) with pH in ponds fertilized with cow dung regardless of the site.

Table 4: Correlation between physicochemical characteristics of water and phytoplankton densities

Physicochemical characteristics of water	Phytoplankton density						
	Bamendou	Batié		Dschang	Fokoué		Foumbot
	Cow dung	Pig manure	Wheat bran	Chicken droppings	Pig manure	Not fertilized	Cow dung
Nitrates (NO ₃ ⁻)	0.03	-0.11	-0.17	0,13	-0.22	-0.85**	0.29
Nitrites (NO ₂ ⁻)	0.22	-0.01	-0.10	0,29	-0.22	-0.68**	-0.47
Dissolved Oxygen (O ₂)	0.16	0.18	0.14	-0,99*	0.45	-0.99**	0.31
pH	0.99**	0.23	-0.22	0,18	0.22	-0.93**	0.99**
Orthophosphates (PO ₄ ³⁻)	-0.56	-0.21	-0.52	-0,12	-0.52	0.22	-0.52
Temperature	0.26	0.14	0.12	0,08	-0.22	-0.77**	-0.18
Transparency	-0.32	0.19	-0.41	0.13	-0.22	0.28	0.23

** : significant correlation at $p < 0.01$ (bilateral)

4. DISCUSSION

Phytoplankton species and genus richness was highest in unfertilized Fokoué ponds and animal feed enriched ponds of Batié respectively. This observation supports the findings of Nana et al. [31] in unfertilized ponds and may be linked to the influence of the use of fertilisers on the physicochemical characteristics of the water. In fact, Patrick [32] revealed a decline in biodiversity from unpolluted to polluted backgrounds. Furthermore, Moss [33] suggested that only species capable of adapting to a high nutrient enrichment of the environment and to an environment presenting ostensible extreme will have an advantage in eutrophic conditions; this ensues an uneven distribution of taxa. The population of taxa unable to adapt decreases and that of taxa able to adapt amplifies, and the proportions of taxa according to the trophic level are thus modified according to the populations of these taxa. Radji et al. [34] similarly reported that the physicochemical characteristics of the water influenced the dynamics of phytoplankton populations. The highest species richness in unfertilised ponds showed the great variability of species living in a pure or dystrophic environment. The imbalance observed in the fertilised environments would be the effect of the fertilisers brought continuously which modifies the physicochemical characteristics of the water.

The distribution of species showed that of the 220 species identified, 3 species (*Chlamydomonas globosa*, *Kirchneriella obesa* and *Pediastrum duplex*) are registered in all sites regardless the type of fertiliser. *Kirchneriella obesa*, *Pediastrum duplex* and *Microcystis aeruginosa* were the most represented in unfertilised ponds. This result is similar to that reported by Nana et al. [16] who observed the majority of *Merismopedia elegans*, *Microcystis aeruginosa* and *Nostoc entophyllum* in unfertilised ponds compared to ponds fertilised with different doses of chicken manures. This tendency to mainly have the species *M. aeruginosa* would be explained by the fact that it constitutes the most common toxic cyanobacterial blooms in freshwaters.

The distribution of phytoplankton genera disclosed that among the genera identified in all sites regardless of the nature of the fertiliser dispensed, *Chlamydomonas* and *Pediastrum* had been listed. This result is explained by their ability to encyst and remain immobile in bodies of water respectively.

The highest richness of phytoplankton families was recorded in the Dschang ponds fertilised with chicken droppings. This result is similar to that reported by Nana et al. [31] in ponds fertilised with chicken manures and pig manure and may be related to the high organic load of this enricher.

The highest density was recorded in the ponds of Dschang fertilised with chicken droppings. These results are contradictory to those observed by Nana et al. [31] and may be linked to the production conditions of phytoplankton organisms and the impact of collection sites. Indeed, the highest densities in the ponds of Dschang might be linked to the henhouse on stilts which would offer higher concentrations of nutrient salts, the basis of phytoplankton production.

The diversity indices were relatively low in the unfertilised ponds, as compared to the ponds fertilised with pig manure in Fokoué and chicken droppings in Dschang. This result could be elucidated by a high specific frequency of *Kirchneriella obesa* (17.70%), *Ulothrix bipyronoidosa* (26.22%) and *Microcystis incerta* (41.02%) respectively in these ponds. The low values of the diversity indices in these ponds are similar to those testified by Kemta et al. [13] in the Yaoundé Municipal Lake and may be related to a young population with a high potential of multiplication with a predominance of one species or a small number of species.

5. CONCLUSION

The phytoplankton species, genera, families and phyla richness were upset by the sampling sites and types of fertilisers. To be sure, the richness of phytoplankton species and genera was higher in the unfertilized ponds of Fokoué and the ponds of Batié fertilised with animal feed respectively. The highest values of phytoplankton phyla richness were detected in Dschang, Fokoué, Foubot and in the ponds receiving animal feed from Batié. Phytoplankton density, Shannon-Weaver and Piéluou evenness indices, were significantly disturbed by sampling sites and type of fertiliser.

REFERENCES

- [1] Kirkpatrick C, Diakosavvas D. Food insecurity and foreign-exchange constraints in sub-Saharan Africa. *The Journal of Modern African Studies* 1985;23:239-50.
- [2] Schulten G. Hunger technology and society: post-harvest losses in tropical Africa and their prevention. *Food and Nutrition Bulletin* 1982;4:1-7.
- [3] Guengant J-P. How to benefit from the demographic dividend. *Namely* 2011;9.
- [4] Aall C. Fish Protein Resources for Human Consumption: Summary of a Seminar Held in Colombo, Sri Lanka. *Food and Nutrition Bulletin* 1982;4:1-8.
- [5] Azam F, Malfatti F. Microbial structuring of marine ecosystems. *Nature Reviews Microbiology* 2007;5:782-91.
- [6] Grogan N. Structure, functioning and dynamics of phytoplankton in Lake Taabo (Côte d'Ivoire) 2012.
- [7] Testard P, Pourriot R, Miquelis A, Rougier C. Functioning of the fluvial ecosystem: organization and role of zooplankton. *PIREN-Seine Synthesis Report* 1993:3-38.
- [8] Garnier J, Billen G, Sanchez N, Leporcq B. Ecological functioning of the Marne reservoir (upper Seine basin, France). *Regulated Rivers: Research & Management: An International Journal Devoted to River Research and Management* 2000;16:51-71.

- [9] Rolland A. Dynamics and diversity of phytoplankton in the Marne reservoir (Seine watershed): University of Savoie; 2009.
- [10] Henson SA, Cael B, Allen SR, Dutkiewicz S. Future phytoplankton diversity in a changing climate. *Naturecommunications* 2021;12:1-8.
- [11] Cuna E, Alcocer J, Gaytán M, Caballero M. Phytoplankton Biodiversity in Two Tropical, High Mountain Lakes in Central Mexico. *Diversity* 2022;14:42.
- [12] Roldán G, Ruiz E, Wetzel R, Gopal B. *Limnology in developing countries*. 2001.
- [13] Kemka N, Njiné T, Togouet S, Niyitegeka D, Nola M, Monkiedje A, et al. Phytoplankton of the Yaounde municipal lake (Cameroon): ecological succession and population structure. *Revue des Sciences de l'Eau/Journal of Water Science* 2004;17:301-16.
- [14] Fonge B, Tening A, Egbe E, Yinda G, Fongod A, Achu R. Phytoplankton diversity and abundance in Ndop wetland plain, Cameroon. *African Journal of Environmental Science and Technology* 2012;6:247-57.
- [15] Fonge BA, Tabot PT, Mange CA, Mumbang C. Phytoplankton community structure and physico-chemical characteristics of streams flowing through an agro-plantation complex in Tiko, Cameroon. *Journal of Ecology and The Natural Environment* 2015;7:170-9.
- [16] Nana TA, Songmo B, Efole ET, Fonkwa G, Kom MF. Comparative effect of the dose of pig manure used as pond fertilizer on the richness and distribution of phytoplankton taxa. *Journal of Applied Biosciences* 2020;152:15630-9.
- [17] Agadjihouede H, Bonou C, Chikou A, Laleye P. Comparative production of zooplankton in ponds fertilized with poultry droppings and cow dung. *International Journal of Biological and Chemical Sciences* 2010;4.
- [18] APHA. Standard methods for the examination of water and wastewater. American Public Health Association 2005;21st Edition: American Water Works Association/Water Environment Federation, Washington DC:2-58.
- [19] Pihan J, Landragin G. Model for predicting the acute toxicity of river waters to fish: With 11 figures and 1 table in the text. *Internationale Vereinigung für theoretische und angewandte Limnologie: Verhandlungen* 1985;22:2457-62.
- [20] Nguetsop V, Fonkou T, Lekeufack M, Pinta J. Seaweed assemblages and relationships with some environmental parameters in two marshy sites in West Cameroon. *Water Science Review/Journal of Water Science* 2009;22:15-27.
- [21] Ward HB, Whipple GC. *Fresh-water biology*: John Wiley & sons, Incorporated; 1918.
- [22] Bourrelly P. Five years of freshwater algology (1950–1954). *Bulletin of the Botanical Society of France* 1955;102:134-90.
- [23] Bourrelly P, Couté A. Some freshwater algae from French Guiana. *Amazoniana: Limnologia et Oecologia Regionalis Systematis Fluminis Amazonas* 1982;7:221-92.
- [24] Bourrelly P, Couté A, Thérézien Y. Freshwater algae from Kerguelen and Crozet islands: With 2 tables in the text. *Internationale Vereinigung für theoretische und angewandte Limnologie: Verhandlungen* 1978;20:2347-50.

- [25] CARDINAL C. Planktonic algae from the Seine basin (with the exception of Cyanophyceae and Diatoms). 1979.
- [26] Utermöhl H. Zur vervollkommnung der quantitativen phytoplankton-methodik: Mit 1 Tabelle und 15 abbildungen im Text und auf 1 Tafel. Internationale Vereinigung für theoretische und angewandte Limnologie: Mitteilungen 1958;9:1-38.
- [27] Shannon CE. A mathematical theory of communication. The Bell System Technical Journal 1948;27:379-423.
- [28] Mårtensson R. Species and Biological Diversity-Choices of Diversity Indices and Their Potential Consequences for Nature Conservation. In: Klimaforskning C-CFM-O, editor. Lunds University 2016. p. 28.
- [29] Shekhar TS, Kiran B, Puttaiah E, Shivaraj Y, Mahadevan K. Phytoplankton as index of water quality with reference to industrial pollution. Journal of Environmental Biology 2008;29:233.
- [30] Pielou EC. An introduction to mathematical ecology. New York: Wiley-Interscience; 1969.
- [31] Algeria NT, Thomas EE, Togouet Z, Hubert S, Joseph T. Effects of doses of chicken manure on the biodiversity of zooplankton populations in ponds. 2018.
- [32] Triplet P. Dictionary of biological diversity and nature conservation. Electronic documentation https://societe-zoologique.fr/sites/default/files/2019-02/Dictionnaire-diversite-biologique-conservationnature_2019.pdf 2019.
- [33] Moss BR. Ecology of fresh waters: man and medium, past to future: John Wiley & Sons; 2009.
- [34] Radji R, Bandje A, Issifou L, Edoth T, Kokou K. Diversity and dynamics of phytoplankton assemblages in aquatic ecosystems in southern Togo. Africa Science: International Journal of Science and Technology 2013;9:67-77.