

Temporal variation of temperature inversions and their effect on fine particulate matter (PM_{2.5}) concentrations in a Vietnamese metropolis

ABSTRACT

This study aims to investigate the temporal variation of temperature inversions, including inversion frequency and inversion strength, in Ho Chi Minh City (HCM), a metropolis of Vietnam. The effect of the temperature inversions on PM_{2.5} concentrations in the dry and rainy seasons of HCM City was also identified. The results revealed that the surface inversions, which are temperature inversions below 300 m, were more frequent at 12Z (7 PM local time) in the rainy season. In addition, the stronger inversions were found in the dry season, when the higher PM_{2.5} concentrations were also observed, suggesting an influence of the temperature inversions on an increase in PM_{2.5} concentrations. This suggestion was also supported by the significantly positive correlation between PM_{2.5} concentrations and inversion frequency in the dry season. However, the temperature inversions were believed to have a minor effect on the PM_{2.5} concentrations in the rainy season since no significant correlation between the PM_{2.5} concentrations and inversion frequency or strength was found for this season. The findings of this study contribute to understanding of the temperature inversions and their effect on PM_{2.5} concentrations in a metropolis of Vietnam, a developing country in Southeast Asia. Based on these results, solutions for reducing the PM_{2.5} pollution in the study area can be obtained.

Keywords: Temperature inversion; PM_{2.5}; Ho Chi Minh City; Vietnam.

1. INTRODUCTION

Temperature inversion is a phenomenon when atmospheric temperature increases with height in the troposphere [1]. The temperature inversions can be classified as radiation inversions and subsidence inversions. The radiation inversion, also known as surface temperature inversion, develops mostly at night when the ground becomes cooler due to emissions of longwave radiation [2]. Consequently, the air temperature in the lower planetary boundary layer is lower than that in the overlying layer. The subsidence inversion, which is also called an upper surface temperature inversion, forms when the air layer descend and then is compressed and heated due to an increase in atmospheric pressure [2], resulting in warmer air at higher altitudes.

The temperature inversions can act as a cap, preventing vertical dispersion of air pollutants [3], and thus, pollutant levels increase near the ground during the inversion. The relationship between temperature inversion and air pollution has been reported in several previous studies [4-9]. For instance, concentrations of several pollutants, including NO₂, SO₂, fine particulate matters (PM_{2.5}), were found to be higher when strong inversions occurred in Ha Noi City of Vietnam [4]. Of these pollutants, PM_{2.5}, which is a particle less than 2.5 micrometers in diameter, is a frequently monitored pollutant because of its harmful effect on human health. In particular, the components of PM_{2.5} contain several chemicals, including metals, ions, and organic compounds (e.g., polycyclic aromatic hydrocarbons) that have high toxicities [10]. Added to this, PM_{2.5} can penetrate deep into the human respiratory system through inhalation due to its small size, harm human health, and cause several diseases, such as lung diseases, asthma, and acute respiratory distress [10]. Therefore, it is essential

36 to understand the factors governing the PM_{2.5} levels. These factors include meteorological
37 conditions (e.g., rainfall amount, wind speed, and air temperature), emission sources
38 emitting air pollutants, and temperature inversions trapping the pollutants.
39

40 Ho Chi Minh City (HCM City), located in southeastern Vietnam, is a metropolis and
41 economic center of the country. HCM City has a tropical monsoon climate with two distinct
42 seasons, including dry and rainy seasons covering a period of November–April and May–
43 October, respectively [11]. Air pollutants (e.g., PM_{2.5}, PM₁₀, SO₂, NO_x, and CO) in HCM City
44 have been linked to vehicle exhaust, industrial emissions, and area sources (i.e., households
45 and construction sites) [12,13]. Most previous studies on PM_{2.5} pollution in HCM City have
46 mainly focused on monitoring concentrations, investigating the correlation between
47 concentrations and meteorological conditions, and calculating the emission inventory of
48 PM_{2.5}. The temperature inversions can contribute to PM_{2.5} pollution [4-9]., however, there has
49 been no information related to the temperature inversions in HCM City, such as frequency,
50 strength, and seasonal variation of the inversions.
51

52 This study aims to investigate the temporal variation of the temperature inversions,
53 considering inversion frequency and inversion strength, and determine the effects of the
54 inversions on the PM_{2.5} concentrations in HCM City, Vietnam. Results from this study help to
55 increase understanding of the temperature inversions and their relation to the PM_{2.5} pollution
56 in a metropolis in Vietnam, a developing country in Southeast Asia. Based on these results,
57 solutions for reducing the PM_{2.5} pollution in the study area can be obtained.
58

59 2. METHODOLOGY

60 61 2.1 Study area

62
63 Ho Chi Minh City (HCM City) is one of the municipal cities of Vietnam and this city covers
64 approximately an area of 2,061 km² and is coordinated between latitude 10°46'32"N and
65 longitude 106°42'07"E [11]. There are two seasons in this city, which are dry and rainy
66 seasons. The average rainy day and rainfall amount in the dry season, lasting from
67 November to April of the following year, are 5 days and 40.6 mm, respectively. The rainy
68 season covers a period of May to October and tends to have more rainy days and greater
69 rainfall amount, which are 21 days and 281.2 mm, respectively [14].
70


71 2.2 Data collection

72
73 To determine the temperature inversion, radiosonde data of HCM City were downloaded via
74 the Wyoming Weather website <http://weather.uwyo.edu/upperair/sounding.html>, operated by
75 the Department of Atmospheric Science, College of Engineering, the University of Wyoming.
76 The code and identifier of the station are 48900 and VVTS, respectively (Fig. 1). This station
77 (latitude: 10°48'36"N, longitude:106°39'36"E) is 10 m above sea level and situated in
78 northwestern HCM City. The radiosonde data were gathered twice per day at 00Z (7 AM
79 local time) and 12Z (7 PM local time) from 1 January 2019 to 31 December 2020 and
80 contained values of several meteorological parameters, including potential temperature,
81 wind speed, and wind direction at different air layer heights.
82

83 To understand the effect of the temperature inversions on PM_{2.5} pollution, data on PM_{2.5}
84 concentrations over the study period (i.e., 1 January 2019 to 31 December 2020) in HCM
85 City were obtained from the AirNow monitoring network (<https://airnow.com>), operated by the
86 U.S. Environmental Protection Agency. The technique used for PM_{2.5} monitoring is a beta
87 attenuation monitor, a detection technique of PM_{2.5} based on the absorption of beta radiation
88 by solid particles [15]. In addition, the PM_{2.5} monitoring station is at the HCM City U.S

89 Consultant (latitude: 10°46'58.76"N, longitude: 106°42'2.05"E), located in a downtown area
90 of District 1, HCM City (Fig. 1). The hourly PM_{2.5} concentrations obtained from the AirNow
91 monitoring network, were pre-processed to remove missing and abnormal data (e.g., null
92 and outliers). The processed data were then averaged into annual, seasonal, and monthly
93 concentrations for further usage.

94
95



96
97

98 **Fig. 1. Map of study area showing locations of the radiosonde and PM_{2.5} monitoring**
99 **stations in Ho Chi Minh City, Vietnam. The red line illustrates the city boundary.**

100

101 In addition, the monthly rainfall amount and the total of rainy days per month in HCM City
102 were obtained from the Mac Dinh Chi station (latitude: 10°47'3.72"N, longitude:
103 106°42'1.04"E). This station is approximately 200 m from the PM_{2.5} monitoring station and is
104 operated by the Hydrometeorological Observatory-Southern region.

105

106 **2.3 Determination of temperature inversion, inversion strength, and inversion** 107 **frequency**

108

109 The temperature inversion is identified based on a ratio of the difference in potential
110 temperature, θT (K), and the height difference, θz (m) [16]. A positive value of $\theta T/\theta z$,
111 meaning that potential temperature increases with increasing height, reflects a stable
112 condition of the atmosphere or an occurrence of the temperature inversion [17]. Reversely, if
113 $\theta T/\theta z$ is negative, the atmosphere is unstable. Additionally, if θT equals zero, the potential
114 temperature un-changes with height, and this situation is defined as a neutral condition. In
115 this study, the temperature inversions were checked at several atmospheric layers and were
116 categorized into three types according to their bottom heights, namely surface inversion (SI),
117 elevated inversion (EI), and lower-troposphere inversion (LTI). Particularly, the SIs has a
118 bottom height of 10 to 300 m. The EIs and LTIs refer to the inversions with bottom heights in
119 a range of 300–2,000 m and above 2,000 m, respectively.

120

121 The inversion strength was also calculated to understand the characteristics of the
122 temperature inversion in HCM City. The inversion strength is defined as the change in
123 atmospheric temperature over the first 100 m above from ground level [18] and is expressed
124 as °C/m. Moreover, the inversion frequency was also computed as the percentage of
125 inversion observation over the total events (i.e., inversion and non-inversion events) [16].

126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149


2.4 Data analysis and visualization

This study used Python programming language (<https://www.python.org>) with *numpy* [19] and *pandas* [20] libraries to pre-process the radiosonde and $PM_{2.5}$ concentration data, then compute the temperature inversion, inversion strength, and inversion frequency for the 2019 and 2020. The OriginPro version 2020 (OriginLab, USA) was used to visualize the calculation results. Moreover, the statistical analysis, such as a t-test for hypothesis testing, was performed using *scipy*, a Python library.

3. RESULTS AND DISCUSSION

3.1 Temporal variation of $PM_{2.5}$ concentrations

The $PM_{2.5}$ concentrations in HCM City from 2019 to 2020 are illustrated in Fig. 2. In general, the concentrations of $PM_{2.5}$ observed in 2019 (range: 15.1–42.6 $\mu\text{g}/\text{m}^3$, mean: 25.5 $\mu\text{g}/\text{m}^3$) were 1.3 times higher than those in 2020 (range: 17.2–32.7 $\mu\text{g}/\text{m}^3$, mean: 23.2 $\mu\text{g}/\text{m}^3$). Additionally, there was no statistically significant difference between the concentrations of 2019 and 2020 (Mann-Whitney rank-sum test, $P = 0.442$). Noticeably, the mean $PM_{2.5}$ concentrations of HCM City in 2019 and 2020 were 1.5–5.0 times higher than the WHO and Vietnamese air quality guidelines for annual $PM_{2.5}$ levels, which are 5 [21] and 25 [22] $\mu\text{g}/\text{m}^3$, respectively. Previous studies on monitoring $PM_{2.5}$ in HCM City also reported an exceedance of the monitored concentrations over the Vietnamese criteria level [13,23,24].




150
151
152
153
154
155
156
157
158
159
160
161

Fig. 2. Concentrations of $PM_{2.5}$ in Ho Chi Minh City of Vietnam shown in (a) yearly variation and (b) monthly and seasonal variation.

Regarding seasonal variation, the $PM_{2.5}$ concentrations in the dry season (range: 17.2–42.6 $\mu\text{g}/\text{m}^3$, mean: 25.1 $\mu\text{g}/\text{m}^3$) were 1.1–1.9 times higher than those in the rainy season (range: 15.1–36.7 $\mu\text{g}/\text{m}^3$, mean: 22.1 $\mu\text{g}/\text{m}^3$). A similar observation was also reported in previous studies [23,25]. The lower concentrations in the rainy season would be contributed by wet deposition removing particulate matters in the atmosphere. The greater rainfall amount and more rainy days in the rainy season (May–October) (Fig. 3) would support this interpretation. For monthly variation, the $PM_{2.5}$ concentrations in several months, including October, November, December, and January (27.9 ± 12.4 , 36.1 ± 9.2 , 31.0 ± 3.2 , and 32.7 ± 1.6

162 $\mu\text{g}/\text{m}^3$, respectively) were 1.4–2.0 times higher than those in the other months (20.5 ± 2.5
 163 $\mu\text{g}/\text{m}^3$). The period of October to January exhibited lower rainfall amount (Fig. 3a) and rainy
 164 days (Fig. 3b), therefore, the removal ability of $\text{PM}_{2.5}$ through wet deposition declined and led
 165 to an increase of $\text{PM}_{2.5}$ concentrations in the atmosphere. In addition, the lower wind speed
 166 in this period (1.9–2.5 m/s) (Fig. 3c) would reduce the dispersion of air pollutants and
 167 contribute to the higher $\text{PM}_{2.5}$ concentrations in HCM City from October to January. The
 168 temperature inversion might also cause an elevation of $\text{PM}_{2.5}$ concentrations in this period
 169 (i.e., October to January) and this issue is more discussed in Section 3.3.
 170



171
 172
 173
 174
 175

Fig. 3. Rainfall amount (a), number of rainy days (b), and wind speed (c) of Ho Chi Minh City shown as monthly averages.


176 3.2 Characteristics of temperature inversions

177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190

The frequency and strength of the temperature inversions at 00Z (7 AM local time) and 12Z (7 PM local time) in the dry and rainy seasons of HCM City were determined. In addition, the inversions observed over the study period were classified as surface inversions (SIs), elevated inversions (EIs), and lower-troposphere inversions (LTIs) to understand the temperature inversions at several atmospheric layers. In general, the inversions in HCM City showed the highest frequency at the air layers between 300–2,000 m (i.e., the EIs), followed by the lower tropospheric layer above 2,000 m (i.e., the LTIs) and the surface layer lower than 300 m (i.e., the SIs). The annual frequencies of the SI, EI, and LTI were 8%, 49%, and 42%, respectively. Regarding hourly variation, mean frequencies of the SI, EI, and LTI at 00Z (7 AM local time) were 7%, 50%, and 43%, respectively. Those at 12Z (7 PM local time) were 10%, 49%, and 41%, respectively. The statistically significant difference in the inversion frequency at two-time points (i.e., 00Z and 12Z) was only observed for the SIs (i.e., near surface inversions) (t-test, $P = 0.008$).

191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201


Noticeably, the SIs at 12Z (7 PM local time) were more frequent in the rainy season (Fig. 4b). This observation would be because rain events in the rainy season of HCM City tend to occur from the afternoon until the evening [26]. After the rain events, the air near the ground is cooler than the air in the higher atmospheric layer, causing a high frequency of SIs (i.e., surface inversions) at 12Z (7 PM local time) in the rainy season. Our finding on the occurrence of surface inversions after the rain events is supported by the similar observations in Kolkata of India [25] which has the similar climate to HCM City. However, these SI tends to disappear in the morning when the ground is heated by solar radiation [3], resulting in a low frequency of the SIs at 00Z (7 AM local time) in the rainy season (Fig. 4a).



202
203
204
205
206
207
208
209
210
211
212
213
214
215
216

Fig. 4 Frequency of the surface inversions, elevated inversions, and low-tropospheric inversions at (a) 00Z (7 AM local time) and (b) 12Z (7 PM local time) in the dry and rainy seasons of Ho Chi Minh City.

The inversion strengths, referred to as the change in atmospheric temperature over the first 100 m height from ground level [18], in the dry and rainy seasons of HCM City is shown in Fig. 5. In general, hourly and seasonal variations were observed for the temperature strength. Particularly, at 00Z (7 AM local time), the temperature inversions of the dry season (0.011 ± 0.005 °C/m) were 2.3 times stronger than those of the rainy season (0.005 ± 0.002 °C/m). The greater inversion strength in the dry season would possibly prevent the vertical mixing of air pollutants [27], leading to a trap of pollutants, such as $PM_{2.5}$, near the ground surface. The relationship between the temperature inversions and $PM_{2.5}$ concentrations in HCM City is more discussed in Section 3.3.



217
218
219
220
221
222
223

Fig. 5. Inversion strength observed at 00Z (7 AM local time) and 12Z (7 PM local time) in the dry and rainy seasons of Ho Chi Minh City, Vietnam.

A reverse trend was observed for the inversion strength at 12Z (7 PM local time), in particular, the rainy season showed a stronger inversion (0.013 ± 0.001 °C/m) compared to the dry season (0.009 ± 0.001 °C/m). As mentioned, rain events in the rainy season of HCM

224 city tend to last from the afternoon to the evening [26]. The occurrence of rain could cause
 225 the air near the surface to be cooler than the air in the above layers, expanding the
 226 temperature difference between the air layers and resulting in a stronger inversion at 12Z in
 227 the rainy season (Fig. 5).

228

229 3.3 Effect of temperature inversion on PM_{2.5} concentrations

230

231 To identify the effect of the temperature inversion on PM_{2.5} concentrations, the study period
 232 was classified into inversion and non-inversion days depending on the occurrence of the
 233 temperature inversion on a given day. To evaluate the effect of the temperature inversions
 234 on the PM_{2.5} concentrations, only the inversions below 300 m height (i.e., the SIs) were
 235 considered because the PM_{2.5} concentrations were monitored near the surface layer. The
 236 PM_{2.5} concentrations on the inversion and non-inversion days in the dry and rainy seasons
 237 are illustrated in Fig. 6. As shown in this figure, the mean PM_{2.5} concentration on the
 238 inversion days (dry season: 27.2 ± 9.1 µg/m³, rainy season: 22.2 ± 4.9 µg/m³) were 1.2 times
 239 higher than those on the non-inversion days (dry season: 23.0 ± 8.9 µg/m³, rainy season:
 240 18.4 ± 4.2 µg/m³), reflecting a contribution of the temperature inversion to an increase of
 241 PM_{2.5} concentrations in HCM City. This suggestion is supported by the significantly positive
 242 correlation between PM_{2.5} concentrations and inversion frequency, especially in the dry
 243 season (Table 1).

244

245 **Table 1. Spearman correlation between PM_{2.5} concentrations and inversion frequency**
 246 **in the dry and rainy seasons.**


247

	PM _{2.5} (dry season)	PM _{2.5} (rainy season)
Inversion strength	0.2507	-0.3859
Inversion frequency	0.7004*	0.2362
Pressure	0.5718	0.4750
Wind speed	-0.8186*	-0.7960
Potential temperature	-0.3657	-0.4737

248

249

*: correlation is significant at the 0.05 level.



250

251

252

253

254

255


256

257

Fig. 6. Concentrations of PM_{2.5} on the inversion and non-inversion days in the dry and rainy seasons of Ho Chi Minh City.

Additionally, the monthly PM_{2.5} concentrations on the inversion and non-inversion days and the monthly inversion frequency are shown in Fig. 6. The higher inversion frequencies were observed in several months of the dry season, including November, December, January, and February. Added to this, the concentrations of PM_{2.5} on the inversion days of these

258 months increased markedly and exceeded the WHO and Vietnamese air quality guidelines
 259 for annual $PM_{2.5}$ levels (5 and $25 \mu\text{g}/\text{m}^3$, respectively) (Fig. 7). This finding suggests that the
 260 temperature inversions would affect the concentrations of $PM_{2.5}$ in HCM City and this
 261 influence could be more considerable in the dry season when the inversions were stronger
 262 and more frequent (Figs. 3 and 4).
 263



264
 265
 266
 267
 268

Fig. 7. Monthly concentrations of $PM_{2.5}$ on the inversion and non-inversion days. The solid line illustrates the inversion frequency in each month.

269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281

In addition, the temperature inversions discourage the vertical mixing of $PM_{2.5}$ [5], causing this pollutant to be horizontal dispersed in the air [28] and resulting in an increase of $PM_{2.5}$ concentrations near the ground surface. Thus, the disappearance of temperature inversions promoted the vertical dispersion of $PM_{2.5}$ and led to a decrease in $PM_{2.5}$ concentrations on the non-inversion days. Regarding the rainy season, the $PM_{2.5}$ concentrations did not show a significant correlation to the inversion frequency and strength (Table 1), suggesting that other meteorological parameters, such as rainfall amount and rainfall frequency, rather than the temperature inversions could importantly impact the $PM_{2.5}$ concentrations in the rainy season. The wet deposition during the rain events would possibly remove particles in the air [25] and would mainly lower the $PM_{2.5}$ concentrations in the rainy season. In addition, the stronger wind speed in the rainy season (e.g., August to October) (Fig. 3) would enhance the dispersion of air pollutants and reduce the $PM_{2.5}$ concentrations.

282
 283
 284

4. CONCLUSION

285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297

In this study, temporal variation of the temperature inversions and their effect on the $PM_{2.5}$ concentrations in a metropolis having a tropical monsoon climate in Vietnam were investigated. For the $PM_{2.5}$ concentrations, the dry season (i.e., November to April of the following year) showed higher concentrations compared to the rainy season (i.e., May to October). The surface inversions, which are temperature inversions below 300 m, were more frequent at 12Z (7 PM local time) in the rainy season. This observation could be stemmed from the cooler air near the ground after rain events, which tend to last from the afternoon to the evening in the rainy season of HCM City.

The results also revealed a stronger inversion in the morning of the dry season, which could prevent the vertical mixing of air pollutants and lead to air pollution. This suggestion was supported by the higher $PM_{2.5}$ concentrations observed on inversion days and the significantly positive correlation between $PM_{2.5}$ concentrations and inversion frequency,

298 especially in the dry season. For the rainy season, the temperature inversions were believed
299 to have a minor effect on the PM_{2.5} concentrations since no significant correlation between
300 the PM_{2.5} concentrations and inversion frequency and strength was found for this season.
301 Based on the results of this study, solutions for reducing the PM_{2.5} pollution in the study area
302 can be obtained, such as the development of public transport to reduce the vehicle density
303 and a limitation of private vehicles in the dry season when the stronger temperature
304 inversions were observed.

305

306 **COMPETING INTERESTS**

307

308 Authors have declared that no competing interests exist.

309

310 **REFERENCES**

311

312 1. H. R. Phillpot and J. W. Zillman, The surface temperature inversion over the Antarctic

313 Continent, *J. Geophys. Res.*, 1970, **75**, 4161–4169.

314 2. C. D. Whiteman, Breakup of Temperature Inversions in Deep Mountain Valleys: Part I.

315 Observations, *J. Appl. Meteorol. Climatol.*, 1982, **21**, 270–289.

316 3. J. W. Enz, V. Hofman and A. Thostenson, *Air temperature inversions: Causes,*

317 *chracteristics and potential effects on pesticide spray drift*, 2019.

318 4. T. T. Trinh, T. T. Trinh, T. T. Le, T. D. H. Nguyen and B. M. Tu, Temperature inversion

319 and air pollution relationship, and its effects on human health in Hanoi City, Vietnam,

320 *Environ. Geochem. Health*, 2019, **41**, 929–937.

321 5. J. Wallace, D. Corr and P. Kanaroglou, Topographic and spatial impacts of temperature

322 inversions on air quality using mobile air pollution surveys, *STOTEN.*, 2010, **408**, 5086–

323 5096.

324 6. K. F. G. Olofson, P. U. Andersson, M. Hallquist, E. Ljungström, L. Tang, D. Chen and J.

325 B. C. Pettersson, Urban aerosol evolution and particle formation during wintertime

326 temperature inversions, *Atmos. Environ.*, 2009, **43**, 340–346.

327 7. S. D. Chambers, F. Wang, A. G. Williams, D. Xiaodong, H. Zhang, G. Lonati, J. Crawford,

328 A. D. Griffiths, A. Ianniello and I. Allegrini, Quantifying the influences of atmospheric

329 stability on air pollution in Lanzhou, China, using a radon-based stability monitor, *Atmos.*

330 *Environ.*, 2015, **107**, 233243.

331 8. M. Coccia, The effects of atmospheric stability with low wind speed and of air pollution on

332 the accelerated transmission dynamics of COVID-19, *Int. J. Environ. Stud.*, 2021, **78**, 1-

333 27.

334 9. N. Reiminger, X. Jurado, J. Vazquez, C. Wemmert, N. Blond, M. Dufresne and J. Wertel,

335 Effects of wind speed and atmospheric stability on the air pollution reduction rate induced

336 by noise barriers, *J. Wind Eng. Ind. Aerodyn.*, 2020, **200**, 104160.

337 10. Y. F. Xing, Y. H. Xu, M. H. Shi and Y. X. Lian, The impact of PM_{2.5} on the human

338 respiratory system, *J. Thorac. Dis*, 2016, **8**, E69–74.

339 11. E. Givental, Ho Chi Minh City: Contested public and private space in the Vietnamese

340 metropolis, *Focus Geogr.*, 2013, **56**, 32–44.

341 12. Q. B. Ho, H. N. K. Vu, T. T. Nguyen, T. T. H. Nguyen and T. T. T. Nguyen, A combination

342 of bottom-up and top-down approaches for calculating of air emission for developing

343 countries: a case of Ho Chi Minh City, Vietnam, *Air Qual. Atmos. Health*, 2019, **12**, 1059–

344 1072.

345 13. T. T. Q. Nguyen, W. Takeuchi, P. Misra and S. Hayashida, Technical note: Emission

346 mapping of key sectors in Ho Chi Minh City, Vietnam, using satellite-derived urban land

347 use data, *Atmos. Chem. Phys.*, 2021, **21**, 2795–2818.

348 14. V. Q. Doan and H. Kusaka, Projections of urban climate in the 2050s in a fast-growing

349 city in Southeast Asia: The greater Ho Chi Minh City metropolitan area, Vietnam, *Int. J.*

350 *Climatol.* 2018, **38**, 4155–4171.

- 351 15.K. Shukla and S. G. Aggarwal, A Technical Overview on Beta-Attenuation Method for the
352 Monitoring of Particulate Matter in Ambient Air, *Aerosol Air Qual. Res.*, 2022, **22**, 220195.
- 353 16.X. Feng, S. Wei and S. Wang, Temperature inversions in the atmospheric boundary layer
354 and lower troposphere over the Sichuan Basin, China: Climatology and impacts on air
355 pollution, *STOTEN.*, 2020, **726**, 138579.
- 356 17.C. D. Whiteman and T. B. McKee, Breakup of Temperature Inversions in Deep Mountain
357 Valleys: Part II. Thermodynamic Model, *J. Appl. Meteorol. Climatol.*, 1982, **21**, 290–302.
- 358 18.EPA, *Discussion paper: Validation of inversion strength estimation method*, 2014.
- 359 19.C. R. Harris, K. J. Millman, S. J. van der Walt, R. Gommers, P. Virtanen, D. Cournapeau,
360 E. Wieser, J. Taylor, S. Berg, N. J. Smith, R. Kern, M. Picus, S. Hoyer, M. H. van
361 Kerkwijk, M. Brett, A. Haldane, J. F. del Río, M. Wiebe, P. Peterson, P. Gérard-Marchant,
362 K. Sheppard, T. Reddy, W. Weckesser, H. Abbasi, C. Gohlke and T. E. Oliphant, Array
363 programming with NumPy, *Nature*, 2020, **585**, 357-362.
- 364 20.W. McKinney, presented in part at the The 9th Python in Science Conference (Scipy
365 2010), 2010.
- 366 21.WHO, *WHO global air quality guidelines: particulate matter (PM_{2.5} and PM₁₀), ozone,
367 nitrogen dioxide, sulfur dioxide and carbon monoxide*, 2021.
- 368 22.MONRE, *National Technical Regulation on Ambient Air Quality (QCVN05:2013/BTNMT)*
369 *(in Vietnamese)*, MONRE (Ministry of Natural Resources and Environment), 2013.
- 370 23.T. T. Hien, N. D. T. Chi, N. T. Nguyen, L. X. Vinh, N. Takenaka and D. H. Huy, Current
371 status of fine particulate matter (PM_{2.5}) in Vietnam's most populous city, Ho Chi Minh City,
372 *Aerosol Air Qual. Res.*, 2019, **19**, 2239–2251.
- 373 24.B. Q. Ho, Modeling PM₁₀ in Ho Chi Minh City, Vietnam and evaluation of its impact on
374 human health, *Sustain. Environ. Res.*, 2017, **27**, 95–102.
- 375 25.S. Talukdar, S. Jana and A. Maitra, Variation of black carbon concentration associated
376 with rain events at a tropical urban location, *Curr. Sci.*, 2014, **107**, 72–78.
- 377 26.H. C. M. C. P. s. Committee, *Report on climate evaluation of Ho Chi Minh City*, 2021.
- 378 27.A. Al-Hemoud, M. Al-Sudairawi, M. Al-Rashidi, W. Behbehani and A. Al-Khayat,
379 Temperature inversion and mixing height: critical indicators for air pollution in hot arid
380 climate, *Nat. Hazards*, 2019, **97**, 139–155.
- 381 28.X. Li, X.-M. Hu, Y. Ma, Y. Wang, L. Li and Z. Zhao, Impact of planetary boundary layer
382 structure on the formation and evolution of air-pollution episodes in Shenyang, Northeast
383 China, *Atmos. Environ.*, 2019, **214**, 116850.
- 384