

Microbial **contamination** and carcass quality of cod *Gadus morhua* obtained from local markets of Imo state

Abstract

The microbial **contamination** of cod *Gadus morhua* obtained from local markets of Imo state revealed that the total viable count for the fish species sampled from cold room ranged from $5.3 \times 10^4 - 9.5 \times 10^4$ cfu/g while that of the open market samples ranged from 8.0×10^4 cfu/g to 2.8×10^5 cfu/g. Total Fungal Count (TFC) had increasing microbial values in the open markets than in the cold rooms and ranged from 1.0×10^3 cfu/g to 7.2×10^3 cfu/g for cold room samples whereas a range of 1.0×10^5 cfu/g to 3.4×10^5 cfu/g was obtained for open markets. Generally, nine bacterial species and nine fungal species were isolated from the cold room and open market samples, including *Micrococcus spp.*, *Klebsiella spp.*, *Pseudomonas aeruginosa*, *Mucor spp.*, *Rhodotorula spp.*, *Aspergillus spp.* among others. Result for proximate composition shows that crude protein cold room value was 17.24 ± 0.52 but significantly increased in the open market samples (19.44 ± 0.94) just as there was significant difference in crude fiber values of cold room (6.98 ± 0.54) and open markets (ranged between 3.04 ± 0.81 and 4.39 ± 1.30). On the other hand, an insignificant increase in values of moisture content, NFE, ash content and ether extract but with values reducing from cold room (1.30 ± 0.62 and 1.06 ± 0.50) to open market (0.12 ± 0.70 and 1.00 ± 1.00) as their minimum value for the later.

Key words: *Gadus morhua*, microbial **contamination**, proximate composition, cold-room, open-market, Imo state

Introduction

Fish, a commodity of high importance, is highly perishable and unless correctly treated after harvesting, can soon become unfit to eat and possibly dangerous to

health through microbial growth, chemical change and breakdown of endogenous enzymes (FAO, 2014). Fish is one of the food commodities that quickly change form by spoilage if not appropriately exposed to a form of processing (Nwazuo *et al.*, 2016). Despite the great importance attributed to fish consumption, it is often overlooked when sustainable and safe food are discussed. Previous studies indicates that the importation of frozen fish plays a key role in bridging the gap between the demand for and supply of fish from domestic sources (Teslim, 2010). This demand for high quality, safe and healthy fish is now on a global increase (Huss, Ababouch and Gram, 2003 & Sen, 2005) yet, low income earning countries like Nigeria preferably export high market value products and import low market value products for domestic consumption as previously stated by Asche *et al.*, (2015).

Gadus morhua (cod) is benthopelagic lean fish, distributed in the North Atlantic and in the Arctic and abundant in cold and temperate shelf areas (FAO, 2018). Cod is one of the top imported fishes in Nigeria with high demand necessitating its importation in its various sizes and variously preserved forms especially, frozen form. Global capture production of cod in 2020 exceeds 1000 kilo tonnes which is nothing to be compared to aquaculture production rated at around 662 kilo tonnes of the same year (FAO 2022). External uncontrolled factors are likely to affect bacterial composition in the wild than in the controlled products. In addition, the removed function of the fish immune system at death allows for bacteria and enzymes to penetrate the walls of the tissues leading to spoilage of the tissue. This action of bacteria is further enhanced by the exposure of fish to unfavourable conditions during storage and handling (Sallam, 2007).

Fish constitute a major delicacy during all occasions in the south eastern part of Nigeria especially the frozen form as reported by Agom, *et al.*, (2012) who noted that some fish like croaker and mackerel has its most supply during the festive season and off the market or scarce during non-festive seasons. With the fish high rate of consumption and the need to avert consequential effects of

consuming non-safe fish of low nutritional value, this paper indirectly instigates the need for improved policies on import and preservation methods, buttress the need for diversification into local production and consumption of fresh live fish and inform the government on the need to put more effort into ensuring that fish sold within Imo state and Nigeria as a whole are safety risk free. This paper therefore exposes the microbial composition and carcass quality of *Gadus morhua* (cod) which is a commonly consumed frozen fish in Imo state.

Materials and methods

Study Area and Experimental Design

The study was carried out in Imo state located in South Eastern Nigeria, constituting of twenty seven local government areas and subdivided into three zones (Imo west- Orlu zone, Imo north- Okigwe zone and Imo East- Owerri zone). Most persons in this region are Christians with almost no phobia for fish. A total of eighty (80) samples were used for the experiment constituting eight samples each of the *Gadus morhua* (cod) collected from a cold room (a major distributor) and three local fish open markets of each of the three representing local government areas of the zones in Imo state- Orlu zone, Owerri zone and Okigwe zone. Representing Orlu zone were three local markets selected from Ohaji Egbema LGA, Owerri Zone was represented by local markets from Mbaitoli LGA while Okigwe zone was represented by local markets from Isiala Mbano LGA.

For experimental purposes, fish samples were coded based on place of collection (Table 1). This served to keep information discrete and remove subjective judgement of samples.

Local Government Areas	Codes		
	Cold room	Open markets	
		I	II

Owerri Municipal	AC				Table 1; Code
Ohaji		IO₁	IO₂	IO₃	
IsialaMbano		AO₁	AO₂	AO₃	
Mbaitolu		AnO₁	AnO₂	AnO₃	

allocation for the place of fish sample collection

Where AC.... Cold room

IO_{1,2,3} – Open market I, II, III in Ohaji

AO_{1,2,3} ... Open market I, II, III in IsialaMbano

AnO_{1,2,3}... Open market I, II, III in Mbaitolu

Fish Identification and Transportation

The fishes were identified with the aid of standard fish identification keys by Reed *et al.*, (1967), Loveque *et al.*, (1991) and FAO (1990) Field identification Guide to the commercial marine resources of the Gulf of Guinea. The length, weight and temperature of the fish samples were recorded and the fish carefully put in well labelled sterile polyethene bags, stocked inside iced boxes and quickly transported to the laboratory for processing and analysis. Temperature close to that of the cold rooms was maintained in the iced boxes using ice blocks.

Microbial Analysis

Microbes were isolated from the skin, tissue, gills and intestines of the sample fishes. Media used for inoculation included Nutrient agar (all-purpose bacteriological media) at 2.8g/100ml, Macconkey agar (for enumeration of coliforms) at 4.9g/100ml, Saboraud Dextrose agar (for enumeration of fungi) at

6.5g/100mls, Salmonella Shigella agar (for enumeration of Salmonella and Shigella) 6.4g/100mls, Manitol Salt agar (for Staphylococcus spp.) at 11.1g/100mls and Centrimide agar (for enumeration of Pseudomonas sp.) at 4.8g/100mls. Media were poured aseptically, samples serially diluted adapting 10-fold serial dilution and samples inoculated by spread plate method. Bacteriological plates were incubated at 37⁰C for 24 hours while mycological plates incubated at 27⁰C (room temperature) for 48 hours.

Proximate Composition of Carcass Quality

Samples of each test fish carcass was analysed chemically in accordance with the Official Analytical Chemists (AOAC, 2005) for such parameters as crude protein, moisture and dry matter, ash, crude fibre, crude fat (or lipid), carbohydrate, caloric value and Nitrogen free extract (NFE) to ensure that the fish meat meet the basic nutritional requirements of consumers. This analysis was carried out at the Food Science and Technology laboratory of the Federal University of Technology Owerri and subsequently at National Root Crops Research Institute Umudike, Abia state. The routine semi micro-kjeldahl procedure was used to determine crude protein, whereas moisture content was determined with a modified version of the AOAC 967-08 (AOAC 2005). The NFE was determined by difference. This was done by subtracting the sum of (% moisture + % crude protein + % ether extract + % crude fiber + % ash) from 100 i.e. $100 - (\%M + \% CP + \% EE + \% CF + \% Ash)$.

Comparative Statistical Analysis

The experiment was designed as a completely randomized experiment (CRD). Consequently the resulting data compared with one way analysis of variance (ANOVA) as described by Steel and Torrie (1996) and Njoku et al., (2014). The observed significant difference in mean values was separated using the LSD multiple range test. The analysis employed the computer statistical package for Social Sciences (SPSS) version 20 window 7.

Results and discussion

Fish, in the form of *Gadusmorhua*, a necessary commodity that bridges the gap of protein supply and hunger alleviation is readily available in its frozen form to meet almost all occasions in Imo state, showed varying levels of increasing microbial content and load as fish moved to the open markets from the cold rooms.

The result of the microbial and fungal loads/occurrence on the fish species from cold room and open markets together with the bacterial and fungal isolates are presented in the tables below;

Table 2a; Mean microbial load of cold room frozen *Gadusmorhua* samples collected from Owerri market

Site of collection	TVC (Cfu/g)	TCC (Cfu/g)	TSC (Cfu/g)	TPC (Cfu/g)	TSSC (Cfu/g)	TFC (Cfu/g)
Skin	6.7×10^4	3.7×10^4	1.3×10^4	4.0×10^2	-	7.2×10^3
Tissue	9.5×10^4	6.0×10^3	6.8×10^3	6.0×10^2	-	1.0×10^3
Gill	8.5×10^4	-	4.0×10^3	-	-	8.2×10^3
Intestine	5.3×10^4	2.0×10^4	1.1×10^3	-	4.0×10^2	2.2×10^3

Table 2b; Mean microbial load of open market frozen fish samples collected from Ohaji LGA markets

Site of collection	TVC (Cfu/g)	TCC (Cfu/g)	TSC (Cfu/g)	TPC (Cfu/g)	TSSC (Cfu/g)	TFC (Cfu/g)
Skin	1.5×10^5	8.6×10^4	6.2×10^5	3.1×10^3	5.6×10^4	3.4×10^5
Tissue	1.4×10^5	2.5×10^4	4.0×10^5	2.8×10^4	9.5×10^4	2.0×10^5
Gill	2.8×10^5	9.4×10^3	1.8×10^5	3.8×10^4	5.7×10^5	2.6×10^5
Intestine	1.2×10^5	-	2.8×10^6	-	-	1.8×10^5

Table 2c; Mean microbial load of open market frozen fish samples collected from IsialaMbano LGA markets

Site of collection	TVC (Cfu/g)	TCC (Cfu/g)	TSC (Cfu/g)	TPC (Cfu/g)	TSSC (Cfu/g)	TFC (Cfu/g)
Skin	9.0×10^4	8.4×10^4	2.9×10^5	2.8×10^3	4.8×10^4	3.5×10^5
Tissue	1.0×10^5	1.5×10^4	2.0×10^5	1.7×10^4	6.0×10^4	1.6×10^5
Gill	1.8×10^5	1.4×10^4	1.3×10^5	2.5×10^4	4.4×10^5	1.8×10^5
Intestine	8.0×10^4	-	2.0×10^6	-	-	1.0×10^5

Table 2d; Mean microbial load of open market frozen fish samples collected from Mbaitoli LGA markets

Site of collection	TVC (Cfu/g)	TCC (Cfu/g)	TSC (Cfu/g)	TPC (Cfu/g)	TSSC (Cfu/g)	TFC (Cfu/g)
--------------------	-------------	-------------	-------------	-------------	--------------	-------------

Skin	1.2×10^5	1.3×10^5	4.4×10^5	2.4×10^4	4.6×10^4	3.0×10^5
Tissue	1.2×10^5	2.0×10^4	3.0×10^5	2.0×10^4	8.5×10^4	1.8×10^5
Gill	2.0×10^5	1.9×10^4	2.0×10^5	2.7×10^4	4.9×10^5	1.0×10^5
Intestine	1.0×10^5	-	2.4×10^6	-	-	1.4×10^5

Where TVC- total viable count, TCC- total coliform count, TSC- total Staphylococcal count, TPC- total pseudomonas count, TSSC- total Salmonella/Shigella count, TFC- Total fungal count and CfU/g- colony forming unit per gram

The total viable count for the fish species sampled from cold room (table 2a) ranged from 5.3×10^4 – 9.5×10^4 cfu/g with the most overall viable count as obvious in the tissue and the least from the intestine. TVC of the open market samples however ranged from 8.0×10^4 cfu/g to 2.8×10^5 cfu/g with samples obtained from Ohaji LGA open markets showing the most viable count as samples from the gills revealed the most TVC. Cold room samples had Total coliform count (TCC) that ranged from not being detected in the gills to 3.7×10^4 cfu/g obtained from the skin. On the other hand, open market samples showed varying TCC ranges from being totally absent (in intestinal samples) to as high as 8.6×10^4 cfu/g, 8.4×10^4 cfu/g and 1.3×10^5 cfu/g in samples of the skin from Ohaji, Mbano and Mbaitoli respectively. Total Staphylococcal count (TSC) likewise recorded increasing values from cold room to open markets with the highest TSC value obtained from the skin (1.3×10^4 cfu/g) of the cold room and from the intestine (2.8×10^6 cfu/g) of the open market sample. Emikpeet *et al.*, 2011 noted that the means of contamination and route of microbial transmission are likely to be through handling. Onsite contamination of fish during transportation, display and handling as reported by Nwazuo *et al.*, 2016; Adams and Moses 2000, explains the increase in microbes as the fish proceeds from cold rooms to the open markets. Of note is the gill isolation of the cold room samples which had isolations (2.0×10^4 cfu/g) but absent in the open market samples. Same is recorded for the Total Salmonella/Shigella Count (TSSC) which were present (4.0×10^2 cfu/g) in cold room samples but entirely absent in the open market samples. Total Fungal Count (TFC) showed increasing values

from cold room with a range of 1.0×10^3 cfu/g - 7.2×10^3 cfu/g to open markets samples with a range of 2.0×10^5 cfu/g – 3.5×10^4 cfu/g the latter obtained from the skin.




Fig 1; Bar chart showing compared microbial indices for cold room and open market *Gadusmorhua* iced fish.

Comparison of mean microbial composition of cold room and open market frozen fish samples

Table 3; Comparative analysis of mean microbial load in cold room and open markets iced fish in Imo state

Sample	Parameters (Cfu/g)	Cold room ($\times 10^5$)	Ohaji ($\times 10^5$)	IsialaMbano ($\times 10^5$)	Mbaitoli ($\times 10^5$)	p-value
<i>Gadusmorhua</i>	TVC	0.75 ± 0.18	1.73 ± 0.72	1.13 ± 0.45	1.35 ± 0.44	0.089
	TCC	0.15 ± 0.16	0.30 ± 0.38	0.28 ± 0.37	0.42 ± 0.59	0.838
	TSC	0.06 ± 0.05	9.95 ± 12.16	6.62 ± 8.94	8.57 ± 10.33	0.462
	TPC	0.00 ± 0.00	0.17 ± 0.18	0.11 ± 0.12	0.17 ± 0.12	0.226
	TSSC	0.00 ± 0.00	1.80 ± 2.62	1.37 ± 2.03	1.55 ± 2.25	0.601
	TFC	0.05 ± 0.03	2.45 ± 0.71 ^a	1.97 ± 1.07 ^b	1.80 ± 0.86 ^c	0.005

Table 4; Comparative analysis of average occurrence of **bacteria in *Gadusmorhua*** in cold room and open markets of Imo state.

Bacterial isolates	Maximum occurrence	Cold room	Open market			p-value
			Ohaji	IsialaMban o	Mbaitoli	
<i>Micrococcus spp.</i>	16	16.00 ± 0.00	11.00 ± 2.65	10.00 ± 5.19	10.00 ± 2.00	0.532
<i>Staphylococcus aureus</i>	16	13.00 ± 0.00	14.67 ± 1.52	13.33 ± 0.57	13.33 ± 0.57	0.340
<i>Bacillus cereus</i>	16	8.00 ± 0.00	5.00 ± 1.00	3.33 ± 1.15	4.00 ± 2.00	0.130
<i>Bacillus subtilis</i>	16	7.00 ± 0.00	11.00 ± 1.00	10.33 ± 1.52	10.67 ± 3.55	0.452
<i>Streptococcus spp.</i>	16	13.00 ± 0.00 ^a	7.33 ± 1.15 ^b	6.00 ± 2.00 ^c	6.00 ± 1.00 ^d	0.024
<i>Escherichia coli</i>	16	6.00 ± 0.00	7.67 ± 1.52	6.67 ± 1.52	7.33 ± 2.08	0.087
<i>Klebsiella spp.</i>	16	3.00 ± 0.00	4.00 ± 1.00	3.33 ± 0.58	3.33 ± 0.57	0.586
<i>Proteus spp.</i>	16	3.00 ± 0.00	4.00 ± 1.00	3.67 ± 1.15	3.33 ± 1.52	0.705
<i>Pseudomonas aeruginosa</i>	16	7.00 ± 0.00	10.67 ± 2.51	10.67 ± 4.72	10.67 ± 4.50	0.861
<i>Salmonella spp.</i>	16	4.00 ± 0.00	15.67 ± 0.57	14.67 ± 1.15	15.33 ± 1.15	0.000

Table 5; Comparative analysis of average occurrence of **fungi in *Gadusmorhua*** in cold room and open markets of Imo state.

Fungal isolates	Maximum occurrence	Cold room	Open market			p-value
			Ohaji	IsialaMban o	Mbaitoli	
<i>Geotrichum sp.</i>	16	4.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	0.00 ± 0.00	NA
<i>Candida spp.</i>	16	13.00 ± 0.00 ^a	0.33 ± 0.577 ^b	0.00 ± 0.00 ^c	0.33 ± 0.57 ^d	0.000
<i>Aspergillusfumigatus</i>	16	0.00 ± 0.0	4.00 ± 1.73	3.33 ± 1.52	3.33 ± 2.08	0.366
<i>Rhizopusstolonifer</i>	16	0.00 ± 0.00 ^a	8.33 ± 1.52 ^b	6.67 ± 1.52 ^c	8.00 ± 2.00 ^d	0.025
<i>Rhodotorula spp.</i>	16	0.00 ± 0.00	4.00 ± 1.00	3.33 ± 1.52	3.33 ± 1.52	0.193
<i>Aspergillusflavus</i>	16	0.00 ± 0.00	8.00 ± 2.64	5.67 ± 1.52	6.67 ± 1.52	0.062
<i>Mucor spp.</i>	16	0.00 ± 0.00	4.00 ± 1.73	2.67 ± 1.15	2.67 ± 1.52	0.240
<i>Aspergillusterreus</i>	16	0.00 ± 0.00	12.00 ± 3.60	10.33 ± 2.51	11.67 ± 3.21	0.068
<i>Aspergillusniger</i>	16	0.00 ± 0.00	4.00 ± 3.46	3.37 ± 2.08	4.00 ± 1.00	0.538

Results from the comparative analysis reveals that TFC values alone showed significance difference at p-value 0.005 across the cold room and open markets collection points. Comparatively, there is no significant difference in the bacterial occurrence of *Micrococcus spp.*, *S.aureus*, *B. cereus*, *B. subtilis*, *E. coli*, *Klebsiellasp*, *Proteus spp.*, and *P. aeruginosa* across cold room and open markets of the three local government areas with p-values of 0.532, 0.340, 0.130, 0.452, 0.087, 0.586, 0.705, and 0.861 respectively but significant difference in values of *Streptococcus spp.* and *Salmonella spp.* Just as was reported in Onyeaunulaet *al.*,2021, and Orji *et al.*, 2014, in their studies on frozen

fish marketed in Imo State and Abakaliki metropolis, similar bacteria were isolated. Values for *Micrococcus spp.*, *Bacillus cereus*, and *Streptococcus spp.* were higher in cold room samples (16.00, 8.00 and 13.00 respectively) than in open markets with 11.00, 5.00 and 7.33 (Ohaji), 10.00, 3.33 and 6.00 (IsialaMbano) and 10.00, 4.00 and 6.00 (Mbaitoli). This may be attributed to the fact that their growth may be significantly reduced at higher temperatures and lower pH which is evident in fish that have left the cold rooms to warmer temperatures and susceptible to other microbial contamination which will likely increase the pH. Onyeaenulaet al., 2021 observed same with *Scomberscombrus* obtained in Imo State local market. Decrease in occurrence of *Bacillus cereus* and *Streptococcus sp.* from cold room to open markets however negates the observations of Onyeaenulaet al., 2021 and may be attributed to the physiochemical composition of *Gadusmorhua*. Fungal occurrence showed a non-significant difference for *A. fumigatus*, *Rhodotorula spp.*, *A. flavus*, *Mucor spp.*, *A. terreus* and *A. niger* across the cold room and open market samples with p-value of 0.366, 0.193, 0.062, 0.240, 0.068 and 0.538 respectively (table 5). *Geotrichum sp.* occurred in only cold room samples but was absent in open markets just as *Candida sp.* and *R. stolonifer* presented statistical significance across cold room and open market samples. Similar observation of fungal isolations was made for *Scomberscombrus* as reported by Onyeaenulaet al., 2021. Of all the nine fungi isolated, *Geotrichum sp.* showed significant difference at $p \leq 0.05$. Adebayo-Tayo et al., 2012, Ibrahim et al., 2015 and Onyeaenulaet al., 2021 reported similar isolations with *Candida sp.* having the most average occurrence as observed in this study. Nishihara et al., 2008 also reported that despite disinfection of fish samples surface, its incubation for up to 14 days at room temperature still presented *Candida sp.* among few other fungi isolated. Isolates of the other fungi from the present study had no occurrence in cold room samples but significantly increased values in open market samples. This further buttress the fact that improper storage, transport and display facilities predispose

these food commodities to fungal growth (Gill and Barbosa, 2011; Adebayo-Tayo *et al.*, 2012)

Proximate composition for frozen fish sold in Imo state markets

Table 6; Proximate analysis for frozen *Gadus morhua* from cold room and open market showing mean values

Fish specie	Parameters	Cold room (AC)			Ohaji			IsialaMbano			Mbaitoli			p-value
		O1	O2	O3	O1	O2	O3	O1	O2	O3	O1	O2	O3	
<i>Gadus morhua</i>	CP	17.24 ±0.52	19.44 ±0.94	19.44 ±0.72	19.44 ± 7.05	19.34 ± 0.53	19.14 ±1.01	19.14 ±0.51	19.13 ±0.71	18.14 ±1.00	17.29 ±0.63	0.042*		
	M	72.31 ±0.57	72.88 ±0.58	73.15 ±1.05	73.27 ± 0.70	74.00 ± 1.06	74.09 ±1.05	73.09 ±0.58	73.66 ±0.91	73.63 ±0.53	71.63 ±0.51	0.671		
	A	1.30± 0.62	0.98± 0.63	0.12± 0.70	0.13± 0.99	0.84± 0.87	0.97± 0.83	0.87± 1.00	0.92± 0.72	0.91± 0.72	0.84± 0.58	0.611		
	EE	1.06± 0.50	1.05± 0.93	1.05± 0.50	1.05± 1.05	1.05± 0.61	1.05± 1.08	1.06± 1.31	1.05± 0.65	1.00± 1.00	1.04± 1.74	0.993		
	CF	6.98± 0.54	3.38± 0.86	4.37± 0.66	3.16± 0.50	3.80± 0.87	4.39± 1.30	4.39± 0.81	3.17± 0.81	3.06± 0.50	3.04± 0.81	0.001*		
	NFE	1.08± 0.60	2.28± 0.94	1.88± 0.92	2.95± 1.78	0.98± 0.73	0.36± 0.50	1.45± 0.73	2.08± 0.52	2.00± 1.11	1.91± 0.54	0.612		

Where CP= crude protein, M= moisture, A= ash, EE=ether extract, CF= crude fiber, NFE= nitrogen free extract p-values with “*” shows significant difference p<0.05




Fig 2: Bar chart comparing mean values for proximate evaluation of fishes sampled from cold room

Result shows that crude protein cold room value was 17.24 ± 0.52 but significantly increased in the open market samples (19.44 ± 0.94) but with a unique value of 17.29 ± 0.63 in one open market of Mbaitoli. The increase in

values of moisture content from the cold room (72.31 ± 0.57) to the open market (74.09 ± 1.05) was insignificant just as was recorded for NFE to be 1.08 ± 0.60 for cold room and 2.95 ± 1.78 as maximum value for open market. On the other hand, an insignificant difference was recorded for ash content and ether extract but with values reducing from cold room (1.30 ± 0.62 and 1.06 ± 0.50) to open market (0.12 ± 0.70 and 1.00 ± 1.00) as their minimum value respectively. Crude fiber showed significant difference in values of cold room (6.98 ± 0.54) and open markets (ranged between 3.04 ± 0.81 and 4.39 ± 1.30), thus a significant difference ($p < 0.05$) was observed in crude protein and crude fiber values of *Gadus morhua*. Proximate composition values obtained from this study agrees with those undertaken by Tawfik (2009) and Nurnadia *et al.*, (2011) with their nutritional properties acceptable especially for those obtained from the cold-rooms. Crude protein mean values indicates an increase in values in fish obtained from the open market to fish obtained from the cold room. The protein content of Atlantic mackerel in Olaolu *et al.*, (2011) supports this finding. Post rigor mortis is also associated in increased water retention capacity of proteins, thus may be responsible for the increased protein content in the open market.

Summary

The study looked at the microbial composition and carcass quality of *Gadus morhua* (a commonly sold and consumed fish) obtained from local markets of Imo state. An effective way of long term preservation of fish is by freezing (Nielsen and Flemming, 2012). The fluctuation of temperature is one of the most significant factors in maintaining quality through the chain of distribution. Just as it is important to quickly cool fish immediately it is caught to inhibit the onset of biochemical and microbiological change, it is equally important to maintain a relatively steady temperature as it is transported and made available to the final consumers. Results revealed that the difference in microbial and carcass quality of fresh and thawed fish may be negligible if fish

was handled with care during and after purchase from the cold rooms, kept frozen as quickly as possible and stored at a stable temperature of about 30°C even at the point of display for sale. This however is almost impossible with the handling habits found in our markets. Handlers and traders act as they deem fit depending on the limited resources available to them and market place involved. Small scale traders stay at cold room sites to share a carton of fish- and in so doing, start the exposure of the fish to microbial contamination accounting for the high levels of microbes in open market samples.

Conclusion

There is a marked recorded difference in microbial and carcass quality of fish in the cold rooms and those in the open markets available to vast majority of consumers. Most of these quality difference has been linked to the handling method used within the state and negligence of the use of best practices. Because frozen fish is exposed to large variations of temperature fluctuations, during storage and transport, short-term storage of the raw material, gentle processing, quick freezing, short-term storage as a semiprocessed product, and short-term cold storage at temperatures lower than 30°C, will result in the best product.

References

- Adams, M.R. and Moss, M.O. (2000). *Food Microbiology*. The Royal Society of Chemistry, Cambridge, 2nded.
- Adebayo-Tayo B. C., Odu N. N., Anyamele L. M., Igwiloh N. J. P. N. and Okonko I. O. (2012). Microbial Quality of Frozen Fish Sold in Uyo Metropolis. *Nature and Science*; 10(3): 71-77. <http://www.sciencepub.net>
- Agom, D. I., Etim, G. C. & Etuk, E. A., (2012). Analysis of wholesale frozen fish markets in Calabar, Cross River State, Nigeria. *Trends in Agricultural Economics*, 5(2): 61-69. Doi:10.3923/tae.2012.61.69
- AOAC (2005). *Official methods of Analysis of the Association of Official Analytical Chemists* (18thed), Maryland, USA, 1550 pp.
- Asche, F., Bellemare, M.F., Roheim, C., Smith, M.D. & Tveteras, S., (2015). Fair enough? Food security and the international trade of seafood. *World Dev.* 67, 151–160.

- Emikpe B. O., Adebisi T. and Adedeji O.B. (2011). Bacterial load on the skin and stomach of *Clarias Gariepinus* and *Oreochromis Niloticus* from Ibadan, South West Nigeria: Public health implications. *J. Microbiol. Biotech. Res.*, 1 (1): 52-59.
- FAO 2022. *Gadus morhua* Linnaeus, 1758. Fisheries and Aquaculture Division [online]. Rome. [Cited Thursday, July 7th 2022].
- FAO, (2014). *The State of World Fisheries and Aquaculture*. Food and Agricultural Organization, Rome, Italy.
- FAO, (2018). *The State of World Fisheries and Aquaculture 2018- Meeting the Sustainable development goals*. Rome.
- Food and Agriculture Organization (FAO), (1990). *Field guide to commercial marine resources of the Gulf of Guinea* RAFR/014/F1/90/02.
- Gil, M.M.; Barbosa, A.L. Microorganisms and safety. In *Practical Food and Research*, 3rd ed.; Cruz, R.M.S., Ed.; CRC Press: Boca Raton, FL, USA, 2011; pp. 195–217. <https://www.fao.org/fishery/en/aqspecies/2218/en>
- Huss, H. H., Ababouch, L., & Gram, L., (2003). *Assessment and Management of Seafood Safety and Quality* Rome. Food and Agriculture Organization of the United Nations. FAO Fisheries Technical Paper 444. p239
- Ibrahim A. S., Amr A. A., Youssef S. A. and Safaa M. E. (2015). Mycological Evaluation of Imported Frozen Fish. *Alexandria Journal of Veterinary Sciences*, 45:127-131
- Loveque, C. Pyugy, O. & Tengel G. G., (1991). The fresh and brackish water fishes of West Africa. Vol.1. Musu Royale de 1. *Afrique Centrale*, Tervurem, Belegue Edition de. 1. ORSTOM. 384pp.
- Nielsen J. and Flemming J. (2012). Quality of Frozen Fish. *Handbook of Meat, Poultry and Seafood Quality*. 1(2) pp577-587. Available at <https://doi.org/10.1002/9781118352434.ch31>
- M. Kamata, T. Koyama and K. Yazawa. 2008. New phospholipase A1-producing bacteria from a Marine Fish. *Marine Biotechnology*, 10: 382 - 387.
- Njoku, D. C. (2014). *Introduction to fisheries Technology*; A Fisheries course in Agriculture for Undergraduates. Martin Continental Publishers, Owerri, Nigeria, 43 pp.
- Nurnadia, A. A., Azrina A and Amin, I. (2011). Proximate Composition and Energetic Value of Selected Marine Fish and Shellfish from the West Coast of Peninsular Malaysia. *International Food Research Journal*, 18, 137-148
- Nwazuo N.I, Keke I.R, Egeruoh A.S, Nwanjo S.O and Ugoeze Q.C (2016) Sanitation and Handling during Processing of Fish in Oguta Fishing Communities and Its Economic Implications. *International Journal of Scientific & Engineering Research*, Volume 7, Issue 5, May-2016. Retrieved from <http://www.ijser.org>.
- Olaolu M. E., G.R. Akande; M.M Salaudeen; A.O. Olusola; R.S. Adeyemi and A.F. Babalola (2011). Technological properties and proximate

- composition of two imported fish species in Nigeria. *Electronic Journal of Environmental, Agricultural and food chemistry* 10 (6):2398-2403
- Onyeonula N. I., Egeruoh A. S., Azoro A. V., Nwanjo S. O. and Mbagwu C. B. (2021). Microbiological Evaluation and Public Health Implication of Frozen *ScomberScombrus* Sold in Selected Markets of Imo State, Nigeria. *The International Journal of Science &Technoledge*. Vol. 9(8): pp 44-51.
- Orji J. O., Okonkwo E. C., and Umezuluike O. I. (2014). Survey of the microbiological Quality of Frozen Fish marketed in Abakaliki Metropolis and its implications for consumer Health. *International Journal of scientific research*. 3 (8): 356-360.
- Reed, W., Burchard, J., Hopson, A. J., Jennes, J. &Yaro, I., (1967). *Fish and Fisheries of Northern Nigeria*. Ministry of Agriculture, Northern Nigeria, Zaira.
- Sallam K. I. 2007. Chemical, sensory and shelf life evaluation of sliced salmon treated with salts of organic acids. *Food Chem*. 2007; 101(2): 592–600.
- Sen D. P., (2005). *Advances in Fish Processing Technology*. Allied publishers, PVT Limited New Delhi.
- Steel RGD, Torrie JH, Dickey DA. (1996). Principles and procedures of statistics. A biometrical approach. 3rd ed. McGraw Hill Book Company Inc., New York, USA.
- Tawfik M. S. (2009). Proximate Composition and Fatty Acids Profiles in most Common Available Fish Species in Saudi Market. *Asian Journal of Clinical Nutrition*, 1, 50-57. Available at <http://docsdrive.com/pdfs/ansinet/ajcn/2009/50-57.pdf>
- Teslim, Bada, (2010). Market delineation study of the fish market in Nigeria; An application of co integration Analysis. *Journal of Agricultural Sciences*. 2(3): 101-112