

Original Research Article

Effects of Blood Meal Inclusion Diet on Growth Performance, Feed Utilization and Survival Rate of *Clarias gariepinus* (Burchell 1822) in Concrete Pond.

Abstract

The inclusion of blood meal as a substitution for commercial fishmeal in the practical fish diets was evaluated in *Clarias gariepinus* juveniles (initial weight: 11.45 ± 0.25 g and initial length: 9.00 ± 0.91 cm) for a period of 39 weeks. The proximate analysis of blood meal had 50.04%, 4.95%, 0.98%, 3.05%, 16.51% and 19.00% representing value of crude protein, lipid, fibre, ash, moisture and nitrogen free extract, respectively. Five isonitrogenous diets (40% crude protein) containing fishmeal which was replaced by blood meal at graded level of 0% as control, 25%, 50%, 75% and 100% were formulated. A total of 300 juveniles were used and randomly distributed into five treatments with each having triplicates of 20 fish per pond respectively. Fish were fed twice daily at 5% body weight in equal proportions at 8.00am and 4.00pm. At the end of the feeding trial, the results showed that growth performance were significant higher ($p < 0.05$) in fish fed control diet (1263.55g) and 25% BM diet (1068.67g), mean length gain (54.16cm and 50.85cm), protein efficiency ratio (31.59% and 24.22%) and percentage weight gain (11035.3% and 8476.03%) compared to other treatments diets with the lowest values of (675.35g, 46.86cm, 11.73% and 4190.63%, respectively) recorded in fish fed 100% BM diet. However, Feed conversion ratio, specific growth rate, average daily growth and survival rate were not significantly difference ($p > 0.05$) in all the dietary treatments. The water quality parameters monitored were within recommended ranged and were not affected by the supplemented diets. It can be concluded that inclusion of blood meal up to 25% can efficiently replace fish meal without any

Comment [ss1]: add

deleterious effects on growth, feed utilization and survival rate. The use of blood meal as an alternative protein source in diets of farmed fish species will reduce the cost of feeds; boost the profitability and sustainability of the fish production.

Keywords: *Clarias gariepinus*, protein, growth performance, blood meal

Introduction

Fish is an important source of rich protein food for human consumption and it has a major source of high-quality dietary essential amino acids, vitamins, minerals and other micro-nutrients. They are the cheapest source of animal protein which accounts for about 38% of Nigeria's total protein requirements (FAO, 2000 and Allan, 2004). Due to its tremendous value, it creates employments, recreation/sports, enhance national health and wealth.

The African catfish, *Clarias gariepinus* (Burchell, 1822) belong to the family *Clariidae* and a native fish species in African countries (Teugels, 1986). It is the most cultured fish species in Nigeria and other developing countries because of its fast growth performance, good feeding conversion, tolerates poor water quality conditions, feeds on agricultural by-products, hardiness and high commercial value (Adewolu *et al*, 2009; Aliu and Dako, 2018).

The interest of every fish farmers is to produce table-sized fish within the shortest possible time (Gabriele *et al*, 2007) with a balanced nutrient composition of the feed. However, feeding is one of the constraints facing fish farming in Nigeria due to its high cost of fish feeds; imported fish feed are very expensive and often scarce thereby reducing the growth performance, profitability and sustainability of the fish culture industry, since its cost constitutes about 60-70% of the production cost (Higgs *et al*, 1994). Protein is a major dietary nutrient component of formulated feeds (Wilson, 2002) and the commonest source is fishmeal which is very expensive. Fish meal is the conventional source of animal protein in

aqua feeds because of its high protein and amino acids profile, vitamins content, high digestibility and growth factors (Hardy and Tacon, 2002).

Due to high quest for protein source, numerous emphases on suitable alternative of various local protein sources has been put into consideration to ensure rapid increase in fish food production to meet the high population demand of fish consumers and aqua-farmers (Dawah *et al*, 2002; Adikwu, 1991 and FAO, 2016). According to Fagbenro (1999), feed cost is the major constraint in aquaculture and to reduce the price of a complete feed, locally available feedstuff should be included in the feed especially agricultural by-products. However, feedstuffs of animal protein origin which can easily be combined with other feed ingredients have attempt to partially or completely replace fish meal component in fish diets with varying degree of success without affecting the growth performance and survival rate since fish utilizes both animal and agricultural by-product (FAO, 2007; Adewolu, 2001; Gabriel *et al*, 2007 and NRC, 2011). They are cheaper, have complementary amino acid profiles, affordable and available in large commercial quantities.

Blood meal is a source of animal protein gotten from abattoirs that has high protein contents of (85%) and heme-iron (Otubusinet *al.*, 2009; FAO, 2012). It has been recognized as one of the most efficiently good substitute for fish meal. Furthermore, it is rich in lysine (7-9%), valine, leucine, phenylalanine and low methionine (Heuze *et al.*, 2015; Hertrampf and Piedad-pascual, 2000 and Agbebi *et al.*, 2009). Several studies have been reported on the potential use of blood meal to replace fish meal component in fish diets such as *Oreochromis niloticus* (Kirimiet *al.*, 2016; Aladetohun and Sogbesan, 2013 and Ogunji, 2001), gilthead sea bream (Nogueira *et al.*, 2012), mudfish fingerlings (Eyo and Olatunde, 1999), largemouth bass (Tidwell *et al.*, 2005) and *Clarias gariepinus* (Agbebi *et al.*, 2009 and Adejoke, 2012) with vary degree of successes. Blood meal provides a cheap and effective replacement for fish meal in fish diets, with no adverse effects on growth rate, survival and feed conversion ratio (Otubusinet *al.*, 2009).

Comment [ss2]: Grammatical error. Replace it.

Comment [ss3]: add

For fish to grow well, a well balanced diet containing the essential nutrient in the right proportions is needed by the fish and at the most economic profit to the farmer using a cheaper alternate protein source to replace the fish meal without compromising the quality, acceptability and palatability of the feed (Adikwu, 2003). Although blood meal has been recognized as a rich protein source and good substitute of fishmeal regarding its effects on growth, survival and yield of fish, there is need to conduct a feeding trial to determine the growth performance of fish this waste by-product. This research is aimed to evaluate the growth performance, feed utilization and survival rate, and also the optimum inclusion level of blood meal in the diets for juveniles of *Clarias gariepinus* in concrete pond.

Materials and methods

Study site:

The research study was conducted at Amas fish farm at Ezihe in Isiala-imbano, Local Government Area of Imo State. It is located in North-East of Imo State within Latitude $5^{\circ} 40' 4''$ N and Longitude of $7^{\circ} 12' 2''$ E. The fish ponds are made of concrete and designed in rectangular forms of (30cm×36cm×52cm) and the main source of water at the farm is bore-hole water.

Processing of blood meal and feedstuff

The blood was collected raw from the slaughter house at Isiala-mbano L.G.A. it was boiled for 30 minutes to reduce the water content and to congeal the blood. It was then oven-dried for 4 hours at a temperature of 105⁰C. The dried blood was then grounded finely using Victorian hand grinding machine. Fishmeal, soybean meal, maize, palm kernel cake and other feedstuffs purchased from commercial mill in Owerri, Imo State were separately milled, screened to fine and analyzed for the proximate composition (AOAC, 2005).

Comment [ss4]: add

Table 1: proximate composition of feed ingredients

Ingredients	Crude Protein (%)	Fiber (%)	Lipids (%)	Ash (%)	NFE (%)	Moisture (%)
Fish meal	58.04	0.72	10.00	10.71	14.24	11.33
Blood meal	50.10	0.98	4.95	3.05	19.00	16.51
Maize flour	10.80	5.78	4.28	6.58	23.30	10.42
S.B.M	48.74	5.10	6.18	4.48	12.03	10.70
P.K.C	30.73	3.60	6.50	4.03	9.52	11.52

S.B. = soya bean meal, P.K.C = palm kernel cake, NFE = nitrogen free extract.

Experimental diets

Base on feed ingredients proximate composition (table 1), five iso-nitrogenous diets containing (40% crude protein) were formulated using Pearson's square method with blood meal replacing fish meal component at 0% served as control, 25%, 50%, 75% and 100% inclusion level respectively (table 2). Prior to processing, the ingredients were milled individually to a fine powder, weighed according to formulation, manually mixed together and moistened thoroughly using the method described by De Silva and Anderson,

(1995). The mixed diets were pelleted using electrical pelletizing machine set at 2mm, 4mm and 6mm, sun-dried for 4days to remove moisture, packaged into separate bags, labeled and stored in a cool dry place. The experimental diets were analyzed in triplicate for the proximate composition (AOAC, 2005) to confirm the formulation (table 3).

Comment [ss5]: add the spaces properly

Table 2: ingredient composition of the experimental diets (g/100g)

Ingredients	Control (0%)	Diet 2 (25% BM)	Diet 3 (50% BM)	Diet 4 (75% BM)	Diet 5 (100% BM)
Fish meal	24.0	18.0	12.0	6.0	----
Blood meal	----	6.0	12.0	18.0	24.0
S.M.B	24.0	24.0	24.0	24.0	24.0
Maize flour	40.0	40.0	40.0	40.0	40.0
Bone meal	2.0	2.0	2.0	2.0	2.0
P.K.C	5.0	5.0	5.0	5.0	5.0
Starch (binder)	2.0	2.0	2.0	2.0	2.0
Salt	0.5	0.5	0.5	0.5	0.5
Vit/min premix	1.5	1.5	1.5	1.5	1.5
Palm oil	1.0	1.0	1.0	1.0	1.0
Total	100.0	100.0	100.0	100.0	100.0

S.M.B = soya bean meal, P.K.C = palm kernel cake. Vitamin/mineral premix supplied per gram of diet. Vitamin A: 20,000 IU; vitamin D₃: 4,000 IU; Vitamin E: 200,000 IU; Vitamin K: 1,200mg; Vitamin B: 10,000mg; VitaminB₂: 3,000mg; Vitamin K: 2.5mg; Vitamin B₁₂: 1,000mg; Niacin: 0.088mg; Folic acid: 5,00mg; Antioxidant: 3,000mg; panthothenic acid: 4,000mg; Biotin: 400mg; Riboflavin: 0.020mg; Vitamin C: 200mg; Iron: 0.300mg; Magnesium: 0.088mg; Manganese: 30mg; Copper: 4g; Zinc:40g; Iodine: 1,000mg; Calcium pantothenate: 11.5mg; Chlorine chloride: 500mg and Selenium:200mg.

Experimental Fish

Three hundred (300) Juveniles of *Clariasgariepinus* with mean body weight of approximately (11.45±0.25g) were used for the study and obtained from a private fish farm in Owerri, Imo State. The fish were transported in 50 liters open plastic rubber half filled

with water to the experimental site. The initial length and initial body weight of the fishes were measured with a meter ruler (to the nearest 0.1cm) and electronic weighing balance (Ohaus-30064419 scout pro, to the nearest 0.01gram) in order to establish the mean individual weight and length at the start of the experiment.

Experimental design and feeding

Fifteen concrete ponds (30cm×36cm×52cm) were used for the feeding trial. The experiment consisted of five treatments (diets) with three replicates. A total of three hundred (300) juveniles of *Clarias gariepinus* with an average weight (11.45 ± 0.25 g) and length (9.21 ± 0.91 cm) were randomly assigned in a completely randomized design (CRD) at a rate of twenty (20) juveniles per replicate. Prior to the experiment, the fish were acclimatized for seven days before the commencement of the experimental feeding in order to increase appetite and utilize the test diets. Fishes were fed at 5% of their total body weight daily (Madu et al., 2003). Feeding was administered twice daily between (8.00 – 8.30am and 4.00 – 4.30 pm) for the period of 39 weeks. Subsequently, growth data were taken fortnightly and quantity of feed fed adjusted in accordance with the fish new, also the leftover feeds together with fecal residues were siphoned out every three days siphoned out and replaced with fresh water described by Njoku (2010).

Growth and feed utilization parameters.

The weights of fish were taken every two weeks. The following growth performance and feed utilization analyzed, calculated and recorded for each treatment using the methods of (Adikwu, 2003; Burel et al., 2000, Ogunji, 2004 and Guroyet et al., 2005). These parameters includes; mean weight gain (MWG), specific growth rate (SGR), feed conversion ratio (FCR), Mean length gain (MLG), average daily growth (ADG), protein efficiency ratio

(PER), percentage weight gain (PWG), protein intake (PI), feed intake (FI) and survival rate (SR),

Mean weight gain (MWG) = final mean weight – initial mean weight (g)

Specific growth rate (SGR %) = $100(\log W_f - \log W_i) / T$

Where; W_1 and W_2 = the logarithms of initial and final mean weights of fish respectively,

Comment [ss6]: correct it as per above mentioned

T = the rearing period in days.

Feed conversion ratio (FCR) = total feed intake (g) / Fish weight gain (g)

Protein efficiency ratio (PER) = fish weight gain (g) / Protein intake (g)

Average daily growth (ADG) = mean weight gain (g) / No of days

Protein intake (PI) = total feed consumed × % protein in feed / 100

Total feed intake (TFI) = initial weight of feed – final weight of feed (g)

Survival Rate (SR) % = No of fish survived / Initial fish stocked × 100

Water quality parameters

Physico-chemical parameters in the various fish ponds were monitored during the experimental period early in the morning using the methods of Boyd, (1979) and APHA *et al*, (2005) prior to siphoning and feeding. The temperature was measured using mercury hand thermometer (Boyd, 1979), pH and dissolve oxygen were measured using Hanna digital multipurpose meter (H12210), and ammonia was measured using spectrophotometer (HACH/2000) to ensure the water quality standards for fish culture were met.

Chemical analysis

Proximate analysis of blood meal, feed ingredients, experimental diets and fish carcass (initial and final) were performed in triplicate according to the methods of AOAC (2005) at

Laboratory of Fisheries and Aquaculture, Federal University of Technology, Owerri, Imo State. Moisture content determined after drying samples in an oven for 4hrs at 105⁰ C until a constant weight was obtained. Crude protein (N×6.25) was determined using the routine semi-Kjeldahl method. Ash content was determined by igniting feed sample in a muffle furnace for 24hrs at 550⁰ C. Crude fat was determined by ether extraction using soxhlet method. Crude fiber was quantified by acid/base digestion followed by ashing the dry residue at 550⁰C in a muffle furnace for 4 hours. Nitrogen free extracts content was determined by subtracting the percentage of moisture, crude protein, ash and fat from 100.

Statistical analysis

All data collected were subjected to one-way analysis of variance (ANOVA) as described by Steel and Torrie (1980). Significant differences between mean were determined by Duncan's multiple range tests ($p < 0.05$) using statistical package for social sciences (SPSS) version 20. Values were expressed as means (\pm S.E)

Results

Table 3: Proximate Composition (% dry matter) of the Experimental Diets.

Parameters	Control ^{***} (0%)	Diet 2 (25%)	Diet 3 (50%)	Diet 4 (75%)	Diet 5 (100%)
C.P (%)	40.00	38.29	38.84	38.25	38.42
Fiber (%)	5.33	5.20	5.14	5.53	5.19
Lipid (%)	12.20	12.09	11.34	12.10	11.28
Ash (%)	7.50	8.17	7.85	7.40	7.62
Moisture (%)	12.17	13.19	11.00	11.54	11.69
NFE (%)	27.32	27.08	27.64	27.12	29.35
D. M (%)	93.16	91.33	91.09	91.13	91.38

C.V (kcal/g)	10.44	10.22	10.34	10.17	10.27
--------------	-------	-------	-------	-------	-------

CP = crude protein, NFE= nitrogen free extract, C.V=caloric value, DM = dry matter.

**Manufacturer's nutrient specification

The proximate composition of the experimental diets (table 3) shows that crude protein was highest in diet 1(40.00%) and lowest in diet 4(38.25%). Crude fiber was highest in diet 4(5.33%) and lowest in diet 3(5.14%). Ash content was highest in diet 2(8.17%) and lowest in diet 4(7.40%). Lipid was highest in diet 1(12.20%) and lowest in diet 5 (11.28%). Dry matter was highest in diet 1(93.16%) and lowest in diet 3(91.09%). NFE was highest in diet 5(29.35%) and lowest in diet 2(27.08%). Caloric value was highest in diet 1(10.44kc/g) and low in diet 4(10.17kc/g).

TABLE 4: Growth response, feed utilization and survival of *C.gariepinus* Juveniles Fed graded level of blood meal.

parameter	T1 (0%)	T2 (25%)	T3 (50%)	T4 (75%)	T5 (100%)
Fish stocked	60	60	60	60	60
Initial mean weight (g)	11.45±0.25	11.43±2.76	11.30±3.17	11.3±3.17	11.20±0.30
Final mean weight (g)	1275.00±0.00 ^a	1080.10±10.03 ^b	980.00±0.07 ^c	786.05±26.05 ^d	675.35±80.85 ^e
Mean weight gain	1263.55±140.59 a	1068.67±142.58 b	968.70±141.50 c	774.68±143.48 d	664.15±142.56 e

Initial mean length	9.26±2.84	9.15±3.04	9.25±2.07	9.21±0.91	9.23±1.67
Final mean length	63.42±2.66 ^a	60.00±0.00 ^b	59.21±8.97 ^c	59.00±0.00 ^d	56.09±5.30 ^e
Mean length gain	54.16±1.56	50.85±1.48	49.96±1.36	49.79±1.28	46.86±1.27
SGR	0.76±0.02	0.71±0.05	0.70±0.10	0.64±0.12	0.64±0.17
ADG	4.68±0.52	3.59±0.57	3.11±0.63	2.18±0.75	1.74±0.81
FCR	2.25±0.11	2.17±0.23	1.59±0.17	1.98±0.25	2.09±0.20
PER	31.59±3.51 ^a	24.22±3.67 ^b	20.98±4.07 ^c	14.73±4.11 ^d	11.73±4.17 ^e
PI	168.94±12.01 ^a	159.28±12.09 ^b	147.70±12.12 ^c	136.67±12.06 ^d	99.53±1.19 ^e
TFI	7650±347.92 ^a	6300±348.90 ^b	4350±346.89 ^c	3900±345.91 ^d	3450±346.75 ^e
SR (%)	100±1.16	100±1.18	98.00±1.20	96.00±1.23	100±1.19

Mean values within the rows with different superscripts are significantly different at ($p \leq 0.05$).

SGR= specific growth rate, ADG= average daily growth, PER = protein efficiency ratio, SR = survival rate, TFI = total feed intake, FCR = feed conversion ratio and PI = protein intake.

Results from the growth performance and feed utilization of *C. gariepinus* (table 4) showed that the initial mean length and initial body weight of fish among the treatment were not significantly different ($p \geq 0.05$), but were significantly different ($p < 0.05$) at the final body weight with the control diet (1275.00g) recorded the highest, followed by diet 2 (1080.10g), diet 3 (980.00g) and diet 4 (786.05g), while diet 5 (675.35g) recorded the least value. Similarly, the control diet had the highest weight gain (1263.55g), followed by diet 2 (1068.67g), diet 3 (968.70g), diet 4 (774.68g), while diet 5 had the lowest weight gain of (664.15g). Highest length gain was recorded in T1 (54.16cm ± 1.56), while lowest length gain was recorded in T5 (46.86cm ± 1.27). Specific growth rate in control diet and diets 2, 3, 4, 5 showed no significant difference ($p \geq 0.05$) from each other with diet 1 (0.76% ± 0.02) having the highest value, while diet 4 (0.64% ± 0.12) having lowest value. The average daily growth for T1 (4.68), T2 (3.59) and T3 (3.11) were not significantly different ($p \geq 0.05$) from each other and also, T4 (2.18) and T5 (1.74) were not significantly different. T1 (4.68) had the highest value and T5 (1.74). The feed conversion ratio (FCR) showed no significant

difference ($p \geq 0.05$) between all the treatments, control diet (2.25%) had the highest value and T3 (1.59%) being the lowest. The protein efficiency ratio (PER) and protein intake (PI) of the treatments showed significant difference ($p \leq 0.05$) from each other, T1 had the highest value of (31.59 and 168.94) while T5 had the least value of (11.79 and 99.53) respectively. Total feed intake (FI) were significant difference ($p \leq 0.05$), the highest feed intake was recorded in T1 (7650g) and lowest feed intake was recorded in T5 (3450g). The survival rate of fish showed no significantly different ($p \geq 0.05$), among the treatments.

Table 5: Water Quality Parameters taken during the feeding experiment of *Clarias gariepinus*

parameters	T1 (0%)	T2 (25%)	T3 (50%)	T4 (75%)	T5 (100%)
DO (mg/l)	5.82±1.08	5.95±1.11	5.81±1.09	6.26±0.81	6.05±0.89
pH	6.34±0.73	6.53±0.77	6.30±0.87	6.26±0.81	5.92±0.79
T(°C)	27.05±0.89	27.13±0.87	26.85±0.79	26.79±0.89	26.80±0.90
NH ₃ N (mg/l)	0.24±0.17 ^a	0.13±0.11 ^b	0.21±0.08	0.09±0.08 ^c	0.09±0.06 ^d

Mean values on the same row with different superscript are significantly different ($p < 0.05$).

DO=dissolved oxygen, NH₃N=ammonia-nitrogen, T= temperature.

Table 5 gives the results of water quality parameters obtained during the experimental period, no significant difference ($P>0.05$) was observed in dissolved oxygen, temperature and pH, but was no significantly different ($p<0.05$) in ammonia-nitrogen. Dissolved oxygen ranged from $5.81\pm 1.09\text{mg/l}$ – $6.26\pm 0.81\text{mg/l}$, pH ranged from 5.92 ± 0.79 – 6.53 ± 0.77 , Temperature ranged from $26.79\pm 0.89^{\circ}\text{C}$ – $27.13\pm 0.87^{\circ}\text{C}$ and ammonia-nitrogen ranged $0.09\pm 0.06\text{mg/l}$ – $0.24\pm 0.17\text{mg/l}$.

Table 6: Proximate carcass composition of experimental fish.

Parameters	Initial	Control (0%)	T2 (25%)	T3 (50%)	T4 (75%)	T5 (100%)
C. P.	54.30±0.18	69.50±0.06	68.00±0.11	67.50±0.40	68.30±0.19	69.10±0.32
Lipid	9.22±0.07	10.46±0.40	10.25±0.13	10.35±0.30	10.26±0.06 ^a	10.43±0.20
Ash	7.27±0.18	8.11±0.03	8.15±0.13	8.07±0.07	8.23±0.06	8.29±0.25
fibre	10.40±0.35	9.16±0.10	9.30±0.12	8.51±0.36	9.45±0.27	9.24±0.04
NFE	10.26±0.14	13.88±0.56	13.59±0.45	13.40±0.24	13.43±0.23	13.33±0.19
M.C	49.22±0.03	54.19±0.16	53.41±0.33	50.20±0.17	50.20±0.16	53.19±0.08

Dry matter	80.18±0.11	91.52±0.36	91.64±0.55	91.12±0.08	91.13±0.13	89.32±0.24
------------	------------	------------	------------	------------	------------	------------

Mean values without superscript are significant different ($p < 0.05$). NFE= nitrogen free extract, M.C = moisture content, C.P. = crude protein.

Proximate composition of the experimental fish before and after the study is presented in table 6. There was an increase in fish carcass in respect to dry matter, crude protein, lipid, ash, N.F.E, moisture content and lower fibre content at the end of experiment. Control diet (69.50%) had the highest value, while diet T3 (67.50%) had the least value in the carcass protein content. The carcass lipid content was high in control diet (10.46%) and low in while diet T2 (10.25%). Ash content was high in fish fed with diet T5 (8.29%) and low in diet T3 (8.07%). The moisture content was highest in fish fed with control diet (54.19%) and lowest in fish fed with diet T4 (50.20%). The N.F.E carcass content was highest in control diet (13.88%) and lowest in diet T5 (13.33%). The carcass dry matter of fish fed diet T2 (91.64%) was highest and diet T5 (89.32%). The whole body carcass showed a significant difference ($p < 0.05$) in fishes between the diet treatments and the initial fish sample, except in diet T4 of lipid carcass which showed no significant difference ($p < 0.05$).

Comment [ss7]: correct the spelling

Discussion

The ideal knowledge of the proximate composition of fish feeds (protein, fat, ash, NFE, fibre and moisture) contents are very important because it gives a reflection on the dietary and health importance of these nutrients in fish and also to final consumers which is man (Ogunjiet *al*, 2008; FAO, 2007), fish feeds components play a vital role in growth response, body development, metabolisms, repairing worn-out tissues and survival of fish. Proximate composition of the experimental diets indicated that the crude protein values obtained (38.48% - 40%) in this study were within the protein requirement (30%-40%) recommended

Comment [ss8]: correct the spelling

for growth performance in *Clarias gariepinus* reported by Fagberoet *al*, (2003) and Otubusinet *al*, (2009).

However, the general increase in the body weight of *Clarias gariepinus* in all the treatments indicated that the diets were accepted and utilized, irrespective of their composition, thereby promoting growth, survival and tissue development (Agbabiakaet *al*, 2013). This study revealed that the overall best growth performance and feed utilization was recorded on fish fed the control diet (0%) and was consistently followed by those fed diet 2 (25%), diet 3 (50%) and diet 4 (75%) blood meal inclusion, while the poorest growth performance was obtained with fish fed diet containing (100%) blood meal. Fish fed diets containing blood meal (25%) had significant better growth and feed utilization as compared to fish fed with the control diet than those of blood meal dietary treatments. This finding indicates that dietary supplementation of fishmeal with blood meal protein inclusion in the diet of *Clarias gariepinus* can only be efficiently replaced with 25% blood meal which enhanced fish growth and survival without adverse effects, and this observation agrees with the result of Agbebi *al*, (2009) who reported that a 25% of blood meal substitution of fish meal in diet gave best growth performance. Similarly, Adejoke (2012) on the use of bovine blood and rumen digest in catfish diet to replace fish meal at 0%, 25%, 50%, 75% and 100% where he reported that the best performance was recorded in fish fed with the control diet and treatment diet with inclusion level of 25% of replacement diet. This observation is in contrast with the findings of Olukunle *et al.*, (2002); Otubusin, (2001) and Aliu and Dako (2018).

As the level of blood meal inclusion increased in the diets, the performance of fish reduced. Replacement of 75% – 100% fishmeal resulted in reduced growth and feed utilization when compared with other experimental diets; it probably indicated the absence of some nutrients particularly some of the essential amino acid profile presents in the control diet and 25% diet which promotes growth on the fish juveniles (Ogunji, 2004). Blood meal has been reported to have imbalance amino profile unlike fishmeal; it is an excellent source of crude protein

and high lysine (Sauvant, 2004) but low in methionine and isoleucine. According to Adewolu and Adamson (2011), deficiency in methionine has been reported to lead to reduced fish growth. Poor growth performance indicates that blood meal cannot be used as a sole protein source in diet formulation and might be due to the nature of its imbalance amino profile, non-attractive odour and digestibility which results in low feed consumption and low performance of fish (Kirimiet al 2016). This study agreed with the study done by Otubusin (1987) and Gatlin (2010) but disagrees with Agbebi et al (2009) and Aladetohun and Sogbesan (2013).

In this study, survival rate of fish in all the treatments were high and showed no significant difference ($p \leq 0.05$). High survival rate could be attributed to the good management precaution during the study and the experimental diets did not have any adverse effect on the health and well-being of the fish which agrees with the report of Agbebi et al (2009) and Aladetohun and Sogbesan (2013).

The physiochemical parameters of water in this study were not affected by the diets and were within the range recommended for *Clarias gariepinus* culture (Boyd, 1979; Bhatnagar et al, 2004; Solomon et al, 2013 and Swann 2007).

Analysis of the carcass fish indicate more protein retained in the body at the end of experiment which revealed that incorporation of blood meal in the diets affected significantly ($p < 0.05$) to the body of fish in a positive way, they converted and utilized the protein synthesis from the feed into their body which increases tissue production and real growth. The results showed an increase in the value of crude protein and lipid over the initial fish samples. This agrees with the reports of Ochan et al (2007), Fountoulaki et al (2003), Fafioye et al (2005)

Conclusion

The results of this present study showed that fish fed with fishmeal based diet best performance in terms of growth rate; ironically it is also the most expensive protein source

due its nutrient profile and non-availability. However, Blood meal diet from this study is proved suitable for use in aquaculture sector, those fed on 25% blood meal substitution performed better than 100% blood meal, hence blood meal can substitute fishmeal partially at 25% inclusion and it has the potential to make considerable contribution to growth and survival rate of *Clarias gariepinus* with no adverse effects. The utilization of blood meal as a protein supplement in fish diets will boost the profitability and sustainability of the fish production, reduced the cost of feed and alleviate the problem of disposal of blood in abattoirs and environmental pollution

References

- Adejoke, A.A. (2012). Bovine blood and ruminant digests in catfish *Clarias gariepinus* diet. *European Journal of Scientific Research*. 2012; 83:167-172.
- Adewolu, M.A., Akintola, S.L. and Akinwunmi, O.O. (2009). Growth performance and survival of hybrid African Catfish larvae (*Clarias gariepinus* × *Heterobranchius bidorsalis*) fed different diets. *Zoologist* 7(1): 45-51.
- Adikwu, I.A. (1991). The development of complete diet from local feed stuffs and industrial wastes for the culture of tilapia (*Oreochromis niloticus*) in Nigeria. Technical Report for the International Foundation for Science. 1991; 2:005-114.
- Adikwu, I.A., (2003). A review of Aquaculture Nutrition in Aquaculture Development in Nigeria. In: Eyo, A.A (Ed) *National Workshop on Fish Feed Development and Feeding Practices In Aquaculture*, Organized by Fisheries Society of Nigeria (FISON) In collaboration with National Institute

of Freshwater Fisheries Research (NIFFR) and FOA National Special Programmed For Food Security (FAO-NSPFS) pp 34-40.

- Agbebi, O.T., S.O. Otubusin and F.O. Ogunleye, (2009). Effect of different levels of substitution of fishmeal with blood meal in pelleted feeds on catfish *Clarias gariepinus* (Burchell, 1822) culture in net cages. *European Journal Scientific Research*. 31: 6-10.
- Aladetohun, N.F. and O.A. Sogbesan, (2013). Utilization of blood meal as a protein ingredient from animal waste product in the diet of *Oreochromis niloticus*. *International Journal of Fisheries and Aquaculture*, 5: 234-237.
- Aliu, B.S. and Dako, M.O. (2018). Effect of fishmeal replacement with blood meal on the growth response and utilization of hybrid (*Clarias gariepinus* and *Heterobranchus bidosaris*) fingerlings. *Asian Journal of Research in Agriculture and Forestry*. 2018; 1(3):1-6.
- Allan, G. (2004). Fish for feed vs. fish for food. In: A. G. Brown (Ed) Fish, Aquaculture and Food Security: Sustaining Fish as a Food Supply. Record of a conference conducted by the ATSE Crawford Fund, Parliament House Canberra, 11 August 2004, pp. 20-26.
- AOAC (2005). Official Methods of Analysis (18th edition) Association of Official Analytical, Chemists. 18th Edition. *AOCA international Official Program* Gaithersburg, Madison, USA.
- Burel, C., Boujard, T., Tulli, F. and Kaushik, S.J. (2000). Digestibility of extruded peas extruded lupin and replaced meal in rainbow trout (*Oncorhynchus mykiss*) and turbot (*Psetta maxima*) diet. *Aquaculture journal*, 188: 285-298.
- Dawah, M.A., A.M. Khater, I.M.A. Shaker and N.A. Ibrahim, (2002). Production of *Scenedesmus Bijuga* (Chlorophyceae) in large scale in outdoor tanks and its use in feeding monosex Nile tilapia (*Oreochromis niloticus*) fry. *Journal of Egypt Academic Society*.

- De-Silva, S.S. and Anderson, T.A. (1995). Fish nutrition in Aquaculture. Chapman and Hall, London, U.K, 1195. ISBN 13:97804 1255030, page 319.
- Eyo, A.A. and Olatunde, A.A., (1999). The effect of replacement of soya bean meal with blood meal on the growth of mudfish (*Clariasanguillaris*) fingerlings. Paper presented at the 13th Annual Conference of the Fisheries Society of Nigeria (FISON). New Bussa, Nigeria, November 1999.
- Fagbenro, O.A., Balogun, A.M and Fasakin, E.A. (1999b). Dietary methionine requirement of the African catfish, *Clariasgariepinus*. *Journal of Applied Aquaculture* 8(4), 47-54.
- Food and Agricultural Organization (FAO), (2016). The State of World Fisheries and Aquaculture 2016. Contributing of food security and nutrition for all. Rome, FAO, pp: 200.
- Food and Agriculture Organization (FAO), (2007). Assessment of freshwater fish seed resources for sustainable aquaculture. Rome, Italy: Food and Agricultural Organization.
- Food and Agriculture Organization (FAO), (2012). Food and Agricultural Organization of the United Nations on the state of the world fisheries and aquaculture part 1: World Review of Fisheries and Aquaculture. Rome. 230p.
- Food and Agriculture Organization (FAO), (2000). The State of World Fisheries & Aquaculture 2000. Rome, Italy: Food and Agricultural Organization.
- Gabriel, U.U., Akinrotimi, O.A., Bekibele, D.O., Onunkwo, D.N and Anyanwu, P.E. (2007). Locally produced fish feed: potentials for aquaculture development in sub-sharan Africa. *African Journal of Agricultural Research* 2(7): 287-295.

- Gatlin D.M., (2010). Principles of Fish Nutrition. Southern Regional Aquaculture Center SRAC publication No. 5003.july 2010, pdf, pp.1-8.
- Guroy, D., Emrah, D., Kut, G., and Adem, A. (2005). Influence of feeding frequency and feed intake, growth performance and nutrient utilization in European sea bass (*Dicentrarchus labrax*) fed pelleted or extruded diets. *Turkish Journal of Veterinary Animal Science*, 30, 171-177.
- Hardy, R. W., and Tacon, A. G. J. (2002). Fish meal: Historical uses, production trends and future outlook for sustainable supplies. In R. R. Stickney & P. MacVey (Ed.), *Responsible Marines Aquaculture* (pp. 311-325). CABI, UK.
- Hertrampf, J. W. and Piedad-Pascual, F., (2003). Blood products in handbook on Ingredients for Aquaculture Feeds. Dordrecht, Netherlands: Kluwer Academic Publishers, pp. 69-79.
- Heuzé, V., Tran, G. and Kaushik, S., (2015). Fish meal. Feedipedia, FAO. <http://www.feedipedia.org/node/208>.
- Kirimi, J.G, Musalia, L.M. and Munguti, J.M. (2016). Effects of replacing fish meal with blood meal of chemical composition of supplement for Nile tilapia (*Oreochromis niloticus*). *East African Agricultural and Forestry Journal*. 2016; 1:1-9.19.
- Madu, C.T., Sogbesan, O.A. and Ibiyo, L.M.O. (2003). Some non-conventional fish feed resources in Nigeria. In: Eyo A.A.(Ed.), *Proceeding of the Joint Fisheries Society of Nigeria/National Institute For Freshwater Fisheries Research/FAO National SpecialProgramme For Food Security National workshop on Fish feed development and Feeding Practices in Aquaculture* held at National Institute for Freshwater Fisheries Research, 15th -19th Sept. 2003. New- Bussa: 73-82.
- Njoku, D.C., (2010). Laboratory Manual for Fisheries and Aquaculture Practical (Vol 1). *Martins Continental Publishers*. Owerri, Nigeria. 108pp.

- Nogueira, N., Cordeiro, N., Andrade, C. and Aires, T., (2012). Inclusion of low levels of blood and feather meal in practical diets for Gilthead Sea bream (*Sparus aurata*). *Turkish Journal of Fisheries and Aquatic Sciences*, 12, pp. 641-650.
- Ogunji, J. R., Summan Toor, C. and Schulz, W.K. (2008). Growth performance, Nutrient Utilization of Nile tilapia (*Oreochromis niloticus*) fed housefly maggot meal (Magmeal) diets. *Turkish Journal of Fisheries and Aquatic Sciences*, 8(1): 141-147.
- Ogunji, J.O and Wirth, M. (2001). Alternative protein sources as substitutes for fish meal in the diet of young tilapia *Oreochromis niloticus* (linn). Israeli Journal of Aquaculture-Bamidgah, 53(1), 34-43.
- Ogunji, J.O., (2004). Alternative protein source in diets for farmed *Tilapia*. Animal Science Comparative Reviews Number 13; CAB International Publishing (Oxford, UK). *Nutrition Abstracts and Reviews: Series B*, 74(8): 23-32
- Olukunle, A.O, Ogunsanmi, A.O, Taiwo, J.O, and Samuel, A.A. (2002). The nutritional value of cow blood on the growth performance haemathology and plasma enzymes of hybrid catfish. *Nigerian Journal of Animal Science*.2006; 5:75-85.
- Otubusin S.O., Ogunleye F.O. and Agbebi O. T. (2009). Feeding trials using local protein sources to replace fishmeal in pelleted feeds in catfish (*Clarias gariepinus* Burchell 1822) Culture. *European Journal of Scientific Research* 31(1):142-147.
- Steel, R.G.D. and Torrie, J.H. (1980). *Principles and Procedures of Statistics. A biometrical approach*. 2nd edition. McGraw-Hill, New York, USA, pp. 20-90.

- Teugels, G. (1986). A systematic revision of the Africa species of the genus (Pisces: *Claridae*). *Annals muse Royal de l' Afrique Centrale*, 247:1-199.
- Wilson, R. P. (2002). Protein and amino acids. In J. E. Halver & R. W. Hardy (Eds.), *Fish Nutrition* (3rd version, pp. 144-179). Elsevier Science, San Diego, USA.
- Otubusin, S.O (1987). Effect of different levels of blood meal in pelleted feeds on Tilapia, *Oreochromis niloticus* production in floating bamboo net cages. *Aquaculture*, 65, 263-266.
- NRC (2011). *Nutrients requirements of fish and shrimps*. National Academy Press. Washington D.C., USA.
- Fagbenro, O.A, Adeparusi, E.O and Fapohunda, O.O (2003). Feedstuff and dietary substitution for farmed fish in Nigeria. In: Eyo A.A.(Ed.), *Proceeding of the Joint Fisheries Society of Nigeria/National Institute For Freshwater Fisheries Research/FAO National Special Programme For Food Security National workshop on Fish feed development and Feeding Practices in Aquaculture* held at National Institute for Freshwater Fisheries Research, New- Bussa 15th -19th Sept. 2008. Pp 60-72.
- APHA. (2005). American Public Health Association, Standard Methods for the Examination of Water and Waste Water. American Public Health Association Washington DC, USA. 1368p.
- Otubusin, S.O (2001). Economics of small scale table Tilapia production in net-cages. *ASSET Series*. 2001; A1:83-90.
- Bhatnagar, A., Jana, S.N., Garg, S.K., Patra, B.C., Singh, G., and Barman, U.K., (2004). Water quality management in aquaculture, in: *Course Manual of Summer School*

on development of sustainable aquaculture technology in fresh and saline waters, CCS Haryana Agricultural, Hisar (India), pp 203-210.

Adewolu, M.A and Adamson, A.A. (2011). *Amaranthus spinosus* leaf meal in the diet of Nile tilapia. *Pakistan Journal of Science and Industrial Research*. 55:335-338.

Sauvant, D. (2004). Tables of composition and nutritional value of feed materials: pigs, poultry, cattle, sheep, Goats, rabbits, horses and fish (2nd ed., p304). Wageningen, Netherlands Wageningen Academic Publishers.

Boyd, C.E., (1979). Water quality in Warm water fish ponds, Agriculture Experiment Station, Auburn, Alabama, pp 359.

Fafioye, O.O., Fagade, S.O., Adebisi, A.A, Jenyo-Oni and Omoyinmi, G.A.K. (2005). Effects of dietary soya beans (*Glycine max* L.) on growth and body composition of *Clariasgariepinus* fingerlings. *Turkish Journal of Fisheries and Aquatic Science*, 5:11-15.

Fountoulaki, E., Alexis, M.N., Nengas, L. and Venon, B. (2003). Effects of dietary arachidonic acid (20:4n-6) on growth, body composition and tissue fatty acid profile of gilthead bream fingerlings (*Sparus aurata*). *Aquaculture Journal*, 255:309-323,

Ochang, S.N., Fagbenro, O.A. and Adebayo, O.T. (2007). Growth performance, body composition, haematology and product quality of African catfish (*Clariasgariepinus*) fed diets with palm oil. *Pakistan Journal of Nutrition*. 6(5): 452-459.

Solomon, W.G.O., Olatunde, A.A. and Matur, B.M. (2013). Some physic-chemical parameters of selected fish ponds in Gwagwalada and Kuje Area Councils, Federal Capital, Territory, Nigeria. *Global Advanced Research Journal of Aquaculture Science*, 2(1) pp.017-022.

Swann, L.D., (2007). A fish farmer's Guide to Understanding Water Quality, Aquaculture Extension Illinois, and Purdue University, Indiana Sea Grant Program Fact Sheet AS-503.

Agbabiaka, L.A, Ezeafulukwe, C.F and Ekeledo, C.B (2013). Recycle dead chicken meal in low cost practical diets for African catfish (*Clariasgariepinus*) fingerling. Agriculture and biological journal of North America, 2151-751.

UNDER PEER REVIEW