

Replacement of Fish Meal with Fermented Soybean Meal in Fish Feed: A Review

ABSTRACT

Soybean meal is a source of vegetable protein substitute for fish meal that is widely used because it has a high protein content, relatively cheap price and easy to obtain. Besides being high in protein, soybean meal also contains anti-nutrients that can inhibit growth, feed effectiveness, and lower nutrient utilization in fish. So it is necessary to process feed ingredients to remove anti-nutrient content. Fermentation can be a solution for the utilization of soybean meal by reducing the anti-nutritional content and crude fibers so as to produce components that can be utilized properly, such as protein, β -glucan, vitamins, mannanoligosaccharides, organic acids, and antibiotics. The type of soybean meal fermentation that is widely used is with *Bacillus* spp., *Aspergillus* spp., *Lactobacillus* spp., and yeast. The purpose of this literature review study was to determine the effect of replacing fish meal with fermented soybean meal on fish feed. Several types of fish that have been fed with fermented soybean meal include seabass, green snapper, mud grouper, rainbow trout, star pomfret, tilapia, and turbot. Giving fermented soybean meal to fish can increase growth rate, intestinal morphology, intestinal microecology, feed digestibility, feed preference, fish innate immunity and produce antimicrobial peptides. Soybean meal fermentation as a substitute for fish meal has been shown to have a good effect on fish so that it can continue to be developed to increase the utilization of soybean meal.

Keywords : Fish feed, fermentation, soybean meal, growth, digestibility

1. INTRODUCTION

Fish meal has a high protein content so that it can be used as the main protein source in the manufacture of carnivorous fish feed with additional contents such as a balanced amino acid content and the content of other growth triggering factors [1]. However, the use of fish meal has a negative impact on the development of a sustainable aquaculture industry and is an obstacle to increasing demand for feed production [2]. The diminishing fishery resources and increasing fish prices are the main factors urging nutritionists to replace fish meal without reducing the protein content and easy to find [3].

Candidate sources of protein as a substitute for fish meal are soy flour extracted with solvents, because of its wide availability and stable supply [2]. Soybean meal is a by-product of processed soybean oil with a high protein content (about 40-50%) and balanced amino acids, so it can be used as the main protein source in the manufacture of animal feed and the food industry [4]. Besides having a high protein content, soybean meal also has a high carbohydrate

Comment [s1]: What percent of protein content? And how much does it cost compared to other feed ingredients? explained in the abstract.

Comment [s2]: What is the percent protein content?

Comment [s3]: what is the negative impact of the use of fish meal? explain in detail.

Comment [s4]: What is the percent protein of fish meal and soybean meal?

content[5]. However, the use of soybean meal as an alternative raw material is still limited because soybean meal contains anti-nutrients such as oligosaccharides, saponins, protease inhibitors and phytic acid [6] which can inhibit growth, physiological conditions and feed utilization so that the nutrients in the feed are not absorbed completely [7].

Comment [s5]: What is the percentage of carbohydrates in soybean meal?

Comment [s6]: Why choose soybean meal? Is the price cheaper than fishmeal and other ingredients? What are the advantages and disadvantages of soybean meal?

Substitution of fishmeal using soybean meal flour often causes stunted growth and lower nutrient utilization in feed [8][9]. Previous studies suggested the reduction or even elimination of anti-nutritional content in soybean meal using various techniques to increase the nutritional value content [10]. So it is necessary to process feed raw materials before being given to fish [11].

Fermentation is the process of chemical changes in certain materials with the help of enzyme activity derived from microorganisms [12]. Make other beneficial components such as probiotics and prebiotics [13]. Making this journal review aims to determine the effect of replacing fishmeal with fermented soybean meal on fish feed.

Comment [s7]: Explain the advantages of fermentation.

Comment [s8]:

2. SOYBEAN MEAL AS A SUBSTITUTE FOR FISH MEAL

Soybean meal is often used as a substitute for fishmeal because it has a relatively high protein content, easy to find and low price [14]. Several studies also said that soybean meal has been widely used as an alternative protein source to replace fishmeal in fish culture feed, because of its high protein content, relatively low cost and easy to find [15]. Besides having a high protein content, soybean meal also has a high carbohydrate content [5]. However, the use of soybean meal in the diet of carnivorous fish such as orange spotted grouper (30%) [16], European seabass (25%), and turbot (30%) [17] needs a benchmark in feeding soybean meal because it can hamper performance, growth of the carnivorous fish.

Comment [s9]: What percent of protein is in fishmeal and soybean meal? How much does a soybean meal cost compared to a fish meal? Is it expensive or cheap compared to other materials?

Comment [s10]: Adding new references, don't quote sentences repeatedly like in the introduction

Comment [s11]: What is the carbohydrate content and other nutritional content? Explained in detail all the nutritional content.

The high use of soybean meal in feed can cause adverse effects on growth, feed utilization and fish health status [15]. Several studies have shown that feeding fermented soybean meal in aquaculture ponds can affect immune function, adverse inflammation responses and stress in fish. [3]. The anti-nutrient content in soybean meal raw materials can inhibit the efficiency of feeding so that the nutritional content is disturbed such as the content of phytic acid [18]. In general, plant-type feed ingredients contain anti-nutrients which are the main source of phosphate and cause low feed solubility in monogastric animals. The strategy that can be used to remove anti-nutritional content in soybean meal is to utilize the fermentation process by microorganisms which has many benefits such as increasing the nutritional content and increasing the solubility of vegetable protein sources by fish [14].

Comment [s12]: What is the maximum percentage of soybean meal used for fish feed? Explain clearly

Comment [s13]: What is the percent content of fermented soybean meal? Can it remove anti-nutrients?

3. FERMENTATION FUNCTION

Fermentation is one method that is widely used in increasing the utilization of fishmeal substitutes with soybean meal [19]. Previous studies have determined that fermentation is an effective method to improve the quality of soybean meal by producing probiotics or prebiotics. So it can improve nutrient digestibility, fish immune system and palatability by eliminating anti-nutrients, oligosaccharides, protein and fiber. [20]. In the soybean meal fermentation process, the most widely used microorganisms are *Bacillus* spp., *Aspergillus* spp., and *Lactobacillus* spp. [21][22]. Several studies also explained that the use of fermented soybean meal with *Bacillus* spp. can increase protein content, tioxidative activity, total amino acids, fish digestibility and can reduce anti-nutrient content and immunoreactivity [9]. So that fermentation is highly recommended to increase the utilization of soybean meal [14].

Comment [s14]: What are the advantages of fermentation? Does it have any effect on feeding? Explained

Comment [s15]: Too few references and discussions. Add the new reference

4. NUTRIENTS IN FERMENTATION OF SOYBEAN MEAL IN FISH FEED

Feed ingredients that are processed by the fermentation process will experience an increase in nutrients such as increased protein and feed digestibility through a decrease in crude fiber and fat levels [23]. Several studies say that fermentation can be used to improve the quality of soybean meal. The nutritional content of fermented soybean meal can improve fish growth performance, digestive physiological conditions, and the coefficient of digesting nutrients. The type of fermentation that is widely used is with the bacterium *Bacillus subtilis* and *Lactobacillus* spp. [7]. List of nutritional content and type of fermentation in several studies can be seen in (Table 2).

Comment [s16]: Too few references and discussions. Add the new reference

Table 1: Nutrient content of fermented soybean meal

Fermentation Type	Crude Protein (%)	Carbohydrates (%)	Ash Content	Lipids Rough (%)	Reference
<i>Bacillus pumillus</i> SE5 42-	-	10 [2]			
<i>Aspergillus layor</i> 60.58	-	2.7 [15]			
<i>Bacillus subtilis</i> E20	48.82	11.85 7.50 [16]			
<i>Bacillus subtilis</i> , <i>Lactobacillus</i> , yeast	65.5	-	-	-	[6]
<i>Lactobacillus</i> spp.	50	-	-	-	[7]
<i>Saccharomyces Cerevisiae</i>	29.7	57.15	5.91	-	[24]

5. FERMENTATION OF SOYBEAN MEAL AS FISH FEED

Soybean meal fermented using microorganisms has been widely used in several types of fish and produces soybean meal with a higher preference as vegetable protein [25]. Fermented soybean meal has been widely used as feed for seabasses [2], greensnapper [26], mud grouper [16], rainbow trout [6], star pomfret [7], tilapia [24], and turbot [15]. Another study mentioned that the use of soybean meal fermentation using *Bacillus* spp. can produce antibiotics [27] and does not cause any morphological changes in the intestines of fish, especially in rainbow trout [25]. Soybean meal fermentation using lactic acid bacteria can increase the acidity of the feed, increase feed preference, and can produce antimicrobial peptides [28]. Soybean meal fermented using yeast can produce useful components such as mannanoligosaccharides, organic acids, vitamins, antibiotics, β -glucans, and chitin [29]. Components produced from soybean meal fermentation using microbes can increase the growth rate, intestinal morphology, intestinal microecology, and innate immunity of fish [30], [31]. The use of fermented soybean meal in fish has different optimum levels (Table 1).

Comment [s17]: Too few references and discussions. Add the new reference

Table 2: Growth and optimum dose of fermented soybean meal

Species	Results (WG%)	Optimum Dose	Reference
<i>Lateolabrax maculatus</i>	12.3	26.9-37.1%	[2]
<i>Micropterus salmoides</i>	14.71	30%	[26]
<i>Epinephelus coioides</i>	34.81	29.32%	[16]
<i>Oncorhynchus mykiss</i>	5.5	40%	[6]
<i>Trachinotus blochii</i>	25.3	35%	[7]
<i>Oreochromis niloticus</i>	21.87	37.4%	[24]
<i>Scophthalmus maximus</i> L.	N/A	45%	[15]

6. CONCLUSION

The use of fermented soybean meal in replacing fish meal is an alternative that can be used because the protein content in fermented soybean meal is not much different from that of fish meal. In addition, fermented soybean meal is also easy to find and the price is relatively cheap. Although the use of fermented soybean meal in fish feed is very promising, there are still not many efficient uses for fish growth. Therefore, it is necessary to determine the species and type of fermentation that are suitable for the use of fermented soybean meal in fish feed.

REFERENCES

- [1] H. Yang, Y. Bian, L. Huang, Q. Lan, L. Ma, X. Li, X. Leng, Effects of Replacing fish meal with fermented soybean meal on the growth performance, intestinal microbiota, morphology and disease resistance of largemouth bass (*Micropterus salmoides*), *Aquac. Reports*. 22(2022)100954. <https://doi.org/10.1016/j.aqrep.2021.100954>.
- [2] S. Rahimnejad, J. Zhang, L. Wang, Y. Sun, C. Zhang, Evaluation of *Bacillus pumilus* SE5 fermented soybean meal as a fish meal replacer in spotted seabass (*Lateolabrax maculatus*) feed, *Aquaculture*. 531(2021) 735975. <https://doi.org/10.1016/j.aquaculture.2020.735975>.
- [3] S. Rahimnejad, S. Hu, K. Song, L. Wang, K. Lu, R. Wu, C. Zhang, Replacement of fish meal with defatted silkworm (*Bombyx mori* L.) pupa meal in diets for Pacific white shrimp (*Litopenaeus vannamei*), *Aquaculture*. 510(2019) 150–159. <https://doi.org/10.1016/j.aquaculture.2019.05.054>.
- [4] F. Lu, E. Alenyorege, N. Ouyang, A. Zhou, H. Ma, Simulated natural and high temperature solid-state fermentation of soybean meal: A comparative study regarding microorganisms, functional properties and structural characteristics, *Lwt*. 159(2022) 113125. <https://doi.org/10.1016/j.lwt.2022.113125>.
- [5] M. Amin, F. H. Taqwa, Y. Yulisman, R. C. Mukti, M. A. Rarassari, R. M. Antika, Effectiveness of Utilization of Local Raw Materials as Fish Feed on Increasing Productivity of Cultivation of Catfish (*Clarias* sp.) in Sakatiga Village, Indralaya District, Regency Ogan Ilir, South Sumatra, *J. Aquac. Fish Heal*. 9(2020) 222. <https://doi.org/10.20473/jafh.v9i3.17969>.
- [6] D. G. Choi, M. He, H. Fang, X. L. Wang, X. Q. Li, X. J. Leng, Replacement of fish meal with two fermented soybean meals in diets for rainbow trout (*Oncorhynchus mykiss*), *Aquac. Nutrient*. 26(2020) 37–46. <https://doi.org/10.1111/anu.12965>.
- [7] H. P. Nguyen, T. Van Do, H. D. Tran, Dietary replacement of fish meal by defatted and fermented soybean meals with taurine supplementation for pompano fish: Effect on growth performance, nutrient digestibility, and biological parameters in a long-term feeding period, *J. Anim. Science*. 98(2020) 1–9. <https://doi.org/10.1093/JAS/SKAA367>.
- [8] Z. Song, H. Li, J. Wang, P. Li, Y. Sun, L. Zhang, Effects of fish meal replacement with soy protein hydrolysate on growth performance, blood biochemistry, gastrointestinal digestion and muscle composition of juvenile starry flounder (*Platichthys stellatus*), *Aquaculture*. 426–427(2014) 96–104. <https://doi.org/10.1016/j.aquaculture.2014.01.002>.
- [9] D. Teng, M. Gao, Y. Yang, B. Liu, Z. Tian, J. Wang, Bio-modification of soybean meal with *Bacillus subtilis* or *Aspergillus oryzae*, *Biocatal. Agric. Biotechnol*. 1(2012) 32–38. <https://doi.org/10.1016/j.bcab.2011.08.005>.

Comment [s18]: The conclusions are not clear. What type of fermentation is good and suitable for what bacteria? not in accordance with the contents see table 1 and table 2

Comment [s19]: Add new reference

- [10] DM Gatlin, FT Barrows, P. Brown, K. Dabrowski, TG Gaylord, RW Hardy, E. Herman, G. Hu, K. Krogdahl, R. Nelson, K. Overturf, M. Rust, W. Sealey, D. Skenberg, EJ Souza, D. Stone, R. Wilson, E. W. Urtele, Expanding the utilization of sustainable plant products in aquafeeds: A review, *Aquac. res.* 38(2007)551–579. <https://doi.org/10.1111/j.1365-2109.2007.01704.x>.
- [11] W. Pamungkas, Fermentation Technology, Alternative Solutions in Efforts to Utilize Local Feed Materials, *Aquaculture Media*. 6(2011) 43. <https://doi.org/10.15578/ma.6.1.2011.43-48>.
- [12] Y. Suryani, I. Hernaman, N. Ningsih, Effect of Addition of Urea and Sulfur to Em-4 Fermented Bioethanol Solid Waste on Protein and Crude Fiber Content, *J. Ilm. Breeder. Integrated*. 5(2017) 13. <https://doi.org/10.23960/jipt.v5i1.p13-17>.
- [13] KJ Hong, CH Lee, WK Sung, *Aspergillus oryzae* GB-107 fermentation improves nutritional quality of food soybeans and feed soybean meals, *J. Med. Food*. 7(2004) 430–435. <https://doi.org/10.1089/jmf.2004.7.430>.
- [14] Z. Wang, M. Yang, L. Wang, K. Lu, K. Song, C. Zhang, *Bacillus subtilis* LCBS1 supplementation and replacement of fish meal with fermented soybean meal in bullfrog (*Lithobates catesbeianus*) diets: Effect on growth performance, feed digestibility and gut health, *Aquaculture*. 545(2021) 737217. <https://doi.org/10.1016/j.aquaculture.2021.737217>.
- [15] C. Li, B. Zhang, H. Zhou, X. Wang, X. Pi, X. Wang, K. Mai, G. He, Beneficial influences of dietary *Aspergillus* or fermented soybean meal on oxidative homeostasis and inflammatory response in turbot (*Scophthalmus maximus* L.), *Fish Shellfish Immunol.* 93 (2019) 8–16. <https://doi.org/10.1016/j.fsi.2019.07.037>.
- [16] YL Shiu, SL Hsieh, WC Guei, YTT sai, CH Chiu, CH Liu, Using *Bacillus subtilis* E20-fermented soybean meal as replacement for fish meal in the diet of orange-spotted grouper (*Epinephelus coioides*, Hamilton), *Aquac. res.* 46(2015) 1403–1416. <https://doi.org/10.1111/are.12294>.
- [17] L. Wang, Y. Jian, Q. Liu, F. Li, L. Chang, Electromagnetohydrodynamic flow and heat transfer of third grade fluids between two micro-parallel plates, *Colloids Surfaces A Physicochem. eng. Asp.* 494(2016) 87–94. <https://doi.org/10.1016/j.colsurfa.2016.01.006>.
- [18] L. Pan, J. Liu, MH Farouk, G. Qin, N. Bao, Y. Zhao, H. Sun, Anti-nutritional characteristics and mechanism of soybean agglutinin, *Biocell.* 45(2021) 451–459. <https://doi.org/10.32604/BIOCELL.2021.014289>.
- [19] B. Zhang, Y. Zhang, M. Cui, M. Zhang, J. Xu, Z. Zhang, Z. Sui, L. Wang, C. Zhang, C. Li, Q. Ma, Comparison of the performance of raw and *Lactobacillus paracasei* fermented soybean meal in diets for turbot (*Scophthalmus maximus* L.): Growth, intestinal morphology, apoptosis, tight junction, and microbiota, *Aquac. Reports*. 24(2022) 101184. <https://doi.org/10.1016/j.aqrep.2022.101184>.
- [20] XF Liang, L. Hu, YC Dong, XFWu, YC Qin, YH Zheng, DD Shi, M. Xue, XF Liang, Substitution of fish meal by fermented soybean meal affects the growth performance and flesh quality of Japanese seabass (*Lateolabrax japonicus*), *Anim. Science Feeds. Technol.* 229(2017) 1–12. <https://doi.org/10.1016/j.anifeeds.2017.03.006>.
- [21] CH Chi, SJ Cho, Improvement of bioactivity of soybean meal by solid-state fermentation with *Bacillus amyloliquefaciens* versus *Lactobacillus* spp. and *Saccharomyces cerevisiae*, *Lwt.* 68(2016) 619–625. <https://doi.org/10.1016/j.lwt.2015.12.002>.

- [22] LSSuryaningrum, Application of microbes in effort to improve the quality of fish feed raw materials through fermentation, *Pros. Biol. Achiev. sustain. Dev. Goals with Biodivers. Confronting Clim. Chang.* (2021) 204–210.
- [23] R. Rostika, R. Safitri, Influence of Fish Feed Containing Corn-Cob Was Fermented By *Trichoderma Sp.*, *Aspergillus Sp.*, *Rhizopus Oligosporus* To The Rate of Growth of Java Barb (*Puntius Gonionotus*), *APCBEE Procedia*. 2(2012) 148–152. <https://doi.org/10.1016/j.apcbee.2012.06.027>.
- [24] MSHassaan, MASoltan, MAAbdel-Moez, Nutritive value of soybean meal after solid state fermentation with *Saccharomyces cerevisiae* for Nile tilapia, *Oreochromis niloticus*, *Anim. Science Feeds. Technol.* 201(2015)89–98. <https://doi.org/10.1016/j.anifeedsci.2015.01.007>.
- [25] ME Barnes, ML Brown, KARosenrater, JR Sewell, An initial investigation Replacing fish meal with a commercial fermented soybean meal product in the diets of juvenile rainbow trout, *Open J. Anim. science*. 02(2012)234–243. <https://doi.org/10.4236/ojas.2012.24033>.
- [26] M. He, X. Li, L. Poolsawat, Z. Guo, W. Yao, C. Zhang, X. Leng, Effects of fish meal replaced by fermented soybean meal on growth performance, intestinal histology and microbiota of largemouth bass (*Micropterus salmoides*), *Aquac. Nutrient*. 26(2020) 1058–1071. <https://doi.org/10.1111/anu.13064>.
- [27] AC Cheng, HLLin, YLShiu, YC Tyan, CH Liu, Isolation and characterization of antimicrobial peptides derived from *Bacillus subtilis* E20-fermented soybean meal and its use for preventing *Vibrio* infection in shrimp aquaculture, *Fish Shellfish Immunol.* 67 (2017) 270–279. <https://doi.org/10.1016/j.fsi.2017.06.006>.
- [28] S. Sanjukta, AK Rai, Production of bioactive peptides during soybean fermentation and their potential health benefits, *Trends Food Sci. Technol.* 50(2016)1–10. <https://doi.org/10.1016/j.tifs.2016.01.010>.
- [29] M. Faustino, J. Durão, CFPereira, MEPintado, APCarvalho, Mannans and mannanoligosaccharides (MOS) from *Saccharomyces cerevisiae*—A sustainable source of functional ingredients, *Carbohydr. Polym.* 272(2021). <https://doi.org/10.1016/j.carbpol.2021.118467>.
- [30] Y. Silva-Carrillo, C. Hernández, RW Hardy, B. González-Rodríguez, S. Castillo-Vargasmachuca, The effect of substituting fish meal with soybean meal on growth, feed efficiency, body composition and blood chemistry in juvenile spotted rose snapper *Lutjanus guttatus* (Steindachner, 1869), *Aquaculture*. 364–365(2012)180–185. <https://doi.org/10.1016/j.aquaculture.2012.08.007>.
- [31] L. Li, J. Song, C. Peng, Z. Yang, L. Wang, J. Lin, L. Li, Z. Huang, B. Gong, Co-occurrence network of microbes linking growth and immunity parameters with the gut microbiota in Nile tilapia (*Oreochromis niloticus*) after feeding with fermented soybean meal, *Aquac. Reports*. 26(2022). <https://doi.org/10.1016/j.aqrep.2022.101280>.