

Review Article

Harnessing the Potentials of Sweetpotato (*Ipomoea batatas* L.) for Enhanced Livelihood and Development

Abstract

This review gives an overview of the potentials of sweetpotato as a versatile and multi-functional crop. It brings together careful reviews of some of the outstanding characteristics and revealing potentials of the crop which have been established and reported by several researches and authors. The contents of this review contains reviews on components such as the various types of sweetpotato based on the flesh color and their various constituent compounds, nutritive features, phenolic acids contained in sweetpotato, the phytochemical and pharmacological properties such as diseases' prevention and promotion of good health through its antioxidant, anticancer, antiulcer, anti-inflammation, cardiovascular properties and the enhancement of the immune system, processing of sweetpotato in to various food products such as vinegar, chips, flakes, cubes, yoghurt, cheese and some industrial utilization in the processing of alcohol and starch, the utilization potentials in animal feed; use as silage, processed to hay, as green-chop, the use of the tuber in poultry feeds and the contributions of the crop towards food security and development through the ability to fight hunger, poverty reduction, income generation, climate resilient and adaptability to vast range of climates. Various authors and researches have investigated and voted sweetpotato as one of the crops with the best production, consumption and utilization attributes which can help the smallholder farmers in the forest, tropical and subtropical zones to achieved food security and households' income stability. Sweetpotato is gaining attention and registering growing demands with appreciating growth in exports and imports in some countries across the globe due to its quality and potentials. The developing countries in particular which have the comparable advantages in the cultivation of sweetpotato should develop schemes and policies that will enable them to tap in to its potentials for livelihood improvement and development.

Keywords: Anthocynin, Carotene, Phytochemical, Pharmacological, Sweetpotato

1 Introduction

Sweet potato is a root-tuber crop which is rated as the 5th and 7th most principal food crop in the less developed countries and globally respectively [1]. Sweetpotato comes next after cassava among the root and tuber crops, making it the second most valuable tuber crop in the world [2]. Sweetpotato is the most prominent among the members of the *Ipomoea* genus, and it is by far the only member in the genus with the characteristics of bearing root tubers of great nutrition and economic values [3]. Sweetpotato is regarded as one of the important crops grown for man's consumption across the globe. Sweetpotato, among other crops is of economic advantage and has been rated as the 15th and the 29th cultivated crop on the account of the amount of yield and the area under cultivation respectively, with the average annual yield of 92, 000000 tons [4-6].

Sweetpotato is arguably an important tuber crop in most areas, cultivated in over 100 countries across the globe and has been regarded as a staple food for the people in the tropical and subtropical regions, which provides nutritional security for both the urban and rural dwellers through enhanced cultivation and utilization [7, 8]. Sweetpotato is grown around the world in about 114 nations, with the global average yield of 14 tons/hectare [9]. Sweetpotato among other staple crops is well adaptable to wide range of climates, rapid and short maturity and promising yield capacity [9].

2 Sweetpotato types, nutritional and chemical constituents

2.1 Types of sweetpotato according to flesh color

There are different types of sweetpotato varieties growing across the globe, with variation in skin and flesh color such as white, cream, orange, purple, pale yellow, etc. these types of sweetpotato do not only vary in the fleshed or skin color, but also possess different levels and kinds of important chemical constituents of enviable relevance in nutrition, medicine and other industrial usage.

Table 1: Types of sweetpotato according the fleshed color and their major compounds

Sweetpotato type	Major bioactive compound contained	Reference
------------------	------------------------------------	-----------

Purple fleshed	Anthocyanins, phenolic acids	[10, 11]
Orange fleshed	β -5 cryptoxanthin, β -carotene, α -carotene, flavonoids	[8,12]
White fleshed	Starch, phenolic acids	[13, 11]
Cream fleshed	Starch, phenolic acids	[13, 11]

2.2 Nutritional components of sweetpotato

Sweetpotato is a rich source of various important nutrient ingredients such as protein, energy, carbohydrates, Lipids, minerals and vitamins [14].




Figure 1: Nutritional composition of the tubers of sweetpotato [15].

2.3 Phenolic acids present in sweetpotato

The various varieties of sweetpotato generally contain a number of phenolic acids which are importantly linked to the unique sensory traits possess by the varieties.




Figure 2: Phenolic acids present in sweetpotato [16, 17].




Figure 3: Phytochemical constituents of sweetpotato [18, 19].

3 Processing, benefits and potentials of sweetpotato

3.1 Processing of sweetpotato

The tubers of sweetpotato can be used to produce starch, which is indigenously used to produce noodles, sugar, vermicelli, syrups, contributing significantly to food products. Additionally the starch and the sugar obtained from it can also be used to make products such as monosodium, citric acid, microbial enzymes, glutamate, lactic acid and alcohol [9, 39]. Sweetpotato are also used in the manufacturing of snacks and bakery foods like chips [40]. The purple and orange fleshed varieties of sweetpotato are reported to have been extensively applied in the making of inartificial anthocyanin colorants and β -carotene in some beverage products on a large scale in Japan [9]. The purple and orange fleshed sweetpotato are used in making beer and wine, due to their characteristics of being rich sources of starch which is convertible to sugar [9]. Sweetpotato provides good industrial raw materials for the producing of alcohol and starch [41].

Maintaining and scaling up the growth and expansion of the sweetpotato sub-sector through processing and formulation of food products is directly proportional to the commercialization and value addition of sweetpotato, due to its richness in polyphenols and dietary fiber, which have the potentials of reducing the digestibility of starch [42, 43].




Figure 4: Food and Industrial products from sweetpotato [9].

Alpha-amylase: the beta-amylase enzyme, oxidase causes the breaking down of the starch component of sweetpotato during processing [44], and the enzymatic reaction of the amylase results to maltose formation during processing leading to the high level of sweetness and sugar content of the end product formed [45].

The Mitsui and Toyota motor companies are collaboratively exploring the potential of sweetpotato in the manufacturing of plastics materials which are bio-degradable, and has also projected sweetpotato as a potential efficient energy source like hydrogen and alcohol in the day ahead of us [46].

3.2 The pharmacological and health potentials of Sweetpotato

There are several therapeutic, pharmacological, phytochemicals and other health related attributes associated with Sweetpotato, which have been studied and their relevance being investigated. Sweetpotato is an excellent source of several phytochemical compounds of

therapeutic importance. Nutritionally, the various parts of sweet potato have great dietary fiber, the stem and leaves have high carotene, vitamin E, vitamin C, vitamin B2 and iron [20]. The sweet potato roots are good sources of vitamin C, starch, β -carotene and sugar [21]. It is reported that the tubers of sweetpotato have high protein level than that of yam and cassava [22].

Table 2: Pharmacological properties of sweetpotato

Property	Mechanism of reaction	References
Anti-oxidant action	Flavonoids and n-hexane, ethanol and ethyl-acetate extracts from leaves help to remove oxygen-based reactive radicals in the cell. Reduces the chances of getting degenerative disorders like senile, dementia, cancer, diabetes, asthma, which are highly caused by the actions of free harmful radicals.	[23, 24] [25]
Boost of immune system	The leaves of purple Sweetpotato regulate the activities of lytic, T-lymphocyte deleterious cell. Generation of antibody production in healthy humans The derivatives of ethyl-acetate from Sweetpotato have immune-modulatory actions	[26] [27]
Anti-cancer property	The polyphenol constituent of sweetpotato extract hinders the proliferation and also leads to apoptosis in prostate cancer cells The anthocyanin in the sweetpotato with purple flesh is potent against the development of cancers of the breast and stomach.	[28, 29]
Anti-diabetic property	The white-fleshed sweetpotato increases the adiponectin level in the blood, a hormone that modifies the metabolism of insulin.	[30] [31]

	Extracts of white-fleshed sweetpotato potentially reduces blood sugar level	
Antimicrobial property	Extracts of ethanol and acetone obtained from the leaves of sweetpotato respectively have antimicrobial inhibitory actions on <i>Pseudomonas aeruginosa</i> and <i>Salmonella typhimurium</i>	[32] [30]
	Liquid extracted from the leaf of sweetpotato highly inhibits the growth of <i>E. coli</i> , <i>Bacillus cereus</i> and <i>Staphylococcus aureus</i>	
Cardiovascular enhancing property	Extract rich in phytochemicals, obtained from the leaves of sweetpotato inhibits the oxidation of less dense lipoprotein in humans.	[33] [34]
	Anthocyanin in purple skinned sweetpotato decreases the chances of getting coronary infections.	
Anti-ulcer property	Extract rich in methanol, obtained from the roots of sweetpotato protects the stomach of rat against ulcer, induced by aspirin.	[35] [36]
	The flour obtained from the roots of sweetpotato inhibits gastric ulcer induced by ethanol.	
Anti-inflammatory property	Inhibition of inflammatory infection of the brain by preventing the inflammatory reactions caused by lipopolysaccharide.	[31]
Haematological property	The leaf of sweetpotato has haematinic action, and it is used in indigenous treatment of anaemia.	[37] [38]
	Powder from the leaf of sweetpotato increases the levels of red blood cells and	

3.3 The uses of sweetpotato in feed and feed ingredients of animals

The vegetative parts and the tuber of sweetpotato are used as silage and fodder for livestock [47]. There was improvement in the growth of young female cattle when fed with roughage diet, supplemented with silage prepared from the vines of sweetpotato [48]. According to Mibach *et al.* [49] substituting of ground-corn for any amount of sweetpotato flour for sheep put on diets of multiple constituents does not interfere with the intake and digestibility of dry matter, nitrogen retention and microbial synthesis of protein. According to Murugan *et al.* [50] averagely 43% of the sweetpotato produced per annum globally is utilized as feed for animals; the vines and other foliar parts of sweetpotato are commonly used as protein supplementary feed and feed materials for farm animals like pigs in several countries across the globe such as Uganda, Korea, Taiwan, Vietnam, Indonesia, Papua New Guinea, China, Philippines and India.

The tubers of sweetpotato have been studied and identified as one of the nutritious and less expensive alternative feed materials for broiler chickens [51, 52]. The addition of about 25% and 20% of the orange-fleshed sweetpotato tuber-meal to layer and broiler chicken feeds respectively do not have any negative impact on the quality of eggs and growth [53]. Sweetpotato-meal inclusive diet has proven to have an enhancing influence on the growth and performance of quails in Japan [54]. The most desirable feed conversion efficiency and growth can be achieved in chicken, when fed with diet containing about 10% of dried sweetpotato-leaves, and can result in more weight gain in broilers in particular compared with other diets containing no sweetpotato leaves [55]. A diet of dried leaves of sweetpotato can increase the level of the yellow color of the egg yolk and the skin of chickens [56]. The meal prepared from the tubers of the orange-fleshed sweetpotato has been suggested with proven results, to be the best substitute for corn at the rate of 12% -16% and 15% and 20% during the starter and grower stages of chicken respectively, with positive influence on chicken characteristics such as ingestion, weight increase and meat attributes [57]. Meal prepared from sweetpotato tubers can be included at the levels of 25% and 50% in the diets of layer and broiler chickens respectively to achieve optimal production [58].

3.4 The potentials of sweetpotato in fostering food security

According to Scott and Maldonado [59] Sweetpotato is largely cultivated in low and middle income earning countries, and because of that, scaling up of sweetpotato cultivation is noted as a greater tool to foster food security and improve the livelihoods and living standards of the economically disadvantaged class of the rural and urban settings.

Sweetpotato plays several fundamental functions, contributing significantly to the “food basket” of the globe, which has a viable potential of reducing the food need of the growing world’s population, decreases poverty, promotes food security and improves livelihoods [2]. Sweetpotato tubers are great hunger-combating food materials which significantly saved millions of people during some daring moments in the history of the world such as the period critical food insecurity at china (in the 1960s), Uganda (during the 1990s) and Japan, that is the time during which there was a development of phytosanitary problems in fields of cassava in Uganda and the destructions of rice farms by typhoons in Japan [60, 2].

In addition to food, Sweetpotato has got multi-industrial uses because of its wonderful attributes such as cheap and simple cultivation, ability to tolerate infections and pests, high output and rich in carbohydrate, minerals and vitamins in both its roots and leaves [61]. The vitamin A precursor (Beta- carotene) rich orange fleshed sweetpotato can importantly contributes to the global fight against the deficiency of vitamin A among people, which has started yielding positive results in improving status quo of vitamin A among lactating and expectant mothers and children in Africa [9, 62].

In the populous semi-arid plains of Eastern Africa, sweetpotato is a cherishable food material used to combat food insecurity by thousands of rural communities [63]. The development and demands of snack-foods developed from sweetpotato have seen a surge in recent years, covering a significant portion of the food commodity market, meeting the taste and preference of consumers [64]. Sweetpotato cultivation is very profitable, with high sale-value and has the potentials of earning its growers more incomes [65].

Okoye [66] has noted that sweetpotato has a good comparative advantage in Nigeria, in terms of its production and small holder farmers earn significant amount of income from the vines, tubers and the leaves. In terms of nutrients and biomass per hectare, sweetpotato is very excellent and tops all other crops worldwide, and as such it is considered as one of the best crops with the

capacity to fight hunger and poverty [63]. Various primary and secondary processed food products such as puree, flour, chips, noodles, pasta and biscuits can be made from the tubers of sweetpotato [67], and therefore, it can be investigated in advance level as a key crop for the formulation of sustainable nutritionally resourced food items.

The tubers of the purple fleshed type of sweetpotato contribute to the keeping of the body stable; they can best substitute rice in our food menu due to their richness in carbohydrate, and also contain more nutrients than the white rice [68]. Ice cream is made from sweetpotato [69] and the purple type sweetpotato can best be used in place of corn in terms of starch for bread making [70].



Figure 5: The roles of sweetpotato in promoting food security

4 Future Research focus and prospects

Sweetpotato (*Ipomoea batatas* L.) is noted globally as one of the great cultivated crops among the roots and tubers, being a cheap and sustainable source of nutrition. It has several underutilized health advantages, market and export potentials and above all, the crop and its products possess greater prospects in the areas of sustainable food, nutrition and income security

for every player along its value chain in Africa, most particularly in the Sub-Saharan African sub region where majority of the rural peasant farming populations are bedeviled with acute hunger, income instability and poverty. The starch of sweetpotato is of promising industrial importance, and can be explored vastly by research to develop more starch production efficient varieties for expansion of the processing industry. The molecular, chemical and mineral richness as well as the medicinal potentials of the edible portions of sweetpotato could be exploited in to detail and well incorporated in to infant foods, other dietary packages for children to improve health and reduce the incidence of infantile minerals and nutritional deficiencies related diseases. The crop is naturally endowed with anthocyanin, anticancer, anti-inflammatory, antibacterial and numerous other disease fighting properties, which can be further explore in medical and pharmaceutical products such as capsules, mixtures, ointment for enhance administering and use. Currently, less priority is given to research focus on the development of more industrially efficient varieties, especially for starch, medicinal and pharmaceutical purposes. Some national and regional agricultural institutions in Africa have demonstrated little interests and weak commitment towards investment in the sweetpotate sub-sector by helping rural and urban growers to expand production and helping to create the enabling local and international market spaces to fully exploit the market and export opportunities of the crop and its related products. The crop is climate change efficient, cheap in production wise and sustainable in income and food stability, which the least developed nations with the comparable advantage in its production can, tapped to improve livelihoods, enhance standard of living, expand and increase export opportunities in international trades as a tool for ensuring rural development and the growth of national economies

5 Conclusions

This comprehensive review has revealed some of the promising potentials of sweetpotato. These characteristics show how multipurpose and versatile the crop is. It has the capacity to fight hunger, employment and income generations, reduction in poverty and raising the standard of living in the. Sweetpotato has the capacity to improve lives and promote development if the potentials along its value chains from production to processing and utilizations are well harnessed. It is nutritionally rich and can be processed industrially and locally in to several food products and feed forms. Scientific research should direct maximum attention to the development of more industrial and processing efficient varieties. It should be well adopted and its prospects and potentials harnessed for development.

References

- [1] R. Mohanraj, Sweetpotato: Bioactive compounds and health benefits, *In the Bioactive Molecules in Food. Reference Series in Phytochemistry*; Mérillon, J. M., Ramawat, K. G., Eds., Springer International Publishing, Cham, Switzerland, pp.1-16, 2018.
- [2] A.F. El-Sheikha, R.C. Ray, Potential impacts of bioprocessing of sweetpotato: Review, *Critical Review of Food Science and Nutrition*, Vol.57, pp.455-471, 2017.
- [3] C.E. Cartabiano-Leite, O.M. Porcu, A.F. De Casas, Sweet potato (*Ipomoea batatas* L.) nutritional potential and social relevance: a review, *International Journal of Engineering, Research and Applications*, ISSN: 2248-9622, Vol.10, No.6, pp.23-40, 2020.
- [4] F.A.O.STAT. Food and Agriculture Organization Statistics, “Food and Agricultural Association of the United Nations Database,” 2018.
- [5]. F.A.O.STAT. Food and Agriculture Organization Statistics, Food and Agricultural Association of the United Nations – Database, 2019.

- [6] T.H. Mu, P.G. Li, Chapter 2 - Sweet potato: origin and production, Sweet Potato, *Academic Press*, pp.5-25, 2019.
- [7] M.K. Alam, A comprehensive review of sweet potato (*Ipomoea batatas* L.) Lam): Revisiting the associated health benefits, *Trends in Food Science and Technology*, Vol.115, pp.512-529, 2021.
- [8] M.K. Alam, Z.H. Rana, S.N. Islam, Comparison of the proximate composition, total carotenoids and total polyphenol content of nine orange-fleshed sweet potato varieties grown in Bangladesh, *Foods*, Vol.5, No.4, pp.64, 2016.
- [9] V.D. Truong, R.Y. Avula, K.V. Pecota, G.C. Yencho, Sweetpotato Production, Processing, and Nutritional Quality,” Handbook of Vegetables and Vegetable Processing, Volume II, Second Edition. Edited by Muhammad Siddiq and Mark A. Uebersax. *John Wiley and Sons Ltd.*, USA, 2018.
- [10] C.C. Teow, V.D. Truong, R.F. McFeeters, R.L. Thompson, K.V. Pecota, G.C. Yencho, Antioxidant activities, phenolic and β -carotene contents of sweetpotato genotypes with varying flesh colours, *Food Chemistry*, Vol.103, pp.829-838, 2007.
- [11] M.S. Padda, D.H. Picha, Quantification of phenolic acids and antioxidant activity in sweetpotato genotypes, *Science of Horticulture*, Vol.119, pp.17-20, 2008.
- [12] S.Y. Park, S.Y. Lee, J.W. Yang, J.S. Lee, S.D. Oh, S. Oh, S.M. Lee, M.H. Lim, S.K. Park, J.S. Jang, *et al.* Comparative analysis of phytochemicals and polar metabolites from colored sweet potato (*Ipomoea batatas*, L.) tubers, *Food Science and Biotechnology*, Vol.25, pp.283-291, 2016.
- [13] S. Tumwegamire, R. Kapinga, P.R. Rubaihayo, D.R. LaBonte, W.J. Grüneberg, G. Burgos, T.Z. Felde, R. Carpio, E. Pawelzik, R.O.M. Mwanga, Evaluation of dry matter, protein, starch, sucrose, β -Carotene, Iron, Zinc, Calcium, and magnesium in East African sweetpotato (*Ipomoea batatas* L.) germplasm, *Horticultural Science*, Vol.46, pp.348-357, 2011.

- [14] U.S.D.A. United States Department of Agriculture, Agricultural Research Service, Nutrient Data Laboratory, *USDA National Nutrient Database for Standard Reference*, Release 28, 2015.
- [15] S.W. Fanta, S. Neela, Review on Nutritional Composition of Orange-fleshed Sweet Potato and Its Role in Management of Vitamin A Deficiency, *Food Science and Nutrition*, Vol.7, pp.1920-1945, 2019.
- [16] J. Musilová, J. Bystrická, J. Árvay, L. Harangozo Polyphenols and phenolic acids in sweet potato (*Ipomoea batatas*, L.) roots, *Potravin. Slovak Journal of Food Science*, Vol.11, pp.82-87, 2017.
- [17] R.G.O. Rumbaoa, D.F. Cornago, I.M. Geronimo, Phenolic content and antioxidant capacity of Philippine sweet potato (*Ipomoea batatas* L.) varieties, *Food Chemistry*, Vol. 113, pp.1133-1138, 2009.
- [18] R. Mohanraj, S. Sivasankar, Sweet potato (*Ipomoea batatas* L.) a valuable medicinal food: A Review, *Journal of Medicine and Food*, Vol.17, pp.733-741, 2014.
- [19] J. Woolfe, Sweet Potato: An Untapped Food Resource, *Cambridge University Press*, Cambridge, UK, pp.134-157, 1992.
- [20] H. Ishida, H. Suzuno, N. Sugiyama, S. Innami, T. Tadokoro, A. Maekawa, Nutritive evaluation on chemical components of leaves, stalks and stems of sweet potatoes (*Ipomoea batatas* poir), *Food Chemistry*, Vol.68, Issue.3, pp.359-67, 2000.
- [21] A.F. Sanoussi, A. Dansi, H. Ahissou, A. Adebawale, L.O. Sanni, A. Orobiyi, M. Dansi, P. Azokpota, A. Sanni, Possibilities of sweet potato (*Ipomoea batatas* L.) Value chain upgrading as revealed by physico-chemical composition of ten elites landraces of Benin, *African Journal of Biotechnology*, Vol.15, No.13, pp.481-9, 2016.
- [22] B.O. Oloo, A. Shitandi, S. Mahungu, J.B. Malinga, B.R. Ogata, Effects of lactic acid fermentation on the retention of beta carotene content in orange fleshed sweet potatoes, *International Journal of Food Studies*, Vol.3, pp.13-33, 2014.

- [23] Y. Tang, W. Cai, B. Xu, Profiles of phenolics, carotenoids and antioxidative capacities of thermal processed white, yellow, orange and purple sweet potatoes grown in Guilin, China, *Food Science and Human Wellness*, Vol.4, pp.123-132, 2015.
- [24] H. Ji, H. Zhang, H. Li, Y. Li, Analysis on the nutrition composition and antioxidant activity of different types of sweet potato cultivars, *Food and Nutrition Science*, Vol. 6, pp.161, 2015.
- [25] T.M. Florence, The role of free radicals in disease *Australian and New Zealand Journal of Agriculture, Food Chemistry and Ophthalmology*, Vol.23, pp.3-7, 1995.
- [26] Y.C. Chen, H.S. Chang, H.M. Lai, S.T. Jeng, Characterization of the wound inducible protein IPOMOELIN from sweet potato, *Plant Cell Environment*, Vol.28, pp.251-259, 2005.
- [27] C.M. Chen, S.C. Li, Y.Y. Liao, J.F. Liu, J. F., Shyur, P.R. Chen, C.Y. Chen, In vitro immunomodulatory effect of *Ipomoea batatas* L. tuber bioactives, *The FASEB Journal*, pp.862-869, 2013.
- [28] M. Sugata, C.Y. Lin, Y.C. Shih, Anti-inflammatory and anticancer activities of Taiwanese purple-fleshed sweet potatoes (*Ipomoea batatas* L. Lam) extracts, *BioMed Research International*, Vol.2015, Article ID 768093, pp.10, 2015.
- [29] L.S. Wang, G.D. Stoner, Anthocyanins and their role in cancer prevention, *Cancer Letters*, Vol.269, pp.281-290, 2008.
- [30] B. Ludvik, M. Hanefeld, G. Pacini, Improved metabolic control by *Ipomoea batatas* (Caiapo) is associated with increased adiponectin and decreased fibrinogen levels in type 2 diabetic subjects, *Diabetes Obesity and Metabolism*, Vol.10, pp.586-592, 2008.
- [31] T.B. Ayeleso, K. Ramachela, E. Mukwevho, A review of therapeutic potentials of sweet potato: Pharmacological activities and influence of the cultivar, *Tropical Journal of Pharmaceutical Research*, Vol.15, No.12, pp.2751-2761, 2016.

- [32] V.B. Adsul, E. Khatiwora, R. Torane, N.R. Deshpande, Antimicrobial activities of *Ipomoea carnea* leaves, *Journal of Natural Product and Plant Resources*, Vol. 2, Issue.5, pp.597-600, 2012.
- [33] M. Nagai, M. Tani, Y. Kishimoto, M. Iizuka, E. Saita, M. Toyozaki, K. Kondo Sweet potato (*Ipomoea batatas* L.) leaves suppressed oxidation of low density lipoprotein (LDL) in vitro and in human subjects, *Journal of Clinical Biochemistry and Nutrition*, Vol.48, pp.203, 2011.
- [34] G. Mazza, Anthocyanins and heart health, *Ann I Super Sanita*, Vol.43, pp.369, 2007.
- [35] V. Panda, M. Sonkamble, Anti-ulcer activity of *Ipomoea batatas* L. tubers (sweet potato), *Functional Foods in Health and Disease*, Vol.2, pp.48-61, 2012.
- [36] R. Sathish, M. Rani, K. Natarajan, Study of ulcer protective effect of *Ipomoea batatas* L. dietary tuberous roots (Sweet Potato), *Iranian Journal of Pharmacology and Therapeutics Collection*, Vol.11, pp.36-39, 2012.
- [37] E.O. Osime, G.E. Ediale, C.E. Omoti, A.A. Famodu, Effect of sweet potato leaf (*Ipomoea batatas* L.) extract on some haematological parameters using rabbits, *Journal of Medicine and Biomedical Research*, Vol.7, No.68, pp.12-15, 2008.
- [38] J.F. Montejo, J.A.B. Mondonedo, M.G.A. Lee, M.B. Ples, R.J.S. Vitor, Hematological effects of *Ipomoea batatas* (camote) and *Phyllanthus niruri* (sampa-sampalukan) from Philippines in the ICR mice (*Mus musculus*), *Asian Pacific Journal of Tropical Biomedicine*, Vol.5, pp.29-33, 2015.
- [39] G. Padmaja, Uses and nutritional data of sweetpotato, *In the Sweetpotato (Loebenstein G, Thottapilly G, editors, Springer, Dordrecht, Netherlands, pp.189-233, 2009.*
- [40] J.G. Waramboi, M.J. Gidley, P.A. Sopade, Influence of extrusion on expansion, functional and digestibility properties of whole sweetpotato flour, *Food Science and Technology*, Vol.59, pp.1136-1145, 2014.

- [41] Y.B. Karan, O.G. Şanlı, The assessment of yield and quality traits of sweet potato (*Ipomoea batatas* L.) Genotypes in middle Black Sea region, Turkey, *PLoS ONE*, Vol.16, No.9, e0257703, 2021.
- [42] S. Wang, S. Nie, F. Zhu, Chemical constituents and health effects of sweet potato, *Food Research International*, Vol.89, pp.90-116. 2016.
- [43] F. Zhu, Interactions between starch and phenolic compound, *Trends in Food Science and Technology*, Vol.43, pp.129-143, 2015.
- [44] V. Truong, R.Y. Avula, Sweet potato purees and powders for functional food ingredients, *Sweetpotato: postharvest aspects in food*, pp.117-61, 2010.
- [45] T. Kiribuchi, K. Kubota, Studies on cooking of sweetpotato (part 1) changes in sugar content and beta-amylase activity during cooking, *Kaseigaku Zasshi*, Vol.27, No.6, pp.418-22, 1976.
- [46] S. Islam, Nutritional and Medicinal Qualities of Sweetpotato Tops and Leaves. Plant Science, *In the Cooperative Extension Program, University of Arkansas at Pine Bluff.*, Arkansas, FSA6135, 2014.
- [47] I. Etela, G.A. Kalio, Yields components and 48-h rumen dry matter degradation of three sweet potato varieties in Ndama steers as influenced by date of harvesting, *Journal of Agriculture and Social Research*, Vol.11, Issue.2, pp.15-21, 2011.
- [48] A.I.M. Ali, S.E. Wassie, R.G. Joergensen, D. Korir, J.P. Goopy, K. Butterbach-Bahl, L. Merbold, U. Dickhoefer, E. Schlecht, Feed Quality and Feeding Level Effects on Faecal Composition in East African Cattle Farming Systems, *Animals*, Vol.11, pp.564, 2021.
- [49] M. Mibach, C.F. Demarco, A. A. Barbosa, L. De Oliveira, M. N. Corrêa, F.A.B.D. Pino, V.R. Rabassa, E. Schmitt, G.V. Kozloski, C.C. Brauner, Sweet Potato Flour as a Replacement for Ground Corn as an Energetic Concentrate, *Animal Production*, ISSN e1678-4596. *Ciência Rural, Santa Maria*, Vol.51, No.10, e20200838, 2021.

- [50] S. Murugan, S.K. Paramasivam, M. Nedunchezhiyan, Sweet Potato as Animal Feed and Fodder. Global Science Books, *Fruit, Vegetable and Cereal Science and Biotechnology*, Vol.6, Issue.1, pp.106-114, 2012.
- [51] R.C. Beckford, J.R. Bartlett, Inclusion levels of sweet potato root meal in the diet of broilers I. Effect on performance, organ weights, and carcass quality, *Poultry Science*, Vol.94, Issue.6, pp.1316-1322, 2015.
- [52] O. Mozafari, S. Ghazi, M.M. Moeini, The effects of different levels of edible potato (*Solanum tuberosum*) replacing maize on performance, serum metabolite and immune system of broiler chicks, *Iran Journal of Animal Science*, Vol.3, Issue.3, pp.583-588, 2013.
- [53] C.O. Okereke, E. Oti, E.C. Nwauzor, Effect of dietary inclusion of orange-fleshed sweet potato meal on egg quality of laying hens, *In the Proceedings of 14th Annual Conference of Animal Science Association of Nigeria*, Nigeria, pp.382-384, 2009.
- [54] J.A. Edache, U. Musa, M.U.M Ehizokale, J.O. Esilonu, E.J. Okpala, P.D. Karsin, A. Yisa, N.J. Zwandor, Replacement value of sweet potato (*Ipomea Batatas* L.) Meal for maize in practical diets fed to quail (*Cotumix cotumix japonica*) chicks, *Nigeria Journal of Animal Production*, Vol.36, No.11, pp.34-40, 2009.
- [55] W. Tsega, B. Tamir, The effect of increasing levels of dried leaves of sweet potato (*Ipomoea batatas*) on dry matter intake and body weight gain performance of broiler finisher Chickens, *Livestock Research for Rural Development*, Vol.21, Article #208, 2009.
- [56] V. Heuzé, G. Tran, P. Hassoun, Sweet Potato (*Ipomoea batatas* L.) Forage, *Feedipedia.org*, 2013.
- [57] K. Mosebework, T. Negesse, A. Nurfeta, Effect of Replacing Maize (*Zea Mays* L.) with Orange Fleshed Sweet Potato Tuber (*Ipomoea Batatas* L.) on Performance of Cobb 500 Broiler Chickens, *Agricultural Research & Technology: Open Access Journal*, Vol.13, No.5, pp.555-895, 2018.

- [58] T. Tagesse, Review on Feed Resources with More Emphasis on Use of Cassava and Sweet Potato Meal in Chicken Diet: Ethiopia, *Animal and Veterinary Sciences*, Vol.9, Issue.2, pp. 32-38, 2021.
- [59] G.J. Scott, L. Maldonado, Sweetpotato for the New Millennium: Trends in Production and Utilization in Developing Countries, *CIP Program Repository*, Vol.98, pp.329-335, 1998.
- F.C. Amagloh, B. Yada, G.A. Tumuhimbise, F.K. Amagloh, A.N. Kaaya, The Potential of Sweetpotato as a Functional Food in Sub-Saharan Africa and Its Implications for Health: A Review, *Molecules*, Vol.26, pp.2971, 2021.
- [60] D. Bach, A.C. Bedin, L.G. Lacerda, A. Nogueira, I. M. Demiate, Sweet Potato (*Ipomoea batatas* L.): a Versatile Raw Material for the Food Industry. Review-Food/Feed Science and Technology, *Brazilian Archives of Biology and Technology*, Vol.64, pp.1678-4324, 2021.
- [61] F. Rozi, N. Prasetyaswati, D.A.A. Elisabeth, Study on sweet potato market behavior in supporting food security, *In the IOP Conference Series, Earth and Environmental Science*, No.756, 012082, 2021.
- [62] J.W. Low, M. Arimond, N. Osman, B. Cunguara, F. Zano, D. Tschirley, A food-based approach introducing orange-fleshed sweetpotato increased vitamin A intake and serum retinol concentrations in young children in rural Mozambique, *Journal of Nutrition*, Vol.137, pp.1320-1327, 2007.
- [63] J. Low, J. Lynam, B. Lemaga, C. Crissman, I. Barker, G. Thiele, S. Namanda, C. Wheatley, M. Andrade, Chapter 16, Sweetpotato in Sub-Saharan Africa. G. Loebenstein, G. Thollappilly (eds.), the Sweetpotato © *Springer Science-Business Media* B.Y., pp.359, 2009.
- [64] T.H. Mu, H.N. Sun, M.M. Ma, Sweet potato snack foods, *Chemistry, Processing and Nutrition*, pp.303-324, 2019.

- [65] B. Dereje, A. Girma, D. Mamo, T. Chalchisa, Functional properties of sweet potato flour and its role in product development: a review, *International Journal of Food Properties*, Vol.23 Issue.1, pp.1639-1662, 2020.
- [66] A.C. Okoye, Gender Deferential in market competitiveness among sweetpotato smallholder farmers in South Eastern Nigeria, *A Ph.D. dissertation presented in the Department of Agribusiness and Management, Michael Okpara University of Agriculture, Umudike, Nigeria*, pp.10-65, 2018.
- [67] N.A. Giri, B. Ambedkar, B.K. Sakhale, Sweet potato (*Ipomoea batatas* L.): A valuable tropical tuber crop: A review. 182, *The Pharmaceutical Innovation Journal*, Vol.8, Issue.6, pp.182-191, 2019.
- [68] H. Wijaya, A. Slay, N. Abdullah, Ice cream products made from processed purple sweet potatoes: a product organoleptic study, *In the IOP Conference Series, Earth and Environmental Science*, No.807, 042074, 2021.
- [69] D.R. Affandi, S. Handajani, Es krim ubi jalar ungu (*Ipomoea batatas* L.): Tinjauan sifat sensoris, fisik, kimia, dan aktivitas antioksidannya *J. Teknol. Has. Pertan*, 4, 2011.
- [70] R.M. Anugrah, E. Suryani, Kandungan Gizi Donat dengan Penambahan Ubi Ungu (*Ipomoea batatas* L.), *Sebagai Makanan Jajanan Berbasis Pangan Lokal Bagi Anak Sekolah J. Gizi*, 9, pp.150-158, 2020.
- [71] T.B.M. Lirag, Determinants of Profitability of Sweet Potato Production in Camarines Sur, Philippines, *International Journal of Advanced Science, Engineering, Information and Technology*, Vol.9, pp.467-472, 2019.
- [72] J.O. Virgínoe Silva, M.F. Almeida, M. Da Conceição Alvim-Ferraz, J.M. Dias, Integrated production of biodiesel and bioethanol from sweet potato. *Renew, Energy*, Vol.124, pp.114-120, 2018.
- [73] N.N. Mudege, S. Mayanja, T. Muzhingi, Women and men farmer perceptions of economic and health benefits of orange fleshed sweet potato (OFSP) in Phalombe and Chikwawa districts in Malawi, *Food Security*, Vol.9, pp.387-400, 2017.

- [74] S. Neela, S.W. Fanta, Review on nutritional composition of orange-fleshed sweet potato and its role in management of vitamin A deficiency, *Food Science and Nutrition*, Vol.7, pp.1920-1945, 2019.
- [75] S. Laurie, M. Faber, P. Adebola, A. Belete, Biofortification of sweet potato for food and nutrition security in South Africa, *Food Research International*, Vol.76, pp.962-970, 2015.
- [76] F.A.O. Food and Agriculture Organization, Land and Water, 2020.
- [77] U.N.I.D.O. United Nations Industrial Development Organization, Baseline Report of Clean Cooking Fuels in the East African Community (EAC) Region, pp.2-34, 2020.
- [78] P. Sakai, S. Afionis, N. Favretto, L.C. Stringer, C. Ward, M. Sakai, P.H. Weirich Neto, C.H. Rocha, J. Alberti Gomes, N.M. De Souza, *et al.* Understanding the Implications of Alternative Bioenergy Crops to Support Smallholder Farmers in Brazil, *Sustainability*, Vol.12, pp.2146, 2020.
- [79] M.H. Su, C.H. Huang, W.T. Li, C.T. Tso, H.S. Lur, Water footprint analysis of bioethanol energy crops in Taiwan, *Journal of Cleaner Production*, Vol.88, pp.132-138, 2015.