

**Metabolic Syndrome and its association Cardiometabolic risk factors;
Prevalence among apparently healthy adults in a rural community in
southwestern Nigeria**

Abstract

Aim: The study was carried out to assess the prevalence of cardiometabolic risk factors associated with metabolic syndrome among apparently healthy adults in rural settlements.

Study design: Descriptive cross-sectional study with

Place and Duration: Ejigbo, a rural settlement in Osun State, Southwestern Nigeria, between September and December, 2019.

Methodology: 271 apparently healthy individuals were recruited using a multistage sampling technique with the WHO STEPS Instrument questionnaire, Level of significance set at $P= 0.05$

Results: The prevalences of overweight and obesity were 96 (35.4%) and 66 (24.4%) respectively. The prevalence of hypertension among respondents was 83 (30.6%). High plasma levels of Total Cholesterol 64 (23.6%), triglyceride 20 (7.4%), LDL-Cholesterol 28 (10.3%) and very high LDL-Cholesterol 31 (11.5%) were found among the respondents. While 72 (26.6%) of the respondents have a low level of HDL-Cholesterol. The prevalence of impaired glucose tolerance among the respondents was 34 (12.6%) while 45 (16.6%) had diabetic plasma glucose levels and 82 (30.3%) for metabolic syndromes.

Conclusion: Cardiometabolic risk factors, were found to be very high among rural dwellers, especially people without metabolic syndromes. Public awareness about those risk factors should be intensified. Routine medical check-ups and screening should be encouraged.

Keywords: Cardiometabolic, Metabolic syndromes, Pre-hypertension, Impaired glucose

Introduction

Metabolic syndrome is the aggregation of several known cardiovascular risk factors. These include insulin resistance, obesity, atherogenic dyslipidemia and hypertension. These conditions have interrelated mechanisms, pathways and underlying mediators.¹ It is a worldwide epidemic disorder with a high socio-economic impact and due to its association with increased morbidity and mortality.² The main causes are increasing urbanization, nutrition changes, and reduced physical activity. Metabolic syndrome consists of several metabolic and physiological abnormalities, including abdominal obesity, impaired glucose tolerance, hypertriglyceridemia, decreased high-density lipoprotein cholesterol (HDL-C), and arterial hypertension. This syndrome increases the risk of developing type 2 diabetes mellitus by twofold and cardiovascular diseases (CVDs) by fivefold compared with apparently healthy persons.^{1,2,3}

It estimated by National Health and Nutrition Examination, that approximately 30% of overweight and 60% of obese men and women meet the criteria for a diagnosis of Metabolic syndrome.³ It is estimated by the Framingham study that approximately 80% of essential hypertension in men and 65% in women directly attributed to obesity.⁴ Hypertension has been observed as one of the major risk factors associated with cardiovascular diseases, and a component of metabolic syndrome.⁵ In individuals with uncontrolled blood pressure, the prevalence of metabolic syndrome and type 2 diabetes is significantly increased compared with those with controlled blood pressure.⁶ Each component of the Metabolic syndrome is an independent risk factor for developing cardiovascular disease and producing a spectrum of

vascular and cardiac diseases.³ The mean overall prevalence of metabolic syndrome in Nigeria is 31.7%, 27.9% and 28.1% according to the World Health Organization (WHO), Adult Treatment Panel III (ATP III) and International Diabetes Federation (IDF) criteria, respectively ⁷

The prevalence and distribution of its components depend largely on the geographical location, prevailing lifestyle, diets and the extent of modernization.^{1,8} and most studies were in urban communities and among age group ≥ 35 years.^{7, 9} Thus, the study was carried out in a rural settlement to assess the prevalence of cardiometabolic risk factors associated with metabolic syndrome among apparently healthy adults.

Materials and Methods

It was a cross-sectional study conducted at Ejigbo, a rural settlement in Osun State, Southwestern Nigeria. Apparently healthy 271 individuals were recruited into the study using a multistage sampling technique. Adults who have lived in Ejigbo for at least one year included in the study. Individuals with a history of cardiovascular diseases, cancer, type 2 diabetes mellitus, renal diseases, pregnancy, and taking medications for hypertension or dyslipidemia excluded from the study.

In the first stage, out of the 30 Local Government Areas (LGAs) in Osun State, Ejigbo was selected from the list by simple random method (balloting technique). The Second stage involved the selection of 6 wards out of the existing 11 electoral wards from a list obtained from the LGA secretariat using a simple random sampling method (balloting technique). The third stage all households with eligible respondents in each of the selected electoral wards, were then identified. They were instructed on overnight fasting, questionnaires administered. Where a

household has more than one eligible respondent, one of them was chosen via a simple random sampling method.

WHO STEPS Instrument (core and Expanded) questionnaires administered to obtain information from each participant. Anthropometric measurements were taken. About 10mls of fasting blood samples were taken following an aseptic procedure for phlebotomy into two separate bottles – lithium heparin and fluoride oxalate. The samples were centrifuged at x3000g, separated and the plasma obtained was stored at -4°C in the laboratory. The stored samples were analyzed in batches using ready-to-use commercial kits manufactured by Randox Laboratories Ltd, Crumlin, County Antrim United Kingdom. All the analyses were done following the manufacturer's instructions and along with controls.

Lipid parameters – High-density lipoprotein cholesterol (HDL- C), Triglycerides and Total cholesterol (TC) were analyzed using the enzymatic method employed by the kits.¹⁰ Low-density lipoprotein cholesterol (LDL-C) was calculated by the use of the Friedewald equation.¹¹ Fasting plasma glucose was determined using enzymatic oxidation in the presence of glucose oxidase, urea by the Urease-Berthelot method, uric acid by enzymatic colourimetric method and creatinine by the modified Jaffe Method.^{12,13} Lipid profile was categorized according to Adult Treatment Panel III (ATP III) classification.¹⁴ while blood pressure was classified into normal, pre-hypertension and hypertension according to Joint National Committee (JNC 7).¹⁵

In this study, International Diabetic Foundation IDF criteria for metabolic syndrome were employed to define the presence of the syndrome in the study population because of its flexibility in requiring obesity and not necessarily insulin resistance but still retained other criteria like the NCEP ATP III definition. The key features are hyperglycemia/insulin resistance,

visceral obesity, atherogenic dyslipidemia and hypertension, and at least any three of the five criteria define metabolic syndrome.¹ The Statistical Package for Social Sciences (SPSS) version 23 (SPSS Inc., Chicago, IL, IBM Version) was used for entry and analysis of the data obtained. Univariate analysis of all the variables measured was first carried out. Data were presented using frequency distribution tables and charts. Association between metabolic syndrome and other categorical variables assessed using a chi-square. For every cell with an expected value less than 5, Fisher's Exact Test was used to determine the statistical significance. In the multivariate analysis, a stepwise model of binary logistic regression analysis was done to determine the predictors of future metabolic syndrome. Variables imputed into the logistic model were selected based on their level of significance during bi-variate analysis. Adjusted odds ratio and 95% confidence interval were obtained to identify determinants of metabolic syndrome. The level of significance was set at $P = 0.05$ for this study.

Ethical approval was obtained from Bowen University Teaching Hospital's (BUTH) ethical review committee. The respondents were informed about the nature of the study and that participation was completely voluntary. Written informed consent was obtained from all selected respondents before recruitment into the study. All information gathered was kept confidential and participants were identified using only serial numbers.

Results

Table1 Socio-economic status of the respondents (n=271)

Variable	Frequency	Percentage (%)
Age (years)		
Range: 18 – 92		
<40	38	14.0
40 – 49	40	14.8
50 -59	49	18.1
60 – 69	73	26.9
≥70	71	26.2
Sex		
Male	110	40.6
Female	161	59.4
Education		
Less Secondary School	141	52.0
Secondary or More	130	48.0
Marital Status		
Currently married	177	65.3
Currently unmarried	94	34.7
Ethnic Group		
Yoruba	266	98.2
Igbo	3	1.1
Others	2	0.7

The youngest and oldest in the study population were 18 years and 92 years respectively. Majorities were 60 years and above; 73 (26.9%), and 71 (26.9%) respectively. Only 38 of the study participants representing 14% were below 40 years. Over half of the respondents were females 161(54.4%) compared with males; 110 (40.6%). About 141(52%) of the study participants had less than a secondary school education. 177 (65.3%) were currently married while 94 (34.7%) were single. Nearly all 266 (98.2%) respondents were from the Yoruba ethnic group.

Table 2: Distribution of Nutritional and Cardiometabolic risk factors among the respondents (N=271)

Variables	Frequency	Percentage (%)
Body Mass Index (Kg/m²)		
Underweight	7	2.6
Normal	102	37.6
Overweight	96	35.4
Obesity	66	24.4
Waist Height Ratio		
Low risk (<0.5)	70	25.8
High Risk (≥0.5)	201	74.2
Waist Hip Ratio		
Low Risk	93	34.3
High Risk	178	65.7
Waist Circumference (cm)		
Normal	113	41.7
At risk	53	19.6
Significant Risk (abdominal Obesity)	105	38.7
Blood Pressure (mmHg)		
Systolic		
Normal	70	25.8
Pre-HT	100	36.9
HT	101	37.3
Diastolic		
Normal	130	48.0
Pre-HT	69	25.5
HT	72	26.6
Combined		
Normal	120	44.3
Pre-HT	68	25.1
HT	83	30.6
Total Cholesterol (mmol/L)		
Desirable	159	58.7
Borderline	48	17.7
High	64	23.6
Triglyceride (mmol/L)		
Optimal	191	70.4
Normal	33	12.2
Borderline	27	10.0
High	20	7.4
LDL-Cholesterol (mmol/L)		
Optimal	117	43.2
Near-Optimal	48	17.7
Borderline	47	17.3
High	28	10.3

Very High	31	11.5
HDL-Cholesterol (mmol/L)		
Optimal	132	48.7
Borderline	67	24.7
Risk factor	72	26.6
Fasting Blood Glucose (mmol/L)		
Normal	192	70.8
Impaired glucose tolerance	34	12.6
Diabetes mellitus	45	16.6
Urea (mmol/L)		
Normal Level	253	93.4
Abnormal	18	6.6
Creatinine (mmol/L)		
Normal	231	86.2
Abnormal	37	13.8
Uric Acid (mmol/L)		
Normal	269	99.2
Abnormal	2	0.8

The prevalences of overweight and obesity were 96 (35.4%) and 66 (24.4%) respectively. The majority of the respondents 201 (74.2%) and 178 (65.7%) had high risk according to Waist Height Ratio and Waist Hip Ratio respectively. 105 (38.7%) had significant risk (abdominal Obesity) while 53 (19.6%) were at risk using their Waist Circumference. Pre-hypertensive category among the study participants were 100 (36.9%), 69 (25.5%) and 68 (25.1%) for systolic, diastolic and combined measurements respectively. The prevalence of hypertension among respondents was 101 (37.3%), 72 (26.6%) and 83 (30.6%) for systolic, diastolic and combined respectively. Prevalence of dyslipidemias was 64 (23.6%), 20 (7.4%), 28 (10.3%) and 31 (11.5%) for high plasma levels of total Cholesterol, triglyceride, LDL-Cholesterol and very high LDL-Cholesterol respectively. While 72 (26.6%) of HDL-Cholesterol were at risk. The prevalence of impaired glucose tolerance among the respondents was 34 (12.6%) while 45 (16.6%) had diabetic plasma glucose levels. A significant proportion of the respondents had normal plasma levels of urea 253 (93.4), creatinine 231 (86.2) and uric acid 269 (99.2).




Figure 1 Prevalence of Metabolic Syndrome (n=271)

Figure 1 shows a prevalence of 82 (30.3%) for metabolic syndrome.

UNDER PEER REVIEW

Table 3: Factors associated with Metabolic syndrome at the bivariate analysis level (N = 271)

Variable	Metabolic Syndrome n (%)		p-value
	Absent (n = 189)	Present (n = 82)	
Age (in years)	58.00 ± 16.00	59 ± 14.00	0.534 ^a
Sex			0.002*
Male	88 (80.0)	22 (20.0)	
Female	101 (62.7)	60 (37.3)	
Education			0.147 ^b
Less Secondary School	104 (73.8)	37 (26.2)	
Secondary or More	85 (65.4)	45 (34.6)	
Marital Status			0.877 ^b
Currently married	124 (70.1)	53 (29.9)	
Currently unmarried	65 (69.1)	29 (30.9)	
Ethnic Group			0.052 ^b
Yoruba	186 (69.9)	80 (30.1)	
Igbo	3 (100)	0 (0)	
Others	0 (0)	2 (100)	
BMI			<0.001*^b
Underweight	7 (100)	0 (0)	
Normal	89 (87.3)	13 (12.7)	
Overweight	70 (72.9)	26 (27.1)	
Obesity	23 (34.8)	43 (65.2)	
Waist-to-height ratio			<0.001*^b
Low risk (<0.5)	66 (94.3)	4 (5.7)	
High Risk (≥0.5)	123 (61.2)	78 (38.8)	
Waist-hip ratio			0.005*^b
Low Risk	75 (80.6)	18 (19.4)	
High Risk	114 (64.0)	64 (36.0)	
Waist circumference			<0.001*^b
Normal	106 (93.8)	7 (6.2)	
At risk	42 (79.2)	11 (20.0)	
Abdominal Obesity	41 (39.0)	64 (61.0)	
Blood Pressure			<0.001*^b
Systolic			
Normal	63 (90.0)	7 (10.0)	
Pre- Hypertension	69 (69.0)	31 (31.0)	
Hypertension	57 (56.4)	44 (43.6)	
Diastolic			0.001*^b
Normal	105 (80.8)	25 (19.2)	
Pre- Hypertension	42 (60.9)	27 (39.1)	
Hypertension	42(58.3)	30 (41.7)	
Combined			<0.001*^b
Normal	108 (90.0)	12 (10.0)	

Pre-Hypertension	35 (51.5)	33 (48.5)	
Hypertension	46 (55.4)	37 (44.6)	
Fasting blood sugar			<0.001*^b
Normal	160 (83.3)	32 (16.7)	
Impaired glucose tolerance	14 (41.2)	20 (58.8)	
Diabetes mellitus	15 (33.3)	30 (66.7)	
Total cholesterol (mmol/L)	5.13 ± 1.63	5.29 ± 2.04	0.493 ^a
Triglyceride (mmol/L)	0.91 ± 0.54	1.30 ± 1.07	<0.001*^a
HDL-C (mmol/L)	1.69 ± 0.61	1.27 ± 0.68	<0.001*^a
LDL-C (mmol/L)	2.96 ± 1.46	3.35 ± 1.66	0.057 ^a
Creatinine (µmol/L)	89.65 ± 38.17	81.73 ± 32.76	0.103 ^a
Urea (mmol/L)	3.45 ± 1.72	3.45 ± 1.42	0.967 ^a
Uric Acid (mmol/L)	0.344 ± 0.37	0.32 ± 0.12	0.595 ^a

* statistically significant ^a t-test for independent samples used ^b Pearson chi-square test used

There was no significant difference between the mean age among respondents for the absence of 58±16 and the presence of 59±14 of metabolic syndromes. More female respondents 60 (37.3%) had metabolic syndromes compared to males 22 (20%) and the difference was significant, (P= 0.002). There was a significant difference in the BMI for the absence and presence of metabolic syndrome for overweight (70; 72.9% vs 26; 27.1%) and obesity (23; 34.8% vs 43; 65.2%) P-value <0.001. Presence and absence of metabolic syndromes for Waist-to-height ratio low risk (66; 94.3% vs 4 (5.7%); high risk (123; 61.2% vs 78; 38.8%), P-value =0.001 and Waist-hip ratio low risk (75; 80.6% vs 18; 19.4%); high 114; 64% vs 64; 36%), P-value=0.005. A significant difference was observed among participants with the presence and absence of metabolic syndromes for different categories of Waist circumference; normal (7;10% vs 106; 93.8%), At risk (11;20% vs 42;79.2%) and abdominal obesity (64; 61.0%vs 41;39.0%), p-value <0.001. In blood pressure, a statistically significant percentage of the participants have a metabolic syndrome for pre-hypertensive and hypertensive categories; systolic (31;31.0% and 44;43.6%), diastolic (25; 19.2% and 30; 44.7%) and combined (33; 48.5% and 37; 44.6%), p<0.001. A significant percentage of participants with impaired glucose tolerance and diabetic

Mellitus have metabolic syndromes 20 (58.5%) and 30 (66.7%) respectively when compared with normal plasma glucose level 32 (16.7%), P-value<0.001. The plasma level of triglyceride for participants with metabolic syndrome was greater than that of participants with the absence of metabolic syndrome (1.30 ± 1.07 vs 0.91 ± 0.54) and the difference was significant, P-value<0.001. The plasma level of HDL-C for participants with metabolic syndrome was significantly less than those of the participants with the absence of metabolic syndrome (1.27 ± 0.68 vs 1.69 ± 0.61), P-value <0.001. Various degrees of differences were observed in the plasma level of total cholesterol, LDL-C, creatinine, urea and uric acid among participants with metabolic syndrome when compared with participants without metabolic syndrome but these differences were not significant.

Table 4: Predictors of Metabolic Syndrome using binary logistic regression analysis

Variable	Adjusted Odds Ratio	P-Value	95% CI	
			Lower	Upper
Age	0.98	0.314	0.935	1.022
Sex				
Male (R)	1	1		
Female	1.29	0.763	0.245	6.819
Waist circumference				
Normal (R)	1	1		
At risk	0.72	0.775	0.075	6.936
Abdominal Obesity	75.65	0.001*	6.158	929.202
Fasting blood sugar				
Normal (R)	1	1		
Impaired glucose tolerance	57.07	<0.001*	7.160	454.906
Diabetes mellitus	63.07	<0.001*	8.246	482.298
Systolic blood pressure				
Normal (R)	1	1		
Pre-hypertension	26.05	0.001*	3.510	193.291
Hypertension	27.7	0.002	3.334	230.096
Diastolic blood pressure				
Normal (R)	1	1		
Prehypertension	1.79	0.403	0.457	7.020
Hypertension	2.19	0.350	0.424	11.305
Waist-to-height ratio				
Low risk (R)	1	1		
High Risk	5.92	0.185	0.424	11.305
Waist-hip ratio				
Low risk (R)	1	1		
High risk	0.707	0.655	0.155	3.222
Body mass index				
Normal (R)	1	1		
Overweight	0.566	0.448	0.130	2.459
Obesity	2.207	0.324	0.457	10.663
Total cholesterol	1.242	0.871	0.090	17.182
Triglyceride	9.543	0.011*	1.678	54.290
LDLC	0.764	0.838	0.058	10.058
HDLC	0.046	0.038*	0.003	0.845
Creatinine	0.989	0.173	0.972	1.005
Urea	0.924	0.608	0.684	1.248
Uric acid	1.431	0.616	0.353	5.806

* statistically significant

The binary logistic regression analysis shows that the respondents with high triglyceride levels had about 10 times the likelihood of developing metabolic syndrome compared with those who had normal plasma levels (OR:9.54; CI:1.68-54.26).

Discussion.

The presence of metabolic syndrome increases the risk and mortality of cardiovascular diseases.¹⁶ This study focused on the risk factors associated with metabolic syndrome in a rural community. The age range for this study was between 18 - 92 years and the majority of the study participants were above 60 years. The majority of the respondents were either overweight or obese and at high risk for developing cardiovascular diseases considering their Waist Height Ratio and Waist-hip ratio. It was similar to other studies that have raised concerns about the increase in cardiovascular risk factors in semi-urban and rural communities^{17, 18} largely due to changes in lifestyle and urbanization.^{19, 20} In the current study, the prevalence of pre-hypertension and hypertension was 100 (36.9%) and 101 (37.3%) respectively. An urban study by Ejiroghene and Henry in Delta State South-South Nigeria reported a higher prevalence of 42.64% for pre-hypertension but obtained a 29.3% prevalence for hypertension among their respondents.²¹ A similar result was observed from a systemic analysis by Davies *et al* which puts the prevalence at 30.9% and 30.6% for pre-hypertension and hypertension respectively.²² The high percentage of pre-hypertension in the community calls for concern. This on its own, is a cardiovascular risk factor and when left unattended to, will definitely increase the incidence of hypertension significantly in no time and consequently, increases morbidity and mortality.²¹ Therefore, the present level of awareness for pre-hypertension must be improved especially among rural dwellers with emphasis on routine medical check-ups.

In the pattern of dyslipidemias among the studied population, more people have low HDL-C (26.6%), followed closely by a high level of TC (23.6%) when compared with a very high level of LDL-C (11.5%) and triglyceride (7.4%). Previous studies have reported inconsistent findings; Odenigbo and Oguejilor in Asaba found a high prevalence for low HDL-C (60%), LDL-C (51%), TC (23%) and TG (5%) levels. Agboola-Abu and Onabolu in Okada reported TC (60.4%) and TG (5%) levels. In Owerri, Osuji et al found low HDL-C (37.6%), TG (34.1%) and TC (31.4%) levels. While Sani, *et al*, in Katsina also found HDL-C (59.3%), TC (28.3%), LDC-C (25.7%) and TG (15.0%) levels.^{23,24,25} All of these studies were conducted among apparently healthy individuals. The disparities between these previous studies and the current study may also be due to sociocultural differences in the study locations.^{26,27} The prevalence of impaired glucose tolerance was 12.6% while for individuals with diabetic glucose levels were 16.6%. Davies Adeloye et al obtained a pooled prevalence of 15.0% for impaired glucose tolerance and 5.8% for diabetics in a systematic review and meta-analysis study.²² ISI Ogbu et al in Owerri Imo State Nigeria observed a prevalence of 15.5% for impaired glucose tolerance and 8.6% for unreported diabetes.²⁷ Like, pre-hypertension above, the prevalence of impaired glucose tolerance is quite high. Impaired glucose tolerance had been shown by studies to be associated with increased cardiovascular morbidity and mortality.^{28,29,30}

In this study, the prevalence of metabolic syndrome was 30.3%. Findings from other studies with the same IDF definition; V.M.Oguoma *et al* in a systematic review obtained 31.3%, Yetunde et al in Ogun State 36.8%, 35.1% in North-Western Nigeria by Sabir *et al* while a lower prevalence of 21.7% was reported in South-west, Nigeria by Adejumo *et al*.^{7,26,31,32} The prevalence in this study, though conducted in a rural area, was similar to those obtained from studies done in urban

communities. The probable reason for this may be due to its proximity to big cities and the effects of urbanization.

There was no significant difference in the mean age between persons with metabolic syndromes and persons without metabolic syndromes. Though, metabolic syndrome was said to be more among the middle age and elderly respondents.³³ The prevalence of metabolic syndrome among females 60, (37.3%) was significantly higher than males 22, (20%). This finding was similar to other studies which found that the prevalence of metabolic syndromes was higher in females than males. This significant difference was partially due to the occurrence of risk factors for metabolic syndromes which was higher in females than males.^{33,34,35}

While comparing the risk factors for metabolic syndromes among persons that have already met the criteria and those that have not, it was surprising to find that most people that have not met the full criteria for metabolic syndromes have significantly high individual risk factors for metabolic syndromes. This category of respondents was also at high risk of the development of cardiovascular diseases. The presence of one risk factor, unattended to, is consequently an invitation for others, thus the risk factors have been described as a cluster.^{1,4}

There were significant differences in anthropometric parameters between those with and those without metabolic syndromes; Waist-to-height ratio, and Waist circumference with p-value < 0.001 while Waist-hip-ratio has a p-value of 0.005. Findings from several studies have shown anthropometric parameters as simple and the best tools or indicators of metabolic syndromes. These can also be used as reliable screening tools.^{36,37,38,39}

The study population were stratified into normal, pre-hypertension and hypertension using American Heart Foundation (AHF) criteria, significant number of people without metabolic

syndromes were pre-hypertensive and hypertensive when compared with people with metabolic syndromes. Hypertension is the most important and independent risk factor for metabolic syndrome. People with uncontrolled hypertension are at greater risk of developing cardiovascular diseases than people with controlled blood pressure.^{5,6}

The incidence of impaired glucose tolerance and diabetes mellitus was significantly high in individuals with metabolic syndromes. Diabetics were said to be more prevalent among all age groups and closely associated with other risk factors such as hypertension, obesity and dyslipidemia.^{40,41,42}. Impaired glucose is a pre-diabetic state and both together indicate insulin resistance which is at the central pathophysiology of metabolic syndromes.^{1,43}

Dyslipidemias is an integral part of metabolic syndromes components, especially low HDL-Cholesterol levels and hypertriglyceridemia.^{44,45} This was also observed in this study significantly low plasma levels of HDL-cholesterol and high levels of triglyceride among participants with metabolic syndromes.

Conclusion

The prevalence of metabolic syndromes was found to be similar to that of urban settlements but the prevalence of cardiometabolic risk factors was higher than expected especially among people without metabolic syndromes. This calls for concerns about the increase in cardiovascular risk factors among rural dwellers. Public awareness about those risk factors should be intensified and routine medical check-ups and screening should be encouraged by government policy and non-governmental organizations vis-a-vis available and subsidized/ free medical services.

Data Availability

Data can be made available upon request

References

1. Huang, P. L. A comprehensive definition for metabolic syndrome. *DMM Dis. Model. Mech.* **2**, 231–237 (2009).
2. Espósito RC, Medeiros PJ, Silva FS, Oliveira AG, Soares Aragão CF, Oliveira Rocha HA, Moreira SA, F. S. V. Prevalence of the metabolic syndrome according to different criteria I DMSO. *Diabetes, Metab. Syndr. Obes. Targets Ther.* **11**, 401–408 (2018).
3. Tune, J. D., Goodwill, A. G., Sassoon, D. J. & Mather, K. J. Cardiovascular consequences of metabolic syndrome. *Translational Research* vol. 183 57–70 (2017).
4. Mendizábal, Y., Llorens, S. & Nava, E. Hypertension in Metabolic Syndrome : Vascular Pathophysiology. *Int. J. Hypertens.* **2013**, (2013).
5. Hussein H. Alhawari, Sameeha Al-Shelleh, Hussam H. Alhawari, Aseel Al-Saudi, Dina Aljbour Al-Majali, Leen Al-Faris, S. A. A. Blood Pressure and Its Association with Gender, Body Mass Index, Smoking, and Family History among University Students. *Int. J. Hypertens.* **2018**, (2018).
6. Walter Zidek, Lisa Naditch-Brûlé, Stefano Perlini, Csaba Farsang, S. E. K. Blood pressure control and components of the metabolic syndrome the GOOD survey. *Cardiovasc. Diabetol.* **8**, (2009).
7. V.M.Oguoma, E.U.Nwoseb, R. S. R. Prevalence of cardio-metabolic syndrome in Nigeria a systematic review. *Public Health* **129**, 413–423 (2015).

8. Ce Tan, Yutaka Sasagawa, Ken-ichi Kamo, Takehiro Kukitsu, Sayaka Noda, Kazuma Ishikawa, N. Y. Evaluation of the Japanese Metabolic Syndrome Risk Score (JAMRISC) a newly developed questionnaire used as a screening tool for diagnosing metabolic syndrome and insulin resistance in Japan. *Environ. Health Prev. Med.* **21**, 470–479 (2016).
9. Nalado A. M., Musa B. M., Gezawa I. D., Muhammad H., Ibrahim D. A., Uloko A. E. Prevalence of Metabolic Syndrome Among Apparently Healthy Adults in a Rural Community, in North-Western Nigeria. *Niger. J. Med.* **24**, 323–330 (2015).
10. Portela Rosana Deyse Ponte, Neto Edilson Martins Rodrigues, Júnior Francisco Josimar Girão, Ferreira Jamile Magalhães, De Sousa Alves Renata, De Jesus Ponte Carvalho, Teresa Maria, H. B. A. P. Comparative analysis of two methodologies for determination of HDL cholesterol. *J. Young Pharm.* **10**, 308–312 (2018).
11. Guo-Ming Zhang, Shu-Mei Bai, Gao-Ming Zhang, Xiao-Bo Ma, H. G. A Novel Method for Estimating Low-Density Lipoprotein (LDL) Levels Total Cholesterol and Non-High-Density Lipoprotein (HDL) Can Be Used to Predict Abnormal LDL Level in an Apparently Healthy Population. *Med Sci Monit.* **24**, 1688–1692 (2018).
12. Ekun O. A., Ogunyemi G. A., Azenabor A., Akinloye O. A comparative analysis of glucose oxidase method and three point-of-care measuring devices for glucose determination. *Ife J. Sci.* **20**, 43 (2018).
13. Jumaah, I. A. M. Estimation of uric acid , urea , creatinine and creatinine clearance in the serum of preeclamptic women ريدقتض ماد ك لوبلا , ايروي , نينتايرك و تيفصت نينتايرك ي فلصم ءاسنلا . *Kerbala J. Pharm. Sci.* 183–189 (2012).
14. Bermúdez Valmore J., Bello Luis Miguel, Naguib Ariana, Añez Roberto J., Fortul

- Yanalith et al. Lipid profile reference intervals in individuals from Maracaibo, Venezuela: An insight from the Maracaibo City Metabolic Syndrome prevalence study. *Rev. Latinoam. Hipertens.* **7**, 24–34 (2012).
15. Gyamfi Daniel, Obirikorang Christian, Acheampong Emmanuel, Danquah Kwabena Owusu, Asamoah Evans Adu. et al. Prevalence of pre-hypertension and hypertension and its related risk factors among undergraduate students in a Tertiary institution, Ghana. *Alexandria J. Med.* **54**, 475–480 (2018).
16. Ziolkiewicz Anna, Peter T. Ellison, Susan F. Lipson, Inger Thune, Jesienka. G. Body fat, energy balance and estradiol levels : a study based on hormonal profiles from complete menstrual cycles. *Hum. Reprod.* **23**, (2008).
17. Dahiru T., Ejembi C. L. Clustering of cardiovascular disease risk factors in semi-urban population in northern Nigeria. *Niger. J. Clin. Pract.* **16**, 511–516 (2013).
18. Gyawali Bishal, Mishra Shiva, Raj Ghimire, Saruna Hansen, Martin Rune et al. P. The burden and correlates of multiple cardiometabolic risk factors in a semi-urban population of Nepal: a community-based cross-sectional study. *Sci. Rep.* **9**, 1–10 (2019).
19. Ntentie FR, Azantsa KBG, Ngondi JL, Santy EV1, Dimodi HT, M. A. Urbanization and Metabolic Syndrome in Cameroon: Alertness on Less Urbanised Areas. *Endocrinol. Metab. Syndr.* **03**, (2014).
20. Nowicki Grzegorz Józef, Ślusarska Barbara, Naylor Katarzyna, Prystupa Andrzej, Rudnicka-Drożak Ewa, Halyuk Ulyana, P. P. The relationship between the metabolic syndrome and the place of residence in the local community on the example of the janów Lubelski district in eastern Poland: A population-based study. *Diabetes, Metab. Syndr. Obes. Targets Ther.* **14**, 2041–2056 (2021).

21. Umuerri Ejiroghene, Aiwuyo H. O. Prevalence and correlates of prehypertension and hypertension among adults in Delta State, Nigeria: A cross-sectional community-based study. *Ghana Med. J.* **54**, 48–57 (2020).
22. Adeloje Davies, Owolabi Eyitayo O., Ojji Dike B., Auta Asa, Dewan Mary T., et al. Prevalence, awareness, treatment, and control of hypertension in Nigeria in 1995 and 2020: A systematic analysis of current evidence. *J. Clin. Hypertens.* **23**, 963–977 (2021).
23. Oguejiofor OC, Onwukwe CH, O. C. Dyslipidemia in Nigeria Prevalence and pattern *Ann Afr Med. Ann Afr Med* **11**, 197–202 (2012).
24. Adeloje D., Abaa D. Q., Owolabi E. O. Prevalence of hypercholesterolemia in Nigeria: a systematic review and meta-analysis. *Public Health* **178**, 167–178 (2020).
25. EnajiteI Okaka, E. B. O. Prevalence and pattern of dyslipidemia in a rural community in Southern Nigeria. *African J. Med. Heal. Sci.* **12**, 80 (2013).
26. Sabir Anas Ahmad, Jimoh Abdulgafar Iwuala, Sandra Omozehio Isezuo, Simeon Alabi Bilbis, Lawal Suleiman Aminu, Kaoje Umar Abubakar et al. Metabolic syndrome in urban city of North-Western Nigeria: Prevalence and determinants. *Pan Afr. Med. J.* **23**, 1–7 (2016).
27. Ogbu ISI, Azodo EC, C. A. Prevalence of pre-diabetes and unreported diabetes mellitus in population aged 45 years and above in Owerri Municipality, Imo State Nigeria. *J. Coll. Med.* **17**, 31 (2013).
28. Kim Hong Kyu, Kim Chul Hee, Kim Eun Hee, Bae Sung Jin, Choe Jaewon et al. Impaired fasting glucose and risk of cardiovascular disease in Korean men and women: The Korean Heart Study. *Diabetes Care* **36**, 328–335 (2013).
29. Tsuchida, K. *et al.* Impaired glucose tolerance and future cardiovascular risk after

- coronary revascularization: a 10-year follow-up report. *Acta Diabetol.* **57**, 173–182 (2020).
30. Huang Yuli, Cai Xiaoyan, Mai Weiyi, Li Meijun, Hu Yunzhao. Association between prediabetes and risk of cardiovascular disease and all-cause mortality: Systematic review and meta-analysis. *BMJ* **355**, (2016).
 31. Adeyemi Yetunde Aderonke, Onabanjo Oluseye Olusegun, Sanni Silifat Ajoke, Ugbaja Regina Ngozi, Afolabi David Olaulowa et al. Prevalence of metabolic syndrome among apparently healthy adults in Ogun state, Nigeria. *Nutr. Food Sci.* **47**, 780–794 (2017).
 32. Adejumo Esther, Ogundahunsi Omobolanle, Adejumo Olusola, Sotunsa John, J. O. Prevalence of Metabolic Syndrome in a Rural and Urban Community in South-West Nigeria Using Three Different Definitions. *Int. J. Trop. Dis. Heal.* **24**, 1–9 (2017).
 33. Jiang Boren, Zheng Yanjun, Chen Yingchao, Chen Yi, Li Qin et al. Age and gender-specific distribution of metabolic syndrome components in East China: Role of hypertriglyceridemia in the SPECT-China study. *Lipids Health Dis.* **17**, 1–11 (2018).
 34. Yang-Mo Young, Byung-Cheul Shin, Son Chihyoung, In-Hyuk Ha. An analysis of the associations between gender and metabolic syndrome components in Korean adults: A national cross-sectional study. *BMC Endocr. Disord.* **19**, (2019).
 35. Sangjin Lee, Young Ko, Chanyeong Kwak, Eun-Shil Yim. Gender differences in metabolic syndrome components among the Korean 66-year-old population with metabolic syndrome BMC Geriatrics Full Text. *BMC Geriatr.* **27**, (2016).
 36. Adejumo Esther Ngozi, Adejumo Adedeji Olusola, Azenabor Alfred, Ekun Ayodele Oloruntoba, Enitan Seyi Samson, Adebola Olayimika Kehinde, O. O. A. Anthropometric parameter that best predicts metabolic syndrome in Southwest Nigeria. *Diabetes Metab.*

- Syndr. Clin. Res. Rev.* **13**, 48–54 (2019).
37. Brudecki Janusz, Chtzanowska M. Anthropometric indicators as predictors of the risk of metabolic syndrome in adult working men. *Anthropol. Rev.* **78**, 67–77 (2015).
 38. Wu Lihong, Zhu Wenhua, Qiao Qiaohua, Huang Lijuan, Li Yiqi, C. L. Novel and traditional anthropometric indices for identifying metabolic syndrome in non-overweight/obese adults. *Nutr. Metab.* **18**, (2021).
 39. Guo X, Ding Q, Liang M. Evaluation of Eight Anthropometric Indices for Identification of Metabolic Syndrome in Adults with Diabetes. *Dovepress* **14**, 1431—1443 (2021).
 40. James Merlit, Varghese Treesa P., Vijayan Anjaly, Anagha Pooleriveetil Padikkal, Muhas C. et al. Link between metabolic syndrome and diabetes mellitus: A pathophysiological implication. *Int. J. Sci. Technol. Res.* **9**, 4201–4206 (2020).
 41. Li Wenzhen, Wang Dongming, Wang Xiaojun, Gong Yanhong, Cao Shiyi et al. The association of metabolic syndrome components and diabetes mellitus: Evidence from China National Stroke Screening and Prevention Project. *BMC Public Health* **19**, 1–10 (2019).
 42. Genser Laurent, Casella Mariolo James Rossario, Castagneto-Gissey Lidia, Panagiotopoulos Spyros, Rubino Francesco. Obesity, Type 2 Diabetes, and the Metabolic Syndrome: Pathophysiologic Relationships and Guidelines for Surgical Intervention. *Surg. Clin. North Am.* **96**, 681–701 (2016).
 43. Colak Ayfer, Diniz Gulden. Impaired Glucose Tolerance, Obesity and Inflammatory Mediators. *Glucose Toler.* (2012) doi:10.5772/52378.
 44. Ruotolo Giacomo, Howard B. V. Dyslipidemia of the metabolic syndrome. *Curr. Cardiol. Rep.* **4**, 494–500 (2002).

45. Halpern Alfredo, Mancini Marcio C., Magalhães Maria Eliane C., Fisberg Mauro, Radominski Rosana, Bertolami Marcelo C. et al. Metabolic syndrome, dyslipidemia, hypertension and type 2 diabetes in youth: From diagnosis to treatment. *Diabetol. Metab. Syndr.* **2**, 1–20 (2010).

UNDER PEER REVIEW