

Original Research Article

EFFECTS OF DIFFERENT POTTING MEDIA ON THE GERMINATION AND EARLY GROWTH OF *Newbouldialaavis*-(P.Beauv.) Seem.

ABSTRACT

The research was conducted at the screen house of the Department of Forestry and Wildlife, Faculty of Agriculture, Nnamdi Azikiwe University, Awka, Anambra state, Nigeria. The study was carried out from January 2021 to May 2021 to access the effect of different potting media on germination and early growth of *Newbouldialaavis*. Treatments were laid out in a Complete Randomize Design (CRD) with four replications. The germination parameters measured were: days to germination, germination period and germination percentage while the growth variables measured includes plant height, collar diameter and number of leaves. The potting media used were T1 (topsoil and cow dung), T2 (topsoil and goat dung), T3 (topsoil and poultry dropping), T4(topsoil and sawdust), T5 control (topsoil only). Data collected were subjected to R software (R Core Team, 2014), using the following packages: plyr, car, gvlma, one-way tests, user friendly science, ggplot. Results obtained indicated that T1 had the highest germination percentage (85%) followed by T2 and T3 with germination percentages of 82% and 80% respectively. T4 had the poorest germination percentage (60%). The results also showed that the treatment had direct effect on the seedling early growth; the highest mean collar girth was observed in T1 with 1.85cm while the least was observed in T4 with 0.72 m. The highest mean height was observed in T2 with 22.15 cm while the least was observed in T4 with 7.23 cm. Also, the highest mean number of leaves was observed in T2 with 21leaves while the least was observed in T4 with 7 leaves. Based on the findings of these experiments it is recommended that for maximum growth and optimum germination of *Newbouldialaavis*, T1 and T2 should be used, sawdust had poor germination and growth performance and thereby is not recommended for the early growth of *Newbouldialaavis*seedlings.

Keywords: *Newbouldialaavis*; germination; early growth; potting media; seedlings.

INTRODUCTION

Comment [CES1]: <https://journalajraf.com/index.php/AJRAF/manuscript-submission>

Formatted: Indent: Hanging: 0.01", Space After: 0 pt

Comment [CES2]: 1. Original research papers: Papers that include original empirical data that have not been published anywhere earlier (except as an abstract). Null/negative findings and replication/refutation findings are also welcome. This type of paper normally should not exceed 25 double-spaced pages of text (including references) and should not contain more than 15 figures/tables. We advise a length of 3000-6000 words (including everything).

Comment [CES3]: Adequate all manuscript to the AJRAF guidelines and template. Citations and the figures should be rechecked.

Comment [CES4]: Recheck all he grammar, and verify the comments to improve the Research manuscript.

Formatted: Centered, Indent: Hanging: 0.01", Space After: 0 pt

Comment [CES5]: Review all the data and verify the double spaces between words, phrases and paragraphs. Redo the tables 2, 3 and 4. Check the tables and figures AJRAF template. Recheck the citations in the manuscript and the references list. Adequate them to the AJRAF template.

Formatted: Indent: Hanging: 0.01", Space After: 0 pt

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

Comment [CES6]: Insert a brief introduction and the main objective of the research.

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

Formatted: Font: (Default) Times New Roman, 12 pt, Italic

Formatted: Indent: Hanging: 0.01", Space After: 0 pt

Formatted: Font: Bold

Formatted: Font: (Default) Times New Roman, 12 pt, Highlight

Formatted: Font: (Default) Times New Roman, 12 pt, Highlight

Formatted: Font: (Default) Times New Roman, 12 pt, Highlight

Newbouldialaavis is a medium sized angiosperm in the Bignoniaceae family. It is native to tropical Africa. It is ever green, though its leaves turn somewhat dark purple during the cold seasons (Okeke, 2003). It is popularly known as the tree of life or fertility tree in Nigeria. Most significantly, it is known as a symbol of unity, family symbol of authority and a shrine for ancestral worship (Ehiagbonare and Onyebi (2009)). The plant species is known in English as tree of life or fertility tree. In Nigeria it is known in Yoruba as okoko, Hausa - aduruku, Tiv - Kontor, Benin - Ikhimi, Igbo-Ogirisi, Efik - obot, Urhoho - ogiriki (Gill, 1992). The plant species plays a role in phytomedicine, and the bark, leaves, and root are the parts mostly used. The isolated chemical constituents of the plant as reported by Gill (1992) are alkaloids, tannins and saponins. It is reported to be effective in the treatments of elephantiasis, syphilis, epilepsy, insomnia, convulsions, rheumatic swelling, piles, dysentery, constipation and psychosis (Usman and Osuji, 2007; Odugbemi, 2008).

The tree grows to a height of about 10m-20m high, 60-90cm in girth. Bark greyish to pale brown, fairly smooth, slash cream, fibrous. Branchlet short and twisted, with short knobby twigs often with clustered leaves at their ends. Leaves are up to 50cm long and usually with 3-5 pairs of leaflets. The common stalk being prominently swollen at the point of attachment of each pair, leaflets 15-20cm long by 5-10cm broad, elliptic to broadly elliptic, sometimes slightly oblanceolate, acuminate, cuneate rather leathery, dark green, glabrous, usually with coarse teeth but the margin sometimes entire with 5- 8 pairs of prominent lateral nerves, looped well away from the margin, leaflet stalk very short, stout. The flowers (Dec- Feb) are purplish- pink with darker stripes, trumpet shaped, up to 6cm long, closely crowded in almost spike-like panicles at the ends of short side stalks, calyx about 2.5cm long, splitting down one side. The fruits (plate 2 Figure 1A) usually mature and ripe between (Jan- Feb) having 22-30cm long, pendulous, surface dotted with purplish glands which are attractive to ants, packed with winged seeds (Plate 4 Figure 1B), flat, about 3.5cm long including the wing at each end.

The young leaves of the plants are crushed in little amount of water and the extract is used to treat eye inflammation and redness. It is also administered to stop vaginal bleeding in threatened abortion (Kargbo, 1982). Kargbo (1982) further reported that the species' leaf and root are used in the treatment of round worm infection, elephantiasis, dysentery, malaria, stomachic, migraines, convulsions and the bark is chewed, swallowed to relieve headache. The

Formatted: Indent: First line: 0.4"

Comment [CES7]: Old reference (2003). Please use more recent data. Priority to publication with less than 10 years (2012-2022).

This comment is reinforced for almost all the bibliographic references used in this research.

Use the scholar google and academic database to verify more publications about the species and the studied parameters.

Formatted: Highlight

Comment [CES8]: Adequate this all other citations to template of AJRAF.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

wood is pale brown, durable, evenly textured and hard and it tends to remain alive for a long time even after cutting it. This makes it viable for usage as boundary tree, woodworks, yam stakes, house posts, bridges and firewood.

Also, *Newbouldialaervis* has different symbols and meanings to different countries for example; some villages in Gabon and Ivory Coast plant the tree near the tombs to act as a protective talisman. The Ibibio and Efik people of Nigeria regard the tree as a symbol of their deities thus they tend to plant in their sacred places. The Igbo part of Nigeria refers to the *Newbouldialaervis* (ogilisi) tree as a sacred tree that has some spiritual powers hence, they usually plant it in front of a native doctor, chief's house, or sacred places (Ersan *et al.*, 2008). It is highly used in every traditional rite and as a sacred tree as it is commonly found in sacred places such as shrine and other traditional sacred places.

Newbouldialaervis is a very useful multipurpose tree which provide numerous services which includes wood, fuelwood, poles, medicinal and produce the oxygen we breathe, and serve as raw materials for many industrial products (Ajibesin, 2011). This plant species of great cultural importance and diverse phytomedicinal uses should not be left in the wild. As such this specie needs to be domesticated by using an appropriate media that would ensure both the ~~ermination~~germination and its early growth.

Potting media is a combination of organic and inorganic material mixed in an appropriate or desired proportion depending on the demand of the research. It serves as a medium in which the species would grow and support the root of the species as it grows with the supply of the content nutrient. The nutrient content of a potting media is dependent on the research as each media can be modified or limited. The potting media needs to supply plant with a means of support, good drainage, adequate air circulation and storage of water and nutrient.

According to ~~Juan-Wral~~ (2006), good potting media management is essential to the production of quality tree seedlings, since vigorous growths needed to face the seasonal hazards encounter on the field. Plant requires favorable potting media for ~~#its~~ growth.

To achieve its function, growing media used should have light-weight, good porosity, well-drained but with good water holding capacity, contained required nutrients, slightly acidic with good cation-exchange-capacity, able to maintain a constant volume when wet or dry, free of

Formatted: Highlight

Formatted: Font: Italic, Highlight

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

insects, diseases, and weed seeds, low in silt, clay and ash content, easily stored for long periods of time without changes in physical and chemical properties, and easily handled and blended.

The development of a healthy, good root system needs a media with these good physical properties. Any nutrient or chemical deficiencies in soil can be compensated or supplemented using good potting media.

Germination is usually the growth of a plant contained within a seed. Anozie and Oboho (2020), it leads to the formation of a young seedling, it is also the process of reactivation of metabolic machinery of the seed resulting in the emergence of radicle and plumule (Baskin, 2014).

Early growth of a plant begins after germination. The early stage of a plant is also known as seedling stage and it is one of the most critical phases during a plant's life history. Seedling survival not only exerts an important influence on the size, persistence, and genetic variability of plant populations, but also on the availability of required nutrients at the early stage of growth. (Kitajima & Fenner, 2000; Grime, 2001). People do not have interest on the germination of this species as they believe that trees are natural gift of nature as it grows naturally on its own without planting. This believes poses a threat on the existence of this species.

The tree is characterized with poor or slow rate of seedling growth and it is susceptible to insect attack. Several researchers have carried out research about the medicinal content of the leaves and bark as well as the sap. However, there is little or no information on the effect of different potting media on the germination and early seedling growth of this species in Nigeria. ~~Therefore~~ Therefore, there is the need to carry out this study that will effectively determine the best potting media that can positively increase germination percentage and enhance early seedling growth of this species.

Hence, this research was carried out to investigate the effect of different potting media on the germination and early growth of ~~Newbouldia~~ *Newbouldia laevis*.

A

B



Figure 1. *Newbouldialaavis* species, where (A): aspect of the species fruits and (B) seeds.

Formatted: Centered, Line spacing: 1.5 lines

Formatted Table

Formatted: Font: Italic

MATERIALS AND METHODS

Experimental Site/ Study Area

This study was carried out at the department of forestry and wildlife management, faculty of Agriculture, Nnamdi Azikiwe University, Awka. The Nnamdi Azikiwe University is located in the Eastern part of this country with the longitude of 7.11540E and latitude of 6.24850N. It is a tropical region has average annual temperature of the area of study is 26.3°C, and the rainfall pattern ranging ranges between 1828 mm 2002mm (Ezenwajiet al., (2013). It is a tropical region.

Experimental Design

Completely Randomized Design (CRD) was used for this study. There were five treatment used with four replicate each.

Seed Procurement and preparation

The seeds used in carrying out this study were procured from Nnamdi Azikiwe University, Awka premises. Matured ripe fruits of *Newbouldialaavis* were gathered from phenotypically superior mother tree within Nnamdi Azikiwe University, Awka. Afterwards, the fruit were

Formatted: Normal, Indent: Left: 0", Line spacing: 1.5 lines

opened for seeds extraction. The extracted seeds were air dried at room temperature for 6 hours and then wrapped in a paper to avoid drying up and losing viability as a result of prolonged exposure to the immediate environment and the seeds were planted immediately into the already prepared growing media.

Organic manure procurement and preparation

The poultry dropping, cow dung and goat dung that was used for this experiment were gotten from the animal farm at the Department of Animal Science, Nnamdi Azikiwe University, ~~Awka Anambra State~~. While the sawdust was gotten from saw mill located around EkeAwka Market, ~~Awka, Anambra State~~.

The poultry dropping, saw dust, cow dung, and goat dung were mineralized for two weeks. This was done by mixing them respectively with top soil in the ratio 2:1, one media: two top ~~soilsoils~~, after which the mixture were made moist but not water logged and covered with black nylon. This was done to enable microorganisms act on the media and in the process liberate the nutrients in the form that can be absorbed by the roots of the plants, increase aeration and light reflection. The mineralized media was then sieved and filled into ten 5x3 inches poly pots ready for germination.

Formatted: Indent: Hanging: 0.01"

Potting Media (Treatments) Used

- Treatment 1 (T1) = Topsoil and Cow dung;
- Treatment 2 (T2) = Topsoil and goat dung;
- Treatment 3 (T3) = Topsoil and poultry dropping;
- Treatment 4 (T4) = Topsoil and saw dust;
- Treatment 5 (T5) = Topsoil only (control).

Routine Maintenance of Experimental Plot

Daily watering of the germination poly pot (day and night) was done using watering can. Weeding was also carried out appropriately when necessary.

Data Collection

Data was collected based on germination and growth response assessment as highlighted below:

i. **Germination response:** the effect of different potting media on the germination was assessed by counting the number of seeds that germinated daily. The germinated seeds ~~was~~were counted and recorded from the date of first emergence until there was no more germination.

ii. **Growth response:** For growth assessment, the first three seedlings to germinate in each treatment per replicate were marked and used for growth assessment of the germinated seedlings.

The effect of different potting media on growth of *Newbouldialaavis* seedlings was evaluated by measuring some growth valuable at five weeks after germination. The growth parameters measured in weekly bases includes.

- i. plant height (using measuring tape or ruler);
- ii. Collar girth (collar girth at soil level was measured using veneer caliper;
- iii. Number of leaves (was determined by counting the number of leaves).

Data Analysis

Data were expressed as mean \pm standard deviation (SD) and then analyzed with R (R Core Team, 2014), using the following packages:plyr, car, gvlma, onewaytests, userfriendlyscience, ggplot2. Prior to the comparison of means, the global test for linear model validation was used to test for linear model assumptions. Since the data did not satisfy the linear model assumptions of normality and homogeneity of variance, they were subjected to the Welch ~~anova~~ANOVA followed by Games-Howell (YEAR) post hoc test. Statistical significance was determined at less than 5% probability level.

Formatted: Indent: First line: 0", Line spacing: 1.5 lines

Formatted: Indent: Hanging: 0.01", Line spacing: 1.5 lines

Comment [CES9]: Insert as a reference.

Formatted: Highlight

Expected Result

The study is expected to provide the result that will review the best growth media for best germination and early growth performance of *Newbouldialaevin* in respect to plant height, stem diameter, and leaf production using different potting media.

Soil physio-chemical analysis

The topsoil used in ~~these research~~this research was obtained from the Department of Forestry and wildlife nursery, located at the Faculty of Agriculture, ~~Nnamdi Azikiwe University Awka and~~ was subjected to soil analysis, the analysis which was aimed at accessing the availability of important soil chemical nutrients that are required for plant growth.

Formatted: Indent: Left: -0.01", Hanging: 0.01", Right: 0.05", Space After: 0 pt, Line spacing: 1.5 lines

RESULTS AND DISCUSSION

Comment [CES10]: Avoid use as keywords expressions/words already used in the manuscript title.

Formatted: Justified, Indent: Left: 0", First line: 0", Right: 0.06", Line spacing: 1.5 lines

RESULTS

GERMINATION

The type of germination exhibited by *Newbodialaevin* seeds was epigeal germination. The new seedlings emerged with the hypocotyle. There was no significant difference between the days to emergence as the whole treatment started germinating the same day which was five days after planting. The germination period in *Newbodialaevin* was relatively short as all the treatment completed their germination between 6-9 days (Table 1). Treatment T1 has the highest germination percentage (85%) followed by T2 (82%), T3 (80%), T5 (65%), T4 (60%) (Table 1 and Figure 2) (Fig. 1).

Formatted: Indent: First line: 0.39", Line spacing: 1.5 lines

Formatted: Font: Not Bold

Formatted: Font: Not Bold

Table 1: Germination parameters of *Newbodialaevin* in relation to treatment.

Treatment	Days to Germination	Germination Period (days)	Germination Percentage (%)
Cow dung (T1)	5	5	85
Goat dung (T2)	5	6	82
Poultry dropping (T3)	5	5	80
Sawdust (T4)	5	9	60
Control topsoil only (T5)	5	8	65

- Formatted: Font: Bold
- Formatted: Centered, Line spacing: 1.5 lines
- Formatted: Font: Bold
- Formatted: Font: 11 pt, Bold
- Formatted: Font: Bold
- Formatted: Font: 11 pt, Bold
- Formatted: Font: 11 pt, Bold
- Formatted: Font: Bold
- Formatted: Font: 11 pt, Bold
- Formatted: Font: Bold
- Formatted: Font: 11 pt, Bold
- Formatted: Font: Bold
- Formatted: Left, Line spacing: 1.5 lines
- Formatted Table
- Formatted: Centered, Line spacing: 1.5 lines
- Formatted: Centered, Line spacing: 1.5 lines
- Formatted: Centered, Line spacing: 1.5 lines
- Formatted: Left, Line spacing: 1.5 lines
- Formatted: Centered, Line spacing: 1.5 lines
- Formatted: Left, Line spacing: 1.5 lines
- Formatted: Centered, Line spacing: 1.5 lines
- Formatted: Left, Line spacing: 1.5 lines
- Formatted: Centered, Line spacing: 1.5 lines
- Formatted: Left, Line spacing: 1.5 lines
- Formatted: Centered, Line spacing: 1.5 lines
- Formatted: Font: Bold
- Formatted: Centered, Line spacing: 1.5 lines

Fig.1: Germination percentage of *Newbouldia laevis* under different treatments




Figure 2.1: Germination percentage of *Newbouldialaevis* under different treatments.

Analysis of growth variables

Data were expressed as mean \pm standard deviation (SD) and then analyzed with R (R Core Team, 2014), using the following packages; plyr, car, gvlma, onewaytests, userfriendlyscience, ggplot2. Prior to the comparison of means, the global test for linear model validation was used to test for linear model assumptions. Since the data did not satisfy the linear model assumptions of normality and homogeneity of variance, they were subjected to the Welch ANOVA followed by Games-Howell post hoc test. Statistical significance was determined at less than 5% probability level. Values are mean \pm standard error (SE) of replicate samples. Column that has means with the same letter are not significantly different from each other.

Plant height under different growth media

Table 2 shows the result of the descriptive statistics of plant height (cm) of *Newbouldialaavis* under different treatments. The result shows the mean diameter as 20.739583 \pm 6.309791, 22.154167 \pm 7.316653, 15.958333 \pm 4.189088, 7.227778 \pm 1.339421 and 14.666667 \pm 5.072371 for treatment T1, T2, T3, T4 and T5, respectively.

Table 2. Descriptive statistics for plant height under different growth media.

Seedling growth	Treatment	N	Mean	Std. Err	Std. de

Comment [CES1 1]: Use this template of Table and adequate all the tables 2, 3 and 4.

Formatted: Highlight

Formatted: Highlight

Formatted: Highlight

Seedling growth Treatment N Mean Std. Err Std. de

Variables

Plant	Cow dung (T1)	144	20.739583	6.309791	0.5258159
Height (cm)					
Goat dung (T2)	144	22.154167	7.316653	0.6097211	
Poultry droppings(T3)	144	15.958333	4.189088	0.3490907	
Sawdust (T4)	144	7.227778	1.339421	0.1116184	
Control (topsoil) T5	144	14.666667	5.072371	0.4226976	

Collar girth under different growth media

Table 3 shows the result of the descriptive statistics of collar girth (cm) of *Newbouldialaavis* under different treatments. The result shows the mean collar girth as 1.8486111 ± 0.7576174 , 1.8180556 ± 0.7761887 , 1.3583333 ± 0.6588292 , 0.7159722 ± 0.3638687 , and 1.3833333 ± 0.6317918 for treatment T1, T2, T3, T4 and T5 respectively.

Table 3. Descriptive statistics for collar girth under different growth media.

Formatted: Highlight

Formatted: Highlight

Seedling growth	Treatment	N	Mean	Std. Err	Std. de
-----------------	-----------	---	------	----------	---------

Variables

Collar girth (cm)	Cow dung (T1)	144	1.8486111	0.7576174	0.06313478
	Goat dung (T2)	144	1.8180556	0.7761887	0.06468239
	Poultry droppings (T3)	144	1.3583333	0.6588292	0.05490244
	Sawdust (T4)	144	0.7159722	0.3638687	0.03032239
	Control (topsoil T5)	144	1.3833333	0.6317918	0.05264931

Number of leaves under different growth media

Table4 shows the result of the descriptive statistics of number of leaves of *Newbouldialaervis* under different treatments. The result shows the mean diameter as 18.43056±9.549761, 20.63194±10.573583, 13.98611±6.230969, 6.75000±2.827191, and 15.94444±7.914559 for treatment T1, T2, T3, T4 and T5 respectively.

Table 4. Descriptive statistics for number of leaves under different growth media

Formatted: Highlight

Formatted: Highlight

seedling growth	Treatment	N	Mean	Std. Err	Std. de
-----------------	-----------	---	------	----------	---------

Variables

Number	Cow dung (T1)	144	18.43056	9.549761	0.7958134
of					
leaves	Goat dung(T2)	144	20.63194	10.573583	0.8811319
	Poultry droppings(T3)	144	13.98611	6.230969	0.519247
	Sawdust (T4)	144	6.75000	2.827191	0.2355992
	Control (Topsoil T5)	144	15.94444	7.914559	0.6595466

Analysis of variance

From the analysis of variance in Table 5, it shows that there is significant difference between the height and treatments since the probability (0.001) is less than the alpha value. This analysis also shows that the collar girth and number of leaves has a probability value of (0.01) which is less than the alpha value of (0.05). this shows that the collar girth and number of leaves are significant respectively.

Table 5. The result of ANOVA for the *Newbouldialaevis*.

Growth variables	Source of Variation	Sum of Squares	Df	Mean Square	F	Sig.
Height (cm)	Between Groups	20009.86	4	5002.46	180.7	.0001
	Within Groups	19793.82	715	27.68		
	Total	39803.68	719			
Collar girth (cm)	Between Groups	121.37	4	30.34	70.82	.001
	Within Groups	306.32	715	0.043		
	Total	427.67	719			
Leaf count	Between Groups	16323.77	4	4080.94	65.3	.001
	Within Groups	44681.33	715	62.49		
	Total	61005.1	719			

Height of *Newbouldialaevis* under different organic manure

Table-6 shows the results of mean separations for seedling height under different growth media. The seedlings which were raised under T2 (Top soil + goatdung) produced the highest height with mean of 22.15 ± 7.32 , followed by T1 (Top soil + cowdung) with mean of

20.74±6.31, (Top soil + poultry droppings) with mean of 15.96±4.19 and T5 (Top soil only) with mean of 14.67±5.07. Seedlings raised under T4 (Top soil and sawdust) produced the least height with mean of 7.23±1.34. The result further revealed that there was significant difference observed between seedlings raised under different treatments.

Table 6. Seedling height of *Newbouldialaavis* under different organic manure.

Treatment	N	Mean ± SE (cm) $\alpha=0.05$
T1 = Topsoil+Cow dung	144	20.74 ± 6.31b
T2 = Topsoil+Goat dung	144	22.15 ± 7.32a
T3 = Topsoil+Poultry droppings	144	15.96 ± 4.19c
T4 = Topsoil +Sawdust	144	7.23 ± 1.3de
T5 = Control (top soil only)	144	14.67 ± 5.07cd

Collar girth(cm) of *Newbouldialaavis* under different organic manure

Table 7 shows the results of mean separations for seedling collar girth under different growth media. The seedlings which were raised under T1 (Top soil + cowdung) produced the highest collar girth with mean of 1.85±0.76, followed by T2 (Top soil + goatdung) with mean of 1.82±0.78, (Top soil only) with mean of 1.38±0.63 and T3 (Top soil + poultry droppings) with mean of 1.36±0.66. Seedlings raised under T4 (Top soil and sawdust) produced the least collar girth with mean of 0.72±0.36. The result further revealed that there was significant difference observed in the collar girth of seedlings of seedlings raised under different treatments.

Formatted: Indent: First line: 0.39", Line spacing: 1.5 lines

Table 7. Seedling collar girth(cm) of *Newbouldialaavis* under different organic manure

Treatment	N	Mean± SE (cm) alpha=0.05
T1=Topsoil+Cow dung	144	1.85 ± 0.76a
T2=Topsoil+Goat dung	144	1.82 ± 0.78ab
T3=Topsoil+Poultry droppings	144	1.36 ± 0.66bc
T4= Topsoil +Sawdust	144	0.72 ± 0.36abcd
T5 =Control(top soil only)	144	1.38 ± 0.63bd

Formatted: Centered, Line spacing: 1.5 lines

Formatted Table

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Font: 12 pt

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Font: 12 pt

Number of leaves of *Newbouldialaavis* under different organic manure

Table-8 shows the results of mean separations for number of ~~seedling~~ ~~seedlings~~ ~~leaves~~ ~~under~~ ~~leaves~~ ~~under~~ different growth media. The seedlings which were raised under T2 (Top soil + goatdung) produced the highest number of leaves with mean of 21±10.57, followed by T1 (Top soil + cowung) with mean of 1.18±9.55, (Top soil only) with mean of 16±7.91 and T3 (Top soil + poultry droppings) with mean of 14±6.23. Seedlings raised under T4 (Top soil and sawdust) produced the least number of leaves with mean of 7±2.82. The result further revealed that there ~~were~~ ~~was~~ significant difference observed in the number ~~of~~ ~~seedling~~ ~~of~~ ~~seedling~~ leaves raised ~~under~~ ~~different~~ ~~under~~ ~~different~~ ~~treatments~~ ~~-~~ ~~treatments~~.

Formatted: Line spacing: 1.5 lines

Table-8. Seedling number of leaves of *Newbouldialaavis* under different organic manure.

Formatted: Indent: Left: 0", First line: 0", Line spacing: 1.5 lines

Treatment	N	Mean ± SE alpha=0.05
T1=Topsoil+Cow dung	144	18 ± 9.55b
T2=Topsoil+Goat dung	144	21 ± 10.57a
T3=Topsoil+Poultry droppings	144	14 ± 6.23cd
T4= Topsoil +Sawdust	144	7 ± 2.83abc
T5 =Control(top soil only)	144	16 ± 7.91b c

Formatted: Centered, Line spacing: 1.5 lines

Formatted Table

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Font: 12 pt

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Font: 12 pt

Formatted: Font: 12 pt

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Font: 12 pt

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Font: 12 pt


Formatted: Centered, Line spacing: 1.5 lines

Formatted: Font: 12 pt

Effect of Different Treatments on Girth Measurement

The highest mean girth was observed with the cow treatment (1.85 m) while the least was observed with the sawdust treatment (0.72 m). The difference in the mean girth between the different treatments was statistically significant ($P < 0.001$). Figure 2 shows that the difference in the mean girth for the cow treatment and that of the goat treatment were not statistically significant. Both were, however, significantly higher than that of poultry, sawdust, and control. The mean girth for the sawdust treatment was observed to be significantly lower than those of the poultry treatment and the control.

Formatted: Indent: First line: 0.4", Line spacing: 1.5 lines



Formatted: Right: 0.06", Line spacing: 1.5 lines

Figure 2. Effect of the different treatments on girth measurement

Comment [CES12]: Correct the Figures number in all manuscript text.

Effect of Different Treatments on Height

The highest mean height was observed with the goat treatment (22.15 m) while the least was observed with the sawdust treatment (7.23 m). The difference in the mean height between the different treatments was statistically significant ($P < 0.001$). The difference in the mean height for the cow treatment and that of the goat treatment were not statistically significant. Both were, however, significantly higher than those of poultry, sawdust, and control. The mean height for the sawdust treatment was observed to be significantly lower than those of the poultry treatment and the control (Figure 3).

Formatted: Indent: First line: 0.4"





Figure 3. Effect of the different treatments on height

Effect of Different Treatments on the Number of Leaves

The highest mean number of leaves was observed with the goat treatment (21) while the least was observed with the sawdust treatment (7). The difference in the mean number of leaves between the different treatments was statistically significant ($P < 0.001$). The mean number of leaves for the goat treatment was significantly higher than those of poultry, sawdust and control, but not that of the cow treatment. The mean number of leaves for the sawdust treatment was observed to be significantly lower than those of the other treatment and the control (Figure 4).



Formatted: Centered, Right: -0.04", Line spacing: 1.5 lines

Figure 4. Effect of the different treatments on the number of leaves.

B5 (%)	82.10	82.30	82.2000	.03162	.07071	.000	.913
P(mgkg+2)	6.23	6.27	6.2500	.00707	.01581	.000	.913

Formatted: Centered, Line spacing: 1.5 lines

Formatted: Centered, Line spacing: 1.5 lines

DISCUSSION

Appropriate potting media greatly influences germination and early growth rate of *Newbouldialaavis*. Infertility in tropical soils is among the main cause of declines in plants productivity (Dania *et al.*, 2014). The potting media (organic amendment) used in this research significantly improves germination and seedling growth of *Newbouldialaavis*.

Formatted: Indent: Left: -0", First line: 0.4", Right: 0.05", Space After: 0 pt, Line spacing: 1.5 lines

Newbouldialaavis seeds used in this research generally performed very well irrespective of the treatment. This increase in germination percentages support the work of Ehiagbonare and Onyebi (2009) who also recorded increase in germination percentage of *Newbouldialaavis* planted in different media.

Formatted: Font: 12 pt, Highlight

However, T1 had the highest germination percentage of 85%. This supports the finding of Vijayalakshmi *et al.* (2017) who obtained highest germination of *Pterocarpus santalinus* when grown with cow dung. This increase according to Naskaret *al.* (2003) might be because, the addition of cow dung increases the mineral status of soil, enhances the resistance of seed against pest and disease; stimulate germination and other beneficial activities such as sulpho-oxidation and phosphorus solubilization. Composted cow manure makes an excellent growing medium for plants and also contains beneficial bacteria, which convert nutrients into easily accessible forms so they can be slowly released without burning tender plant roots. (Gardening Know How 2021).

The least germination percentage was observed in T5 (control) with 60%, the result of the soil analysis reviewed that the top soil used contains some important chemical nutrient required for plant growth and this was in line with findings of Egunatum *et al.* (2020) that good top soil is considered ideal for planting because it retains nutrients well and also retains water while still allowing excess water to drain away even though it does not contain enough manure for better seedling growth. However, germination period was generally short within 5-9 days in all the treatment and this trend was also reported by Ehiagbonare and Onyebi (2009) who also recorded short germination period of *Newbouldialaavis* planted in different media.

Formatted: Indent: Left: -0", First line: 0.4", Right: 0.05", Space After: 0 pt, Line spacing: 1.5 lines

Formatted: Font: 12 pt, Highlight

The highest mean height was observed in goat dung (T2), this might be due to the fact that goat dung supply organic matter in the soil and have a definite benefit in improving soil nutrient as reported by Duarsa *et al.* (1996) who revealed that goat dung improved water holding capacity, soil capacity, soil fertility, microbial fertility, microbial activities and increased crop dry matter. This also concurs with the report of Murwira *et al.* (1995), who stated that goat manure can help increase pH, organic matter content and cation exchange capacity, and improve soil physical characteristics like water holding capacity that results to increase in plant height growth.

Formatted: Font: 12 pt, Highlight

Cow dung also did well as it was observed to perform better than poultry, sawdust, and control as regards to media effect on the seedling height of *Newbouldia laevis*, this according to Miller (1999). The cow dung media is a good treatment for plant cultivation, which contains moisture 79.05%, phosphorous 0.096% and potassium 0.520%.

Comment [CES13]: Informal language

Formatted: Font: 12 pt, Highlight

The least height growth was observed in sawdust. This low seedling height growth according to Miller and Jones (1995) is due to the fact that sawdust and similar material further depleted the soil nitrogen supply by the production of a compound injurious to the soil nitrifying bacteria and the possibility that such a compound was poisonous to plant roots. This supports the finding of Ashiona *et al.* (2017), who obtained lowest height growth of *Eucalyptus saligna* in Sawdust treatment stating that seedlings in Sawdust treatment had severe stunted growth, symptoms associated with deficiency of essential nutrients in the growth media. This was also in accordance with the findings of Garner (2014), who stated that the slow rate of decomposition of sawdust and temporary depression of nitrogen has been the principal objection to its use in the field.

Formatted: Font: 12 pt, Highlight

The highest mean collar diameter was obtained in T1 (cow dung). Cow dung has long been recognized as perhaps the desirable animal manures because of its high nutrient and organic matter content. This finding corresponds to that of Agbo-Adediran *et al.* (2020), who obtained best result in stem girth of *Entandrophragma angolense* using cow dung. Offiong *et al.* (2010) and Mukhtar (2016) also recorded high performance of stem diameter in cow dung manure application with *Adansonia digitata* and *Tetrapleura tetraptera* seedlings. But this disagrees with the findings of Yahaya *et al.* (2016), whose cow dung treatment on vegetative growth of *Moringa oleifera* obtained a low stem girth diameter result.

Formatted: Font: 12 pt, Highlight

Addition of cow dung increases the organic carbon content of degraded soil which may lead to the increasing activity of beneficial soil microorganisms as well as the fertility status of soil by increasing the availability of nutrients for plants growth; (Akande *et al.*, 2006; Mehedi *et al.*, 2011; Gudugi, 2013; Akande *et al.*, 2006; Mehedi *et al.*, 2011). Cow dung is good for plant growth, it contains moisture 79.05%, potassium 0.520% and phosphorus 0.096% (Miller, 1999).

Formatted: Font: 12 pt, Highlight

Formatted: Font: 12 pt, Highlight

However, there was no significant difference between the effect of treatment with cow dung and goat dung. It was observed that goat dung also did well on the collar diameter more than poultry, sawdust, and control media. This might be to the fact that goat manure has desirable characteristics to increase soil fertility, and this was in accordance with the work of Malavolta *et al.* (2002), that goat manure has a better structure (more solid and much less aqueous) generating better aeration and faster fermentation which makes it ideal for plant growth.

Formatted: Font: 12 pt, Highlight

The least collar diameter was observed in T4 (sawdust). This supports the finding of Midgley (1990) who opined that if decomposed sawdust decreases plant growth, the decrease could be attributed to the wide carbon: nitrogen ratio in the soil. Newton and Daniloff (1997) also stated that the depression of growth and yield following the addition of low nitrogen materials to the soil may be largely overcome by applying 75lbs of nitrate of soda per ton of sawdust used.

Formatted: Font: 12 pt, Highlight

Formatted: Font: 12 pt, Highlight

The highest mean of leaf count was obtained in T2 (goat dung). "Goat manure contains high content of nitrogen as compared to cows, sheep, and poultry manure. This nitrogen further enhances the growth of plants and crops by nitrogen fixation; hence it increases the yield of crops at least by 20 per cent%," said Wangare (2018). Goat manure treatments increased soil N, P, K, Ca, Mg and pH and leaf N, P, K, Ca and Mg growth and yield parameters as number of leaves is significantly increased by goat manure treatments (M.A. Awodun *et al.*, 2007). This was followed by T1 (cow dung), this result is in line with the findings of Alas (2000) who discovered that goat dung kept the soil loose, porous, increased moisture content, microbial activities and provided proper aeration and as a result plant nutrient became available for growth and development of leaves which ultimately increased number of leaves. However, this oppose the findings of Haouvang *et al.* (2017), who found out that cow dung and poultry dung was more effective to number of leaves of *moringa oliefera* than goat dung.

Formatted: Font: 12 pt, Highlight

The least average leaf count was obtained in T4 (sawdust). This may be due to insufficient nitrogen content in the saw dust as reported by Miller and Jones (1995). Turk (1943) found out in greenhouse tests that nitrogen added in sufficient quantity to give the sawdust the equivalent of about 2% by weight of nitrogen would overcome the detrimental effect of the sawdust in soils low in available nitrogen. Sawdust is also a largely sterile material that does not harbor microorganisms due to its low nutrient content (Oekaleboet *al.*, 2002).

Formatted: Font: 12 pt, Highlight

Formatted: Font: 12 pt, Highlight

CONCLUSIONS

The study revealed that growth media has effect on both the germination and early growth of *Newbodialaavis*. The treatment T1 (topsoil and cow dung) has the highest germination percentage of (85%) followed by T2 (82%) however, there was no significant difference in the days to germination for all treatments as their germinations started the same day (the 5th day after sowing of seed). T1 and T3 media recorded the most minimum number of germination period (5 days) followed by T2 (6 days) and T5 (8 days), while T4 had the longest germination period of 9 days.

The T1 (topsoil and cow dung) had the best in germination percentage and collar girth, followed by T2 (topsoil and goat dung) which had the highest seedling height growth and number of leaves. However, the least result was observed in T4 (topsoil and sawdust) both in germination percentage, seedling height, collar diameter and number of leaves.

Therefore, it can be concluded that for maximum germination and optimum growth rate, treatment T1 (topsoil and cow dung) is the best for raising the seedlings of *Newbodialaavis*.

RECOMMENDATION

Based on the results obtained, it is therefore recommended that

- i. for maximum germination and optimum early growth of *Newbodialaavis* cow dung media is recommended;
- ii. where cow dung is not available, goat dung should be used;

iii. sawdust had the lowest germination percentage and poor growth performance; hence it is not recommended for the optimum germination and early growth of *Newbouldialaavis* seedlings:-

iv. Where only germination of *Newbouldialaavis* seedlings are required, ~~any~~ of any of the treatment (T1, ~~T1~~, T2, T3, T5) should be used as they all have germination percentage of 65% and above.

This study is an asset to domestication and regeneration of *Newbouldialaavis* seedlings and can easily be adopted by resource poor farmers and tree growers.

REFERENCES

Agbo-Adediran O. A., Adenuga, D.A., Odeyale, O.C., Musa, F.B. and Agboola, F.O. (2020). *Entandrophragma angolense* seedling. *Journal of Research in Forestry, Wildlife and Environment*, volume 12, no. 3, pp 34-40

Akande, M.O., Oluwatoyinbo, F.I., Kayode, C.O. and Olowokere, F.A. (2006). Response of maize (*Zea mays*) and okra (*Abelmoschus esculentus*) intercrop relayed with cowpea (*Vigna unguiculata*) to different levels of cow dung amended phosphate rock. *World J. Agric. Sci.*, 2: 119.

Aladesanmi, A.J., R. Nia and A. Nahrstedt, (1998). New pyrazole alkaloids from the root bark of *Newbouldialaavis*. *Planta Med.*, 64: 90-9.

Anozie E.L and Oboho E.G (2019); The effects of seed source and different pre-sowing treatment on germination of *Canarium schweinfurthii* Engl. Seeds. *Asian journal of research in Agriculture and forestry* 4(4):1-11.

Anozie E.L, Ibeh K.G, Ndulue N.B, Nwachukwu A.L, Ume C.L (2020): Growth response of *Dennettia. tripetala* G.BAKER to different organic manure at early stage; *European journal of Agriculture and forest research* vol8; 3 PP 17-26.

Arancon, N.Q., A. Pant, T. Radovich, N.V. Hue, J.K. Potter and C.E. Converse, (2012). Seed germination and seedling growth of tomato and lettuce as affected by vermicompost water extracts (teas). *Hort. Sci.*, 47: 1722–1728

Comment [CES14]: Adequate to the references template of AJRAF.

Ashiono, F.A., Kamiri, H.K., and Kinyanjui, M.J. (2017). Effects of saw dust, Forest Soil and Cow Dung Mixtures on growth Characteristics of Blue Gum (*Eucalyptus Saligna*) seedlings in South Kinangop Forest, Nyandarua, Kenya. *Open Journal of Forestry*, 7,373-387

B. Rosie Lerner, "What is Loam?", *Purdue University Consumer Horticulture*, 6 January 2000. Retrieved 5 March 2017

Barwick. M. (2004). *Tropical and Subtropical Trees - A Worldwide Encyclopaedic Guide*.

Burkil. H. M. 1985: *The Useful Plants of West Tropical Africa*. Publisher; Royal Botanic Gardens; Kew.

Burkil. H. M.(1985 - 2004): *The Useful Plants of West Tropical Africa*. PublisherRoyal Botanic Gardens; Kew.ISBN Brief descriptions and details of the uses of over 4,000 plants. A superb, if terse, resource, it is also available electronically on the Web - see <http://www.aluka.org/>.

Dania S.O. Dania, P. Akpansubi, O.O. Eghagara(2014)Comparative effects of different fertilizer sources on the growth and nutrient content of moringa (*Moringa oleifera*) seedling in a greenhouse trial*Adv. Agric.* 2014 (2014) Article ID 726313

Duarsa MAP, Suama I M, Suama IW, Partama IBG Kusumawati, NNC (1996). The effect of goat manure and soil moisture content on tiller number and leaf yield of *Brachiariadecwnbens* over three growth cycles. *Proceedings of 8thAustralian Agronomy Conference, Toowoomba, 1996*.

Ehiagbonare J.E and Onyebi H.I (2009); Seed storage of evaluation, imbibition capacity assessment and seed pre-sowing treatment studies on *Newbouldialaervis* (P.Beauv) ex bureau;scientific research and Essay .vol 4 (5)pp 453-456.

Ersan.Y, Ari.I, Koc.E (2008).Ethanol extracts of *Newbouldialaervis* leaves and stem on serum lipid extract of *Newbouldialaervis* . *Afri.J. Trad. Compl.Alternat. Med.*, 4(4): 476-48

Gafner, S., J.L. Wolfender, M. Nianga and K. Hostettmann, (1997): Phenylpropanoid glycosides from *Newbouldialaervis* roots. *Phytochemistry*, 44: 687-690.

Garner, E (2014). Sawdust as a Mulch and Soil Amendment for Rhododendrons and Azaleas. A. N. Roberts, & A. R. S. Bulletin (Eds.), *Journal American Rhododendron Society (eJournals)*, 5, 58

Germann, R., M. Kaloga, D. Ferreira, J.P. Marais and H. Kolodziej, (2006). Newbouldioside A-C phenylethanoid glycosides from the stembark of *Newbouldialaervis*. *Phytochemistry*, 67: 805-811.

Gudugi, I.A.S. 2013. Effect of cowdung and variety on the growth and yield of okra (*Abelmoschus esculentus* L.). *Eur. J. Exp. Biol.*, 3: 495-498.

Formatted: Font: 12 pt, Portuguese (Brazil), Highlight

Huxley, A., (1992): The New Royal Horticulture Society Dictionary of Gardening. *MacMillan Press, London*, pp: 474-94.

Iwu, (2000). Handbook of African Medicinal Plants. London: CRC Press, Inc; 2000. p.19

Keay *et al.*, (1964) - "Nigerian Trees" Scientific Research Publishing Keay, R.W.J. (1989) Trees of Nigeria. A Revised Version of "Nigerian Trees" (Keay *et al.*, 1964). Clarendon Press, Oxford.

Khan, T. I. ; Dular, A. K. ; Solomon, D. M., (2003). Biodiversity Conservation in the Thar Desert; with Emphasis on Endemic and Medicinal Plants. *The Environmentalist*, 23: 137-144.

L. C Haouvang, Ngakou A, Yemefack M, Mbailao M (2017) Growth response of *Moringaoleifera* Lam. as affected by various amounts of compost under greenhouse conditions. *AOAS* 62:221–226.

Kitajima K, Fenner M.(2000): Ecology of seedling regeneration. In: M Fenner, ed. Seeds: the ecology of regeneration in plant communities. Wallingford, UK: CABI Publishing, 331– 359.

Landis TD, Morgan N (2009). Growing media alternatives for forest and native plant nurseries. In: Dumroese RK, Riley LE (eds) National proceedings: forest and conservation nursery associations – 2008 (tech coords). US Department of Agriculture, Forest Service, Rocky Mountain Research Station, *Proceedings RMRS-P-58*, Fort Collins, pp 26– 31.

Lewis and Manny (2007): Analysis and determination of the qualitative and quantitative phytochemicals present in *Newbouldialaavis* leaves.

M.A. Awodun , L.I. Omonijo and S.O. Ojeniyi , (2007). Effect of Goat Dung and NPK Fertilizer on Soil and Leaf Nutrient Content, Growth and Yield of Pepper. *International Journal of Soil Science*. 142-147

Malavolta, E., Adubos E, Adubações. I. ,d. São Paulo, SP Nobel, 2002. 200 p.

Mehedi, T.A., Siddique, M.A. and Shahid, S.B. (2012). Effect of urea and cow dung on growth and yield of carrot. *J. Bangladesh Agril. Univ.*, 10: 9–13.

Midgley, G. (1990). Creative methodology design. *Systemist* 12, 108-113

Miller, J.H. and Jones, N. (1995). Organic and compost-based growing media for tree seedling nurseries. World Bank Technical Paper No. 264, Forestry Series.

Miller, D. T. (1999). The norm of self-interest. *American Psychologist*, 54,1053–1060. <http://dx.doi.org/10.1037/0003-066X.54.12.1053>

- Mukhtar, R.B. (2016). Response of *Adansonia digitate* seedling to organic manure in Aliero area of Kebbi State. *Nigerian Journal of Agriculture, Food and Environment*, 12(4): 120-122
- Murwira, K.H., Swift, M.J. and Frost, P.G.H. (1995). Manure as a key resource in sustainable agriculture. In: *Livestock and Sustainable Nutrient Cycling in Mixed Farming Systems of sub-Saharan Africa*. Volume II. Technical Papers. Proceedings of an International Conference held in Addis Ababa, Ethiopia 22-26 November 1993. Powell, J.M., Fernandez-Rivera, S., Williams, T.O. and Renard, C. (Eds.), pp. 131-148. ILCA, Addis Ababa
- Naskar, S.K., Sethuraman, P., Ray, R.C. (2003). Sprouting in plant by cow dung slurry. *Validation of Indigenous Technical Knowledge in Agriculture*. New Delhi, India: Division of Agricultural Extension, Indian Council of Agricultural Research; p. 197-201.
- Newton, G.A. and Daniloff, K.B. (1997). The influence of manures and organic residues on plant growth. *Soil Sci.* 24: 95-99.
- Odugbemi (2008): *A Textbook of medicinal plants from Nigeria*. Unilag press, Lagos. Nigeria.
- Offiong, M.O., Udofia, S.I., Owoh, P.W. and Ekpenyong, G.O. (2010): Effects of fertilizer on the early growth of *Tetrapleura tetraptera* Del. *Nigerian Journal of Agriculture, Food and Environment*, 6(2): 53-59.
- Ogunlana and Ramstad, (1975); Ejele et al., (2012). Analysis and determination of the qualitative and quantitative phytochemicals present in *Newbouldialaavis* leaves.
- Okalebo, J. R., Gathua, K. W. & Woomer, P. L. (2002). *Laboratory Methods of Soil and Plant Analysis: A Kenya Working Manual*, KARI, SSSEA, TSBF, SACRED Africa, Moi University. Second Edition, Nairobi, Kenya. pp. 128.
- Peter H. Raven, Ray F. Evert, Susan E. Eichhorn (2005): *Biology of plants - 7th ed.* New York: W.H. Freeman and Company.
- Reddy, D. V., (1998). The effect of supplementation of green forages (*Cenchrus ciliaris*/*Stylosanthes hamata*/*subabul*) on utilisation of rice straw-poultry droppings-rice bran-fish meal diet in buffalo. *Buffalo J.*, 14 (1): 31-44
- Tor-Anyi et al. (2003): traditional medicine practice amongst the igede people of Nigeria. part ii ... Milne-Redh. (Leguminosae-Caesalpinioideae) Leaves Middle Belt, Southwest.
- Turk, L.M. 1943. The effect of sawdust on plant growth. *Michigan Quart. Bull.* 26(1): 10-22.a
- Uphof, J. C. Th. (1959): *Dictionary of Economic Plants*. Publisher; Weinheim. Publisher; Thames & Hudson, London. ISBN 0-500-51181-0.

Usman and Osuji (2007); Usman *et al* (2007): Phytochemical and in Vitro Antimicrobial Assay of the leaf Extract of *Newbouldialaavis*. The African Journal of Traditional, complementary and Alternative Medicines,4, 476-480.

Vijayalakshmi, K.P. and Renganayaki, P.R. 2017. Effect of Pre-Sowing Treatment on Germination of Red Sanders. Int.J.Curr.Microbiol.App.Sci. 6(4): 168-173.

Wangare (2018). Farm using goat manure as mulch and fertiliser doubling its production; A farm in Ndeiya in kikuyu, kiambu.

Wilson, S.B.,and P.J. Stoffella (2006). "Using Compost for Container Production of Ornamental Wetland and Flatwood Species Native to Florida." *Native Plants Journal* 7(3): 293-300.

Yadav, Y. S.; Siddiqui, A. U.; Aruna Parihar, (2005). Management of root-knot nematode *Meloidogyne incognita* infesting gram through oil cakes. J. Phytolog. Res., 18 (2): 263-264

Yahaya U, Oladele O. N., Suleiman R. A., Oloyede E. O. and Anamayi S. E. (2017) Effects of Cow dung and N. P.K Fertilizer at different levels on the Growth performance and Nutrient Composition of *Moringa oleifera* Ann. Exp. Bio., 2016, 4 (1):3-39

Page 21: [1] Formatted	Carlos Eduardo Silveira	11/28/2022 3:43:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Superscript

Page 21: [1] Formatted	Carlos Eduardo Silveira	11/28/2022 3:43:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Superscript

Page 21: [1] Formatted	Carlos Eduardo Silveira	11/28/2022 3:43:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Superscript

Page 21: [1] Formatted	Carlos Eduardo Silveira	11/28/2022 3:43:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Superscript

Page 21: [1] Formatted	Carlos Eduardo Silveira	11/28/2022 3:43:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Superscript

Page 21: [1] Formatted	Carlos Eduardo Silveira	11/28/2022 3:43:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Superscript

Page 21: [2] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [2] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [3] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [3] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [4] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [4] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Font: 12 pt, Bold

Page 21: [5] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [6] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [6] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [7] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [7] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [8] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [8] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [9] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [9] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [10] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
-------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [10] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
-------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [11] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
-------------------------	-------------------------	-----------------------

Page 21: [11] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
-------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [12] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
-------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [12] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
-------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [13] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
-------------------------	-------------------------	-----------------------

Font: 12 pt, Bold

Page 21: [13] Formatted	Carlos Eduardo Silveira	11/28/2022 3:50:00 PM
-------------------------	-------------------------	-----------------------

Font: 12 pt, Bold