

Original Research Article

Medicinal plants traditionally used for the treatment of skin diseases in Southern Punjab, Pakistan

ABSTRACT

Skin conditions are among the most common health problems that affect people. Modern, allopathic, and natural medicine can all help to promote healthy skin. Human skin problems can be successfully treated using medicinal herbs because they contain a wide variety of bioactive molecules. The goal of this research is to look into the dermatological properties of medicinal plants used by different ethnic communities in southern Punjab, Pakistan and it seeks to establish a foundation for the development of novel medications with anti-skin activity. The snowball sampling method was employed for expert sampling in the southern Punjab districts of Multan and Vehari. Direct interviews with traditional herbalists (Hakeem) and shopkeepers selling medicinal plants (pansar) were done between August and December 2021. After gaining their prior informed agreement, 120 experts from southern Punjab were interviewed. 21 medicinal plants were recorded to be utilized in the treatment of skin problems. Plants are largely members of the Asteraceae and Piperaceae families. Leaves (62%) are the plant portion most frequently used as a treatment for skin conditions. Decoction (23%) is the most frequently used method in the study area. *Aloe barbadensis* miller having the highest Relative Frequency Citation (RFC) value while the frequency citation (FC) value ranged from 20 to 60. The highest value of FC was also recorded for *Aloe barbadensis*. The fidelity level (FL) ranged from 66 to 100%. The documented 15 medicinal plants have a higher FL value than 85%.

Key Words: Medicinal plants, Traditional knowledge, Skin disorders, Quantitative analysis, Southern Punjab, Pakistan

1. INTRODUCTION

Humans use plants to meet a variety of survival needs [1]. Plants give variety of food products, fuelwood, timber, raw material for industries and fodder but they are also a good source of herbal medicines for curing a variety of maladies [2]. Indigenous peoples have traditionally employed conventional medical practices [3]. Around 80% of Asian and African communities are considered to use traditional medicine for their healthcare requirements [3, 4]. Numerous ethnic groups rely on natural resources, particularly medicinal herbs. Herbal remedies have been utilized for the treatment, mitigation, prevention, and management of numerous diseases throughout the world since ancient times [5]. The uses of medicinal plant-based remedies are becoming popular day by day [6]. Herbal medication is the treatment method used by the majority of people living in rural regions in which medicinal plants are used as an alternate of allopathic therapy [7]. Allopathic drugs having more efficacy and give quick recovery when treating many diseases but they might have negative effects if administered incorrectly. Herbal medications are preferred due to their low cost, ease of availability, and lack of negative effects [2].

In tropical countries, infectious illnesses are the primary cause of mortality [8]. This is partly due to the decreased efficacy of conventional treatments as a result of the rise of drug-resistant diseases, especially in developing countries [8]. The skin is the biggest organ in the human body and it is highly susceptible to a variety of changes that can cause skin damage in a variety of ways. Additionally, this barrier can be altered by pathogens, which can result in a variety of skin issues such as wounds, boil, acne, atopic dermatitis, shingles, hives, sunburn, contact dermatitis, diaper Rash and general skin illnesses. As a result, skin conditions can seriously degrade health and provide a risk for mental health problems [9]. These health issues faced by peoples of different ages and are estimated to account for 34% of all occupational diseases experienced [10]. Furthermore, dermatological disorders are many and ubiquitous because they are caused by different pathogens and the specific symptoms appeared at the skin. The effects caused by dermatological disorders are significant throughout the world [11]. The Global Burden of Disease (GBD) 2013 report showed that dermatological disorders are the eighteenth main reason of health problems worldwide [12, 13].

Treatment for any sickness, particularly for skin disorders, is necessary to reduce the burden brought on by skin diseases and to protect the healthy body of human beings. There are two medical systems in this regard: traditional medicine and conventional medicine. In the conventional medical system, various medications were already utilized for treating some symptoms associated with skin illnesses [14]. The hefty price tag and occasionally unfavorable side effects of these synthetic medications continue to raise suspicions even though they are successful in treating skin problems. Additionally, several chemical products have issues with allergy and resistance, which leads researchers to search for alternate solutions [15]. Natural medicines made from plants are becoming more and more popular as an alternative to traditional allopathic medications since they are easily accessible, safe, and affordable for the poor. In reality, it has been discovered that using conventional medicines, particularly botanicals, is crucial for treating skin infections [10]. In many nations across the world, plants were used to cure skin problems, greatly enhancing people's access to basic healthcare [16].

The herbal medication in a traditional way to cure different disorders including dermatological disorders is deeply ingrained in Pakistan. However, ethnobotanical investigations on the treatment of skin ailments in southern Punjab, Pakistan remained spatially scattered. Herbal practitioners in the study region are well-versed in the use of therapeutic herbs, but many are hesitant to share their expertise with others. However, this vital information is in danger of being lost by the next generation. Previous ethnomedical research had been conducted in surrounding locations, but this area remained undiscovered [17, 18, 19, 20, 21, 22, 23, 24, 25]. Therefore, this research was organized to investigate the wild plants used to cure skin disorders in the Multan and Vehari regions of Southern Punjab, Pakistan.

2. MATERIALS AND METHODS

2.1. Study area

This investigation was conducted in two main districts (Multan and Vehari) of Southern Punjab, Pakistan. The latitude and longitude of Multan district are approximately 30° 28' 16" North and 71° 43' 54" East while the latitude and longitude of Vehari district are approximately 30° 22' 21" North and 72° 58' 43" East. Multan has the distinction of being one of Pakistan's oldest cities.

Multan is relatively fertile and plain, with the Chenab River running through Tehsil Muzafargarh on the western side. The area has an arid environment with average 175mm rainfall annually and highest temperature 52 degree centigrade was recorded. District Vehari is situated between Nili Bar and the rivers Ravi, Sutlej, and Beas [24].

2.2. Sampling and Ethnobotanical data collection

The snowball sampling method was employed for expert sampling. An open-ended interview was used followed by [26] and [27]. Before it was utilized in the study, the questionnaire was verified by experts, and a pilot study was conducted. Direct interviews with traditional herbalists and pansar were done between August and December 2021. The regional language was used to collect the data. 120 traditional herbalists and pansar across the two main districts of southern Punjab were interviewed.

2.3. Analysis of Quantitative Data using Indices

Use Value (UV)

To calculate the UV of all medicinal plants following formula is used as used by [28].

$$\frac{U.V \sum u}{4} = N$$

Frequency of Citation (F.C) and (R.F.C)

The following formula was used to calculate the RFC as used by [29].

$$R.F.C = \frac{F.C}{N}$$

Fidelity level (FL)

The following formula was used to calculate the FL as used by [30].

$$F.L (\%) = \frac{N.P}{N} \times 100$$

3. RESULTS

3.1. Socio-demographic Information

Table 1 showed that 120 respondents (Hakeem and pansar) participated from the Multan and Khanewal districts of Southern Punjab. Participants age range from 20 to 100 years old, with 23% being young, 50% being middle-aged, and 27% being over 50. Thirty percent of the participants were illiterate, 35% had completed elementary school, 18% had completed optional schooling, 9% had completed higher education, and just 8% had completed college. Half of the respondents are traditional health practitioners (Hakeem) while the remaining half is shopkeepers who have sold medicinal plants (Pansar). Most respondents lived in suburban (55%) and only 45% lived in urban areas of Multan and Khanewal districts.

Table 1. Socio-demographic information of the respondents (n = 120)

Demographic Information	Divisions	Frequency (f)	Percentage (%)
Age	Young age	28	23%
	Middle age	60	50%
	Above 50 y	32	27%
Education	Illiterate	36	30%
	Literate	84	70%
Source of livelihood	Hakeem	60	50%
	Pansar	60	50%
Locality	Sub urban areas	66	55%
	Urban areas	54	45%

3.2. Documented medicinal plant species

21 medicinal plants belong to 18 different families were reported as showed in fig. 1. Table 2 showed the detail enumeration of collected medicinal plants. Plants mostly belong to Asteraceae and Piperaceae families. The plant part used as a remedy for skin diseases is leaves (62%) followed by roots (19%), flowers (18%), seeds (15%) and whole plant (8%) respectively. Decoction (23%) is the most frequently used method in the study area. Fig. 2 showed that most medicinal plants are unknown or herbs.

Different skin diseases were noticed in the study area. The most common skin problems which are treated with traditional remedies are itching (42%) followed by boils (42%), acne (10%), wounds (3%) and inflammation (3%). It has been noticed that most herbal treatments are made up of a mixture of more than one plant species or portion of a plant, as well as other supplements.

Table 2. Detail enumeration of documented medicinal plants related to skin diseases from the study area

Used as remedy for acne disorder				
Vernacular Name/ Scientific Name	Family	Application Method	Disease	Part Used
Aloe vera/ <i>Aloe barbadensis miller</i>	Asphodelaceae	Ointment	Acne	Whole
Barberries/ <i>Berberis vulgaris</i>	Berberidaceae	Decoction	Acne	Fruit
Arq e Ghulab/ <i>Rosa indica</i>	Rosaceae	Ointment	Acne	Flower
Used as remedy for boil disorder				
Kameela/ <i>Mallotus philippensis</i>	Euphorbiaceae	Ointment	Boil	Flower
Satynasi booti/ <i>Argemone Mexicana</i>	Papaveraceae	Ointment	Boil	Leave, Root
Panwar/ <i>Alexandrian senna</i>	Fabaceae	Decoction	Boil	Seed
Charaita/ <i>Swertia chirayita</i>	Gentians	Decoction	Boil	Whole
Gule Mundi/ <i>Sphaeranthus indicus</i>	Asteraceae	Decoction	Boil	Whole
Shreen/ <i>Albizia lebbek</i>	Fabaceae	Ointment	Boils	Seeds
Wild Mint/ <i>Mentha arvensis</i>	Lamiaceae	Decoction	Boils	Whole
Used as remedy for inflammation disorder				
Turmeric/ <i>Curcuma longa</i>	Zingiberaceae	Paste	Inflammation	Roots
Used as remedy for itching disorder				
Kali Zeeri/ <i>Centratherum anthelminticum</i>	Asteraceae	Decoction, Ointment	Itching	Seed
Kali Mirch/ <i>Piper nigrum</i>	Piperaceae	Decoction, Ointment	Itching	Seed
Bapchi/ <i>Psoralea corylifolia</i>	Fabaceae	Decoction	Itching	Whole
Neem/ <i>Azadirachta indica</i>	Meliaceae	Ointment	Itching	Leave
Sandal Surkh/ <i>Santalum paniculatum</i>	Santalaceae	Decoction	Itching	Bark
Anab/ <i>Ziziphus jujube</i>	Rhamnaceae	Decoction	Itching	Fruit

Kalwanji/ <i>Nigella sativa</i>	Ranunculaceae	Decoction	Itching	Seeds
Hena/ <i>Lawsonia inermis</i>	Lythraceae	Paste	Itching	Seeds
Sheesham/ <i>Dalbergia sissoo</i>	Fabaceae	Ointment	Itching	Leaves , Seeds

Used as remedy for wound disorder

Arnica/ <i>Mountain arnica</i>	Asteraceae	Ointment	Wound	Whole
--------------------------------	------------	----------	-------	-------

UNDER PEER REVIEW




Figure 1. Family wise classification





Figure 2. Life form of medicinal plants collected from the study area

3.3. Ethnobotanical indices data

According to fig. 4, the RFC values of several species ranged from 0.1 to 0.24, with Aloe vera having the highest RFC value. Fig. 3 demonstrated that the FC value ranged from 20 to 60, with Aloe vera once more having the highest value. As seen in fig. 5, FL ranged from 66 to 100%. The FL value of 15 medicinal plants is higher than 85%. The skin infections were divided into four groups according to commonly used classifications of skin problems. The skin disorders include itching, pimples, mumps, measles, wounds, boils, skin consumption, abscesses, aggravation,

skin disturbance, irritation, inflammation and acne. It is observed that most skin disorders were treated by using traditional remedies in which wild medicinal plants are used. Most local people visited traditional healthcare centers for the treatment of skin-related issues and belief in



herbal medicines.

Figure 3. Frequency of citation of all collected medicinal plants from the study area

UNDER PUBLICATION




Figure 4. RFCs value of all documented medicinal plants




Figure 5. FL % of all documented medicinal plants

4. DISCUSSION

The skin is the biggest organ in the human body and it is highly susceptible to a variety of changes that can cause skin damage in a variety of ways. Additionally, this barrier can be altered by pathogens, which can result in a variety of skin issues such as wounds, boil, acne, atopic dermatitis, shingles, hives, sunburn, contact dermatitis, diaper Rash and general skin illnesses. As a result, skin conditions can seriously degrade health and provide a risk for mental health problems [9]. These health issues faced by peoples of different ages and are estimated to account for 34% of all occupational diseases experienced [10]. The Global Burden of Disease (GBD) 2013 report showed that dermatological disorders are the eighteenth main reason of health problems worldwide [12, 13].

Medicinal herbs and their extracts are natural treatments that have excellent potential for healing a wide range of ailments, including dermatological problems [31]. Medicinal plants have considerable therapeutic efficacy while having fewer negative effects. They offer a variety of uses in treating dermatological problems, whether as monotherapy or in combination with other treatments [32]. Herbal medication is traditionally used to cure different ailments including dermatological disorders is deeply ingrained in Pakistan. However, ethnobotanical investigations on the treatment of skin ailments in southern Punjab, Pakistan, remained spatially scattered. Herbal practitioners in the study region are well-versed in the use of therapeutic herbs, but many are hesitant to share their expertise with others. However, this vital information is in danger of being lost by the next generation. The purpose of this research study was to investigate the dermatological properties of medicinal plants used by different ethnic communities in southern Punjab, Pakistan and it seeks to establish a foundation for the development of novel medications with anti-skin activity.

Twenty-one (21) medicinal plants belonging to 18 different families were reported which are used for the treatment of skin disorders. Plants mostly belong to Asteraceae and Piperaceae families. The plant part used as a remedy for skin diseases is leaves (62%) followed by roots (19%), flowers (18%), seeds (15%) and whole plant (8%) respectively. These findings are consistent with most ethnobotanical research, which found that when plants are used to cure various skin problems, the leaves are the preferred plant portion [10, 33, 16, 34, 35]. During the survey, three different growth forms were recorded as dominating the research region, and it

was determined that herbs dominate the area by contributing 35%, followed by shrubs (11%), and trees (10%). These findings agreed with those of [36, 37]. Herbs were widely used due to their abundance in the research region. The usage of decoction is favored over other ways of use because it is simpler to prepare, requiring simply the combining of plant parts with water, tea, soup, or milk. It is also because heating plant materials in the form of a decoction accelerate the availability of active phytochemicals [38]. In Pakistan, the usage of plant components in the form of decoction is always favored, and this is confirmed by prior evidence [39, 40, 41].

To determine the popularity and efficacy of the reported plant species used to cure Skin illnesses, UV, RFC, FC, and FL were quantitatively measured. Numerous researchers used the aforementioned quantitative indices to assess the efficacy of the recorded plant species in their investigations [27, 28, 30]. RFC values ranged from 0.1 to 0.24, with *Aloe barbadensis* having the highest RFC value while the FC value ranged from 20 to 60. The highest value of FC was also recorded for *Aloe barbadensis*. It is important to examine ethnomedicinal species with high UVs and RFCs values to determine and show their pharmacological activity [42]. The fidelity level (FL) ranged from 66 to 100%. The documented 15 medicinal plants have higher FL value than 85%. High FL demonstrates the particularity and extensive use of a plant species for a given ailment [43].

5. CONCLUSIONS

21 medicinal plants were recorded to be utilized in the treatment of skin problems. Plants are largely members of the Asteraceae and Piperaceae families. Leaves (62%) are the plant portion most frequently used as a treatment for skin conditions, followed by roots (19%), flowers (18%), seeds (15%), and the entire plant (8%). The most frequently used medicinal plants in the form of decoction (23%). Relative Frequency Citation (RFC) values ranged from 0.1 to 0.24, with *Aloe barbadensis* miller having the highest RFC value while the frequency citation (FC) value ranged from 20 to 60. The highest value of FC was also recorded for *Aloe barbadensis*. The fidelity level (FL) ranged from 66 to 100%. The documented 15 medicinal plants have higher FL value than 85%.

6. REFERENCES

- [1] Munawar T, Anwar K, Bibi Y, & Ahmad F. 2021. "Brighamia insignis" a Hawaiian Endangered Species, Current Status and Future Prospects: A Review: Brighamia insignis" a Hawaiian Endangered Species. Proceedings of the Pakistan Academy of Sciences: B. Life and Environmental Sciences 58(2):17-22.
- [2] Ishtiaq M, Maqbool M, Ajaib M, Ahmed M, Hussain I, Khanam H, Mushtaq W, Hussain T, Azam S, Hayat Bhatti K, Ghani A. 2021. Ethnomedicinal and folklore inventory of wild plants used by rural communities of valley Samahni, District Bhimber Azad Jammu and Kashmir, Pakistan. Plos One 16(1): e0243151.
- [3] Dogara A, Labaran I, Hamad SW, Lema AA, Jakada BH. 2021. Traditional medicinal plants used for the treatment of cancer in Mubi, Adamawa State, Nigeria. Al-Qadisiyah Journal of Pure Science 26(4):258-268.
- [4] Abdul rahman MD, Zakariya AM, Hama HA, Hamad SW, Al-Rawi SS, Bradosty SW, Ibrahim AH. 2022. Ethnopharmacology, biological evaluation, and chemical composition of Ziziphus spina-christi (L.) Desf.: A Review. Advances in Pharmacological and Pharmaceutical Sciences 2022:1-36.
- [5] Ahmad AS. & Sharma R. 2020. Comparative Analysis of Herbal and Allopathic Treatment systems. European Journal of Molecular & Clinical Medicine 7(7):2869-2876.
- [6] Awan AA, Akhtar T, Ahmed MJ, & Murtaza G. 2021. Quantitative ethnobotany of medicinal plants uses in the Jhelum valley, Azad Kashmir, Pakistan. Acta Ecologica Sinica 41(2):88-96.
- [7] Birjees M, Ahmad M, Zafar M, Nawaz S, Jehanzeb S, Ullah F, & Zaman W. 2021. Traditional knowledge of wild medicinal plants used by the inhabitants of Garam Chashma valley, district Chitral, Pakistan. Acta Ecologica Sinica.
- [8] Musa AM, Ibrahim MA, Aliyu AB, Abdullahi MS, Tajuddeen N, Ibrahim H, Oyewale AO. 2015. Chemical composition and antimicrobial activity of hexane leaf extract of Anisopus mannii (Asclepiadaceae). Journal of Intercultural Ethnopharmacology 4(2):129.
- [9] Picardi, A., Lega, I., Tarolla, E., 2013. Suicide risk in skin disorders. Clin. Dermatol. 31, 47–56.
- [10] Abbasi, A.M., Khan, M.A., Ahmad, M., Zafar, M., Jahan, S., Sultana, S., 2010. Ethnopharmacological application of medicinal plants to cure skin diseases and in folk cosmetics among the tribal communities of North-West Frontier Province, Pakistan. J. Ethnopharmacology. 128, 322–335.

- [11] Basra, M.K.A., Shahrukh, M., 2009. Burden of skin diseases. *Expert Rev. Pharmacoecon. Outcomes Res.* 9, 271–283.
- [12] Aldeen, T., Taylor, J., Bates, E., 2020. Assessing comorbidities, quality of life, and the needs of patients with psoriasis and eczema at each consultation; can their needs be assessed and met online? *Clin. Res. Dermatology* 3, 1–8.
- [13] Roderick, H.J., Johns, N.E., Williams, H.C., Bolliger, I.W., Dellavalle, R.P., Margolis, D.J., Marks, R., Naldi, L., Weinstock, M.A., Wulf, S.K., Michaud, C., Murray, C., Naghavi, M., 2014. The global burden of skin disease in 2010: an analysis of the prevalence and impact of skin conditions. *J.I. J. Invest. Dermatol.* 134, 1527–1534.
- [14] Tabassum, N., Hamdani, M., 2014. Plants used to treat skin diseases. *Phcog. Rev.* 8, 52–60.
- [15] Priya, K.S., Gnanamani, A., Radhakrishnan, N., Babu, M., 2002. Healing potential of *Datura alba* on burn wounds in albino rats. *J. Ethnopharmacology.* 83, 193–199.
- [16] De Wet, H., Nciki, S., Vuuren, S.F. van, 2013. Medicinal plants used for the treatment of various skin disorders by a rural community in northern Maputaland, South Africa. *J. Ethnobiology & Ethnomedicine.* 9, 1–10.
- [17] Azhar MF. 2014. Ethnobotanical Potential of Medicinal Shrubs in Socioeconomic Uplift of Cholistan Rangeland Dwellers. Doctoral dissertation, University of Agriculture Faisalabad-Pakistan.
- [18] Azhar MF, Siddiqui MT, Ishaque M, Tanveer A. 2014. Study of ethnobotany and indigenous use of *Calotropis procera* (Ait.) in cholistan desert, Punjab, Pakistan. *Pakistan Journal of Agricultural Research* 52(1):117–126.
- [19] Azhar, MF, Aziz A, Haider MS, Nawaz MF, Zulfikar, MA. 2015. Exploring the ethnobotany of *Haloxylon recurvum* (Khar) and *Haloxylon salicornicum* (Lana) in Cholistan desert, Pakistan. *Pakistan Journal of Agricultural Sciences* 52:1085-1090.
- [20] Azhar, M. F., Aziz, A., Hussain, M., Pirzada, S. A., Ahmad, I., & Rasool, F. 2017. Ethnobotanical studies of *capparis decidua* (forsk.) With special reference to cholistan desert, Pakistan. *J. Agric. Res.* 55(4), 611-618.
- [21] Azhar MF, Aziz A, Siddiqui MT, Zafar S, Abdullah M, Ijaz M, Hussain M. 2018. Evaluation of chemical composition and ethno-botanical uses of *Calligonum polygonoides* L. in Cholistan Desert of Pakistan. *Journal of Medicinal & Spice Plants* 23(3):132-137.

- [22] Azhar, M. F., Naseer, U., Aziz, A., Zafar, S., Qadir, I., Farooq, M., & Anjum, K. 2020. Antioxidant and phytochemical composition of leaves, stem and root extracts of *Withania coagulans* and *Withania somnifera*. *Zeitschrift Arznei-Gewurzpflanzen*, 25, 27-30.
- [23] Abdullah, M., Rafay, M., Farooq Azhar, M., & Yousaf, M. M. 2021. Ecology, Ethnobotany, and Conservation Status of Browse Vegetation from Cholistan Rangelands of Pakistan. *Journal of Rangeland Science*, 11(3), 357-373.
- [24] Hussain, M., Ali, E., Azhar, M. F., Sarfaraz, I. H., Khan, I., Dilbar, D. A., & Imtiaz, M. T. 2022. Phytochemical properties of botanical parts of *eucalyptus camaldulensis* under different edaphic regimes. *Fresenius environmental bulletin*, 31(8), 7788-7795.
- [25] Azhar, M. F., Aziz, A., & Ali, E. 2022. Assessment of medicinal folklores and chemical composition of *Aerva javanica* (Burm. f.) Juss. ex Schult. in Cholistan Desert of Pakistan. *Ethnobotany Research and Applications*, 24, 1-10.
- [26] Abdul rahman M. 2021. Traditional medicinal plants used for the treatment of cancer in Mubi, Adamawa State, Nigeria. *Al-Qadisiyah Journal of Pure Science (QJPS)*. 26(4).
- [27] Kayfi S, Abdulrahman MD. 2021. Ethnopharmacology of Plants in Choman, The Kurdistan Region of Iraq. *Applied Biological Research* 23(4):322-330.
- [28] Mahmoud AD, Abba A. 2021. Ethnomedicinal Survey of Plants used for Management of Inflammatory Diseases in Ringim Local Government, Jigawa State, Nigeria. *Ethnobotany Research and Applications* 22:1-27.
- [29] Abdulrahman MD, Ali AM, Fatihah H, Khandaker MM, Mat N. 2018. Traditional medicinal knowledge of Malays in Terengganu, Peninsular Malaysia. *Malayan Nature Journal*. 70(3):349-364.
- [30] Mahmoud AD, Labaran I, Yunusa A. 2020. Ethnobotany of medicinal plants with antimalarial potential in Northern Nigeria. *Ethnobotany Research and Applications* 19:1-8.
- [31] Moragrega, I., Ríos, J.L., 2021. Medicinal plants in the treatment of depression: evidence from preclinical studies. *Planta Med.* 87, 656–685.
- [32] Li, Y., Huang, J., Lu, J., Ding, Y., Jiang, L., Hu, S., Chen, J., Zeng, Q., 2019. The role and mechanism of Asian medicinal plants in treating skin pigmentary disorders. *J. Ethnopharmacology*. 245, 112173.
- [33] Chaachouay, N., Benkhniq, O., Zidane, L., 2022. Ethnobotanical and ethnomedicinal study of medicinal and aromatic plants used against dermatological diseases by the people of Rif, Morocco. *J. Herb. Med.* 32, 100542.

- [34] Salhi, S., Fadli, M., Zidane, L., Douira, A., 2010. Etudes floristique et ethnobotanique des plantes m´edicinales de la ville de K´enitra (Maroc). *Lazaroa* 31, 133–146.
- [35] Sharma, J., Gairola, S., Sharma, Y.P., Gaur, R.D., 2014. Ethnomedicinal plants used to treat skin diseases by Tharu community of district Udham Singh Nagar, Uttarakhand, India. *J. Ethnopharmacology*. 158, 140–206.
- [36] Shah A, Sarvat R, Shoaib S, Ayodele A, Nadeem M, Qureshi T, Abbas A. 2018. An ethnobotanical survey of medicinal plants used for the treatment of snakebite and scorpion sting among the people of Namal valley, Mianwali district, Punjab, Pakistan. *Applied Ecology and Environmental Research* 16(1):111-143.
- [37] Amjad MS, Zahoor U, Bussmann RW, Altaf M, Gardazi SMH, & Abbasi AM. 2020. Ethnobotanical survey of the medicinal flora of Harighal, Azad Jammu & Kashmir, Pakistan. *Journal of Ethnobiology and Ethnomedicine* 16(1):1- 28.
- [38] Zhang JL, Cui M, He Y, Yu HL, & Guo DA. 2005. Chemical fingerprint and metabolic fingerprint analysis of Danshen injection by HPLC–UV and HPLC–MS methods. *Journal of Pharmaceutical and Biomedical Analysis* 36(5):1029-1035.
- [39] Ahmad M, Sultana S, Fazl-i-Hadi S, Ben Hadda T, Rashid S, Zafar M, Yaseen G. 2014. An Ethnobotanical study of Medicinal Plants in high mountainous region of Chail valley (District Swat-Pakistan). *Journal of Ethnobiology and Ethnomedicine* 10(1):1-18.
- [40] Bibi T. Ahmad M, Tareen RB, Tareen NM, Jabeen R, Rehman S.U., Yaseen G. 2014. Ethnobotany of medicinal plants in district Mastung of Balochistan province-Pakistan. *Journal of Ethnopharmacology* 157:79-89.
- [41] Tariq A, Sadia S, Fan Y, Ali S, Amber R, Mussarat S, Adnan M. 2020. Herbal medicines used to treat diabetes in Southern regions of Pakistan and their pharmacological evidence. *Journal of Herbal Medicine* 21: 100323.
- [42] Yaseen G. 2019. Ethnobotany and Floral Diversity of Medicinal Plants in Deserts of Sindh-Pakistan. Quaid-i-Azam University, Islamabad.
- [43] Shil S, Choudhury MD, Das S. 2014. Indigenous knowledge of medicinal plants used by the Reang tribe of Tripura state of India. *Journal of Ethnopharmacology* 152:135-141.