

Effect of Salicylic Acid and Gibberelic Acid on germination and growth of Bitter Gourd, *Momordica Charantia* L.

Formatted: Centered

ABSTRACT

The current study looked into the effects of different salicylic acid (SA) and gibberelic acid (GA₃) concentrations on the bitter gourd's ability to fast germination and rapid growth. The experiment's findings indicated that lower concentrations of SA and GA₃ had a beneficial effect, however greater levels significantly reduced the bitter gourd's ability to germinate and develop in comparison to the control. Findings of the study showed priming concentrations of 3mM SA, 6mM SA, and 1mM GA₃, the germination percentage, shoot and root length, seedling vigor, and fresh and dry weight of the shoot and root, RWC all markedly improved.

Comment [DS21]: not clear rewrite

Key words: Salicylic acid, Gibberelic acid, Germination, Growth, Bitter gourd

INTRODUCTION

A balanced diet for humans depends heavily on vegetables, which also provide farmers with a source of revenue. *Momordica charantia*, a cucurbitaceous vegetable known for its bitter taste and therapeutic properties, is one of the most consumed cucurbitaceous vegetables in Bangladesh. It is widely grown around the nation, primarily in the summer. Its immature fruit, which also functions as a blood purifier and is very helpful to diabetics, is a great source of nutritional fiber, minerals, and vitamins (C and A) [3]. It also possesses anti-carcinogenic qualities and can be employed as a cytostatic drug against several cancer types [13]. Additionally, it is employed in conventional therapy to treat conditions like perlipidemia, digestive issues, menstrual irregularities, and a number of microbiological diseases [13].

Comment [DS22]: Re-numbering references from smallest to largest

An essential component of production, quality, and ultimately the profitability of vegetable farmers is uniform and quick germination. Even though bitter gourd seeds have a high germinability, rapid and uniform emergence is usually problematic because of the thick seed coat that causes the seed to slowly consume water and produce delayed germination [4]. Priming is one of the strategies that have been used to address this issue. Increasing crop performance through seed priming is an easy, inexpensive and efficient technique. It is referred to as a physiological technique that hydrates and drydries seeds without radicle protrusion into water or the solution of additional priming chemicals to enhance the pre-germinative metabolic process [24, 29, 34]. It has been demonstrated that several horticultural and agricultural croperops profit from it in terms of seed germination, seedling establishment and ultimately productivity [6, 10, 11, 12, 28]. Thus, it has been suggested for a long time as a possible means of enhancing crop performance [7]. Additionally, primed seeds exhibit improved uniformity in the emergence of seedlings and a higher rate of germination, both of which aid in the regular establishment of crops and subsequently the production. Better plant water status management and an improved ability for nutrient uptake are linked to the primed plants' rapid development [2, 18].

Plant hormones, which have tiny molecules and are present in minuscule amounts in cells, are essential for controlling the growth processes of plants. Salicylic acid (SA; o-hydroxy benzoic acid) is one of the crucial plant growth regulators (PGRs) that contributes significantly to the development and growth of plants. For instance, SA priming of cereal seeds has been shown to positively affect plant growth and development in maize [25] and also in rice [5, 9, 15, 23, 30]. Therefore, SA has the potential to enhance yield components as well as biological, physiological and morphological indices in plants [2]. Gibberellins (GA), another significant phytohormone that is a member of the tetracyclic diterpenoid carboxylic acid group, can stimulate plant growth and development by encouraging seed germination and releasing dormancy [8, 10]. They have an impact on growth and development throughout the entire plant life cycle as tiny plant growth molecules. Therefore, the goal of this study is to investigate the ideal SA and GA₃ concentration for boosting bitter gourd seed germination uniformity and seedling growth.

MATERIALS AND METHODS

The experiment was accomplished using local bitter gourd variety Dhakaiya at the Department of Seed Science and Technology in Bangladesh Agricultural University. Initially, uniform-sized seeds were sterilized for 5 minutes with 1% sodium hypochlorite, then washed repeatedly with double-distilled water. Firstly, for priming, the seeds were soaked in 1mM, 2 mM and 3mM GA₃, and 3mM, 6mM and 9mM SA in separate screw-capped bottles. For control, untreated seeds were used. Ten bitter gourd seeds were then placed on petri plates (150*20 diameter) with three layers of Whatman filter papers moistened with 20 ml distilled water and stored in standard laboratory conditions (room temperature was 25±1⁰C and relative humidity was 95%). Three independent duplicates of the experiment were run in a completely randomized block design. List 1: The following treatments were maintained:

Comment [DS23]: Petri dishes

Treatment	Concentration of priming agent
T1	Control
T2	3mM SA
T3	6mM SA
T4	9mM SA
T5	1mM GA ₃
T6	2mM GA ₃
T7	3mM GA ₃

Determination of germination and growth parameters

Germination percentage, vigor index, root length, shoot length, fresh and dry weight and relative water content of leaf were considered as germination and growth parameters and determined in control as well as plants under treatment.

Germination percentage

The number of sprouted seeds were counted daily commencing from [the](#) 1st day to 7th day. After 7th day, final count was done and Germination Percentage (GP) of final day was calculated by the following formula stated in [22]:

Percent germination (PG) = (Total no. of seeds germinated)/ (Total no. of seeds taken for germination) x100

Shoot and root length

Shoot and root length of all sprouting's from each replication were measured on [the](#) 15th day. Shoot length was measured from shoot base to the tip of the longest leaf and root length was measured from root base to the root tip.

Fresh and dry weight of shoot and root

Fresh weight (g) of root and shoot were measured just after harvesting. The dry weight (g) of root and shoot were determined by drying the sample in an oven at 80±2°C for three days' till [attainingattained](#) a constant weight.

Relative water content (RWC)

The procedures of Mostofa and Fujita were used to determine the relative water content (RWC) [17]. After 14 days of planting, leaf samples were collected, and fresh weights (FW) of the leaves were taken. After the leaves had been submerged in water for one or two hours, their turgid weights were measured. Following the removal of excess water from the turgid leaves, the turgid weight (TW) was immediately recorded. To determine the leaves' dry weight, they were subsequently oven-dried for 48 hours at 70 degrees Celsius (DW). The RWC was examined using the following equation:

$$\text{RWC (\%)} = (\text{FW} - \text{DW}) / (\text{TW} - \text{DW}) \times 100$$

Statistical analysis

Utilizing the statistical program Minitab 17.0, a one-way ANOVA was performed on the data gathered for each parameter (Minitab Inc., State College, PA, USA). Pair-wise comparisons revealed statistical differences between the means of various treatments (P 0.05).

RESULTS

Effect of priming on germination and seedling growth of bitter gourd

Germination percentage

The effects of SA priming and GA₃ priming on germination of bitter gourd seedlings are presented in Table 1. Seed priming with SA and GA₃ increased germination percentage considerably. Priming with 3 and 6 mM SA showed 100% germination and with 9mM 87.51%. Also priming with 1 and 2 mM SA showed 100% germination and with 3mM 87.5%. All the concentrations showed increased germination percentage compared with control which only

accounted for 62.33%. Germination percentage improved between 40-60% with both priming agents compared with control.

Seedling growth and vigour index

To evaluate the consequences of SA and GA₃ priming on the growth of bitter gourd seedlings and to find out the effect of these agents on seedling vigor, the shoot and root length was monitored (Table 1).

Priming concentrations had both positive and negative ~~effectseffect~~ on shoot and root length of bitter gourd seedlings. The priming with 1 mM GA₃ showed the highest shoot length and the shoot length was recorded as 24.3cm. Again, priming with 2 and 3mM GA₃ increased the shoot length significantly by 40% and 7%, respectively, in comparison with control plants. Moreover, seed priming with 3 and 6 mM SA enhanced shoot length by 14% and 13% but priming concentration 9mM SA decreased shoot length by 17% compared to control. Likewise, root length showed significant variation with different priming treatments. Priming with 6 mM SA had the highest root length which accounted for 16.35cm and priming with 3 mM SA had the second highest root length which was 15.5cm. All other treatments had decreased value compared with control. Priming with every agent, significantly increased SVI both for SA and GA₃ primed conditions compared with control. Among the treatments, the highest SVI was recorded for priming with 1mM GA₃ and the lowest result was found for 9 mM SA.

Table 1: Effect of concentrations of plant growth regulators i.e salicylic acid and gibberelic acid on germination and seedling properties of bitter gourd

Formatted: Font: Bold

Treatment	Germination Percentage (%)	Shoot Length (cm)	Root Length (cm)	Vigour Index
T1 (Control)	62.33c	15.26667e	13.44c	1789.418
T2 (3mMSA)	100a	17.5c	15.5b	3300c
T3 (6mMSA)	100a	17.3c	16.35333a	3365.333c
T4 (9mMSA)	87.50	12.57333f	10.53f	2021.726e
T5 (1mM GA3)	100a	24.3a	12.3d	4660a
T6 (2mM GA3)	100a	21.44b	11.29667e	4273.667b
T7 (3mM GA3)	87.5b	16.4d	9.27g	2858.633d
Level of Significance	*	*	*	*

In a column, values having similar letter(s) do not differ significantly at 5% level of probability by LSD0.05 test. * indicates significant at 5% level of significance

* indicates significant at 5% level of significance

Fresh and dry weight

A considerable variation in fresh and dry weight was observed as a result of priming of bitter gourd seeds (Fig.). The highest shoot and root fresh weight was observed with priming

concentration of 6mM SA which was 2.45g and 0.49g, respectively. Priming with 3mM SA and 1 mM GA₃ increased shoot fresh weight by 18% and 8%, respectively where a decrease in fresh weight of shoot was observed for other treatment compared with control. Similarly priming with 3 and 6mM SA increased root fresh weight by 63% and 86%, and priming with 1mM GA₃ increased by 31% compared with control.

In case of dry weight the highest shoot and root weight was observed with priming concentration of 6mM SA₂ which was 0.29g and 0.13g, respectively. Priming with 3mM SA and 1 mM GA₃ increased dry weight of shoot by 22% and 9%, respectively where a decrease in shoot dry weight was observed for other treatment compared with control. Moreover root dry weight showed an similar pattern as of increase 84% and 5% with 3mM SA and 1 mM GA₃ priming compared with control. Other concentrations of priming decreased in root dry weight compared with control.




Figure 1. Effect of concentrations of plant growth regulators i.e salicylic acid and gibberelic acid on (a) shoot fresh weigh (b) root fresh weight, (c) shoot dry weight, (d) root dry weight of bitter gourd. Vertical bars are SEM (n=3).

Formatted: Font: Bold

Relative water content (RWC)

A considerable variation of relative water content was observed as a result of priming seeds (Fig. 2). All the concentrations except 9mM SA relative water content increased between 2-11%

whereas in 9mM SA RWC decreased by 0.9% compared to control. The highest RWC was observed in pretreatment with 1mM GA₃.




Figure 2. Effect of concentrations of plant growth regulators i.e salicylic acid and gibberelic acid on relative water content of bitter gourd. Vertical bars are SEM (n=3).

Formatted: Font: Bold

DISCUSSION

The process of seed germination is the first and most critical stage in the growth of a plant because it helps seedlings adapt to their constantly changing environment and increase their production [24, 26, 31]. Due to this feature, several pre-treatment application procedures are frequently utilized to promote seedling germination and establishment. Because the bitter gourd's seed coat is so thick, it takes longer for the seed to soften before germination, which causes a delayed emergence [4]. By assisting the biological processes necessary for germination and weakening the seed coat before to sowing, phytohormone pre-sowing promotes early germination [20]. Early seed germination and emergence cause seedlings to grow quickly and become taller. According to the results of our study, 3mM, 6mM, and 1 mM SA and GA₃ increased germination %, seedling vigor, shoot and root length, fresh weight and dry weight of seedlings, and RWC which positively supports the above statements. On the other hand, bitter gourd germination and growth were negatively impacted by greater SA and GA₃ concentrations. These findings are in line with several studies that have demonstrated the effectiveness of low quantities of signaling molecules in promoting germination and growth [1, 2, 14, 21, 29, 33]. Plants grown from primed seeds grow more quickly than those grown from unprimed plants, resulting in taller seedlings (Fig 3). Similar to this, [19, 27, 16] reported that cucurbits had longer shoots when they were sown after priming than when they were not. The positive impact of priming on plant growth can be attributed to better root development and, as a result, an

Comment [DS24]: rewrite this sentence

increased capacity for utilizing nutrients that permits a higher relative growth rate [18] and better plant water status regulation [2]. The direct impact of pretreatment on the regulation of the cell cycle and the mechanisms of cell elongation can also be the reason.

CONCLUSION

It has been determined that the seed priming concentration has a substantial positive and negative impact on the bitter gourd's quick germination and seedling growth. The germination percentage, shoot and root length, seedling vigor, and fresh and dry weight of the shoot and root all significantly increased with priming concentrations of 3mM SA, 6mM SA, and 1mM GA₃. So it stands to reason that these concentrations might be utilized to ensure consistency in the effective manufacturing of bitter gourd. To confirm our findings, it is strongly advised to carry out the same experiment on a larger scale.

REFERENCES

1. Ahmad I, Basra SMA, Wahid A. Exogenous application of ascorbic acid, salicylic acid and hydrogen peroxide improves the productivity of hybrid maize under at low temperature stress. *Int J Agric Biol.* 2014;16:825–830.
2. Ahmed M, Qadeer U, Ahmed ZI, Hassan FU. Improvement of wheat (*Triticum aestivum*) drought tolerance by seed priming with silicon. *Arch Agron Soil Sci.* 2016; 62(3):299–315.
3. Asna AC, Joseph J, Joseph John K. Botanical Description of Bitter Gourd BT—The Bitter Gourd Genome. In: Kole C, Matsumura H, Behera TK, editors. Cham: Springer International Publishing; 2020. p. 7–31. https://doi.org/10.1007/978-3-030-15062-4_2
4. Baig KK, Ara N, Ali S, Khan BP, Wahab A, Rabbani U. Effect of seed priming on bitter gourd with different sources of phosphorus at various soaking durations. *Pure Appl Biol.* 2020; 9(1):80–90.
5. Basra SMA, Farooq M, Tabassam R, Ahmad N. Physiological and biochemical aspects of pre-sowing seed treatments in fine rice (*Oryza sativa* L.). *Seed Sci Technol.* 2005; 33(3):623–8.
6. Bruce TJA, Matthes MC, Napier JA, Pickett JA. Stressful “memories” of plants: Evidence and possible mechanisms. *Plant Sci.* 2007; 173(6):603–8.
7. Chen K, Arora R. Priming memory invokes seed stress-tolerance. *Environ Exp Bot* [Internet]. 2013; 94:33–45.
8. Farooq M, Basra SM, Wahid A, Ahmad N. Changes in Nutrient-Homeostasis and Reserves Metabolism During Rice Seed Priming: Consequences for Seedling Emergence and Growth. *Agric Sci China* [Internet]. 2010; 9(2):191–8.
9. Farooq M, Basra SMA, Ahmad N. Improving the performance of transplanted rice by seed priming. *Plant Growth Regul.* 2007; 51(2):129–37.

Comment [DS25]: Follow pattern of journal. Correct the all references according to journal pattern. Check all references are traceable to put on the DOI at the end of each reference

10. Farooq M, Basra SMA, Rehman H, Saleem BA. Seed priming enhances the performance of late sown wheat (*Triticum aestivum* L.) by improving chilling tolerance. *J Agron Crop Sci.* 2008; 194(1):55–60.
11. Farooq M, Usman M, Nadeem F, Rehman HU, Wahid A, Basra SMA, et al. Seed priming in field crops: Potential benefits, adoption and challenges. *Crop Pasture Sci.* 2019; 70(9):731–71.
12. Gopalan C, Sastri B. R, Balasubramanian SC. Nutritive Value of Indian Foods. National Institute of Nutrition, Indian Council of Medical Research (ICMR); 1989.
13. Grover JK, Yadav SP. Pharmacological actions and potential uses of *Momordica charantia*: A review. *J Ethnopharmacol.* 2004; 93(1):123–32. <https://doi.org/10.1016/j.jep.2004.03.035> PMID: 15182917
14. Jamil E, Zeb S, Ali QS, Ahmad N, Sajid M, Siddique S, et al. Effect of Seed Soaking On Seed Germination and Growth of Bitter Gourd Cultivars. *Pure Appl Biol.* 2016; 5(1):31–6.
15. Mahboob W, Rehman HU, Basra SMA, Afzal I, Abbas MA, Naeem M, Abbas M. Seed priming improves the performance of late sown spring maize (*Zea mays*) through better crop stand and physiological attributes. *Int J Agric Biol.* 2015;17(3):491–498. doi: 10.17957/IJAB/17.3.14.283. [[CrossRef](#)] [[Google Scholar](#)]
16. Mehta DK, Kanwar HS, Thakur AK, Thakur S, Thakur KS. Standardization of seed hydro-priming duration in bitter gourd, *Momordica charantia* l. *Int J Bio-resource Stress Manag.* 2014; 5(1):98.
17. Mostofa, M.G., Fujita, M. Salicylic acid alleviates copper toxicity in rice (*Oryza sativa* L.) seedlings by up-regulating antioxidative and glyoxalase systems. *Ecotoxicology* 22, 959–973 (2013). <https://doi.org/10.1007/s10646-013-1073-x>
18. Muhammad I, Kolla M, Volker R, Gunter N. Impact of Nutrient Seed Priming on Germination, Seedling Development, Nutritional Status and Grain Yield of Maize. *J Plant Nutr.* 2015; 38(12):1803–21.
19. Ogbuehi HC, Madukwe DK, Ashilonu P. Assessment of Hydro Priming of Seeds on Performance of Morphological Indices of Bambara Groundnut (*Vigna Subterrenea* Linn.) Landrace. *Glob J Biol Agric Heal Sci.* 2013; 2(2):17–22.
20. Parida AK, Das AB. Salt tolerance and salinity effects on plants: a review. *Ecotoxicol Environ Saf.* 2005;60(3):324–349. doi: 10.1016/j.ecoenv.2004.06.010.
21. Pill, W.G. and E. A. Kilian. 2000. Germination and emergence of parsley in response to osmotic or matricseed priming and treatment with gibberellin. *HortSci*, 35(5):907-909.
22. Rauf F, Sagar A, Rahman T, Hossain MA, Kabir H, Hossain AK. Seed Quality Status of Rice Varieties Based on Physical Properties, Seed Health and Proximate Composition. *Int. J. Plant Soil Sci.* 2020;32:14-24.
23. Rehman H, Iqbal H, Basra SMA, Afzal I, Farooq M, Wakeel A, Ning W. Seed priming improves early seedling vigor, growth and productivity of spring maize. *J Integr Agric.*

- 2015;14(9):1745–1754. doi: 10.1016/S2095-3119(14)61000-5. [[CrossRef](#)] [[Google Scholar](#)]
24. Rhaman MS, Rauf F, Tania SS, Khatun M. Seed priming methods: Application in field crops and future perspectives. *Asian Journal of Research in Crop Science*. 2020a;25:8-19.
 25. Ruan S, Xue Q, Tylkowska K. The influence of priming on germination of rice (*Oryza sativa* L.) seeds and seedling emergence and performance in flooded soil. *Seed Sci Technol*. 2002;30:61–67. [[Google Scholar](#)]
 26. Sadeghian SY, Yavari N. Effect of water-deficit stress on germination and early seedling growth in sugar beet. *J Agron Crop Sci*. 2004; 190(2):138–44.
 27. Shakuntala NM, Kavya KP, Macha SI, Kurnalliker V, Patil M. Studies on standardization of water soaking duration on seed quality in cucumber (*Cucumis sativus* L.) seeds. 2020; 9(4):1400–4.
 28. Sher A, Sarwar T, Nawaz A, Ijaz M, Sattar A, Ahmad S. Methods of seed priming. In: *Priming and Pretreatment of Seeds and Seedlings*. Springer; 2019. p. 1–10.
 29. Sung Y, Cantliffe DJ, Nagata RT, Nascimento WM. Structural changes in lettuce seed during germination at high temperature altered by genotype, seed maturation temperature, and seed priming. *J Am Soc Hortic Sci*. 2008; 133(2):300–11.
 30. Szalai G, Pál M, Árendás T, Janda T. Priming seed with salicylic acid increases grain yield and modifies polyamine levels in maize. *Cer Res Commun*. 2016;44:537–548. doi: 10.1556/0806.44.2016.038. [[CrossRef](#)] [[Google Scholar](#)]
 31. Tania SS, Rhaman MS, Hossain MM. Hydro-priming and halo-priming improve seed germination, yield and yield contributing characters of okra (*Abelmoschus esculentus* L.). *Tropical Plant Research*. 2020;7:86-93.
 32. Waqas M, Korres NE, Khan MD, Nizami AS, Deeba F, Ali I, et al. Advances in the concept and methods of seed priming. In: *Priming and pretreatment of seeds and seedlings*. Springer, Singapore; 2019. p. 11–41.
 33. Yarnia, M. and E. F. M. Tabrizi. 2012. Effect of Seed Priming with Different Concentration of GA3, IAA and Kinetin on Azarshahr Onion Germination and Seedling Growth. *J. Basic. Appl. Sci. Res.*, 2(3):2657-2661.
 34. Yibchok-Anun S, Adisakwattana S, Yao CY, Sangvanich P, Roengsumran S, Hsu WH. Slow acting protein extract from fruit pulp of *Momordica charantia* with insulin secretagogue and insulinomimetic activities. *Biol Pharm Bull*. 2006; 29(6):1126–31.