

## **Growth Response of *Pterocarpus mildbraedii* (Harms) Seedlings Inoculated with Arbuscular Mycorrhizal Fungi (AMF) under Salt Stress in Calabar, Nigeria.**

### **ABSTRACT**

**Aim:** To determine the influence of four AMF strains on dry biomass of root, stem and leaf of *P. mildbraedii* under salt stress.

**Place and duration of study:** Calabar in the south-south zone of Nigeria. A study was conducted between February and May, 2021.

**Methodology:** The experiment was a  $9 \times 3$  randomised complete block design comprising of nine treatments with three replicates, totalling 27 pots. The *P. mildbraedii* seedlings were inoculated with four AMF strains (*Glomus intraradices*, *Glomus etunicatum*, *Glomus mossae*, and *Glomus occultum*) and were subjected to a 75 mM level of sodium chloride (NaCl) salt stress. AMF colonization and dry biomass were determined at harvest (12 weeks after emergence).

**Results:** The results clearly indicated that all the salt stressed seedlings had higher AMF root colonization than their unstressed counterparts. Seedlings treated with *G. intraradices* (*Gi*) in combination with NaCl (*GiNaCl*) recorded the highest root colonization of AMF of 96.24% and 66.77% as the lowest in seedlings inoculated with *G. intraradices*. The total dry weight of the inoculated seedlings were significantly ( $P \leq 0.05$ ) higher than the control even under salt stress. However, inoculation with *G. intraradices* recorded the highest total dry weight of 16.03g/plant followed by *G. mossae* with 14.00g/plant.

**Conclusion:** AMF-inoculated plants can successfully cope with different ecological extremes (salinity) and subsequently help improve afforestation in coastal areas of Nigeria. Improvement in dry biomass accumulation is an essential mechanism of salinity tolerance in mycorrhizal plants.

**Keyword:** Arbuscular mycorrhizal fungi, *Pterocarpus mildbraedii*, salt stress, seedlings inoculated, growth response.

### **1. INTRODUCTION**

Coastal areas are important land reserve resources, gaining global interest for their vegetation, conservation, and rehabilitation (Rodrigues *et al.*, 2011). The Calabar coastal area lies in the tropical high rainforest belt of Nigeria with an annual rainfall of 2500 – 3500mm (Iwena, 2008). However, vegetation restoration of local coastal areas has not been effectively accomplished due to shallow water table, regular stormy weather, and high salt accumulation. Salt stress in this area is the underlying and severe limitation that affects the conservation and restoration of vegetation (Osim, 2021a). High salt concentration in plants above certain threshold levels could lead to oxidative damage and ion injury to plants causing metabolic disorder, lessen activities of biocatalysts, suppression of the photosynthetic activity of leaves and disintegration of cell membrane integrity, consequently, resulting in limited plant growth and development. (Kumaret *et al.*, 2015).

Arbuscular mycorrhizal fungi (AMF) are obligate biotrophs that institute an endosymbiotic association with the roots of vascular plants (Choiet *et al.*, 2018). AMF have favorable effects on photosynthetic

activity, decrease disease invasion, improve mineral nutrient uptake in the host plant and confer tolerance to abiotic stresses caused by salinity stress, heavy metals and abrupt temperature changes (Baltazar-Bernalet *et al.*, 2022). The favorable effects of AMF on host plants are the production of polyamines and antioxidants, osmotic adjustment, help with water transport in plants exposed to salinity stress, and maintenance of an ionic balance in the  $\text{Na}^+/\text{K}^+$  ratio through the external mycelium to improve plant growth, survival, and development under salt stress (Salwanet *et al.*, 2019; Thangavelet *et al.*, 2022). However, there is paucity of information on how AMF affect growth and the salt tolerance of indigenous trees species in south-south coastal areas of Nigeria.

There are thirty-five (35) species of *Pterocarpus* that are presently accepted (World Agro. Forestry Centre, 2008). The species are widely distributed, extending across the African continent, but *Pterocarpus mildbraedii* (Harms) is a distinct sub-species confined to Eastern Tanzania (Babaretet *et al.*, 2012). It is an evergreen or semi-deciduous tree belonging to the family *Fabaceae* and found in the lowland, dry evergreen and riverine forest. As an important economic indigenous tree species in Cross River State, Nigeria, *P. mildbraedii* yield valuable timber traded as paduak (Osim, 2021b). The paduak reddish wood is valued for its toughness, stability in use and decorativeness. *P. mildbraedii* is well adapted to grow in local, moderately saline soil and is a likely candidate for coastal areas afforestation project (Lammenset *et al.*, 1995). However, despite its prolific seed production and wide range of uses, *P. mildbraedii* seedlings have low survival rates in the soil of local coastal areas due to high soil salinity, which limits their use for the afforestation of this area (Osim, 2021b). Therefore, the present study was conducted based on the hypothesis that AMF could promote the growth, biomass and salt tolerance of *P. mildbraedii* seedlings. A pot experiment was performed in a field to determine the influence of four AMF strains (*Glomus intraradices*, *Glomus etunicatum*, *Glomus mossae* and *Glomus occultum*) on dry biomass of root, stem and leaf of *P. mildbraedii* under salt stress (75mM NaCl).

## 2. MATERIALS AND METHODS

### 2.1 Study Area

The field work was conducted from February 2021 to May 2021 and lasted for 12 weeks in the University of Calabar, Cross River state, Nigeria. Calabar was in the tropical high rainforest agro-ecology of the equatorial climatic belt of Nigeria (Latitude  $5^{\circ}00'$  and  $5^{\circ}40'N$ , Longitude  $8^{\circ}40'$  and  $8^{\circ}62'E$ ) and is about 70m above sea level (Iwena, 2008). It has a bimodal annual rainfall distribution that ranges from 2500-3500mm with a mean annual temperature range of  $22.2^{\circ}C$  to  $38.2^{\circ}C$  and a relatively high humidity range of 75-90%.

### 2.2 Plant Seedlings, AMF Inocula and Soil

The seeds of *P. mildbraedii* were obtained from ministry of Agriculture Department of Forestry, Calabar in January 2019 and were subjected to pre-treatment according to the methods adopted by ILDIS (2005). Four AMF strains (*Glomus intraradices*, *Glomus etunicatum*, *Glomus mossae* and *Glomus occultum*) of inocula consisting of spores, mycelium and infected root fragments were provided by the International Institute for Tropical Agriculture, Ibadan. Soil sample was collected at 0-20 cm depth and was sterilized by heating using the hot air oven at  $160^{\circ}C$  for 2 hours and used as the growth medium.

### 2.3 Inoculation, Treatments and Experimental Design

The pots (39cm diameter and 49cm deep) were each filled with 9kg sterilized soil and placed on racks under natural field conditions. These were watered to field capacity and left to drain overnight. Fifty grams (50g) of crude inoculum was placed 3cm below the surface of the soil in AMF designated pots before sowing to produce mycorrhizal plants (Rabie and Almadini, 2005). Seedlings of uniform height were selected after thinning at one week after emergence (WAE). The treatments consisted of uninoculated and without NaCl (control), *G. etunicatum* (*Ge*), *G. etunicatum* with NaCl (*GeNaCl*), *G. intraradices* (*Gi*), *G. intraradices* with NaCl (*GiNaCl*), *G. mossae* (*Gm*), *G. mossae* with NaCl (*GmNaCl*), *G. occultum* (*Go*), and *G. occultum* with NaCl (*GoNaCl*). Differential salinization commenced at two weeks after emergence of seedlings to avoid subjecting plants to osmotic shock (Abbaspour, Fallahyan and Fahimi, 2005). The salt seedlings were watered with 50 ml of 75 mM NaCl solution per pot, once a week. The experiment was a 9x3 randomized complete block design which comprised of nine (9) treatments with three (3) replicates totaling 27 pots.

## 2.4 Growth Measurements

Twelve weeks after emergence (12WAE), the seedlings were carefully uprooted after thoroughly watering the soil to loosen it. Each seedling was separated into roots, stems, and leaves and were placed in a separately labeled paper envelope and then oven dried at 70<sup>0</sup>C to constant dry weight (Wanget *al.*, 2019). The total plant dry weight was computed as the sum of the dry weight of root, stem and leaf. The dry weight data obtained were then used to calculate the root: shoot ratio. The root: shoot ratio (R/S) of each seedling was calculated as a ratio of root dry weight to shoot dry weight;

$$\text{Root: Shoot Ratio (R:S)} = \frac{\text{Dry weight of root}}{\text{Dry weight of shoot}} \text{ g/g}$$

## 2.5 AMF Colonization

Fresh feeder root subsamples were taken at the time of harvest to determine the percentage of AMF colonization of the roots. The feeder roots were thoroughly washed in distilled water and fixed in 50% ethanol. Clearing and staining were carried out following the method of Koske and Gemma (1989). The stained roots were assessed for AMF colonization using the gridline intersect method of Giovannetti and Mosse (1980). Stained roots were spread on a gridline plate and viewed under a dissecting microscope at x45 magnification. The percentage root colonization was computed using the following equation;

$$\text{Root colonization (\%)} = \frac{\text{Number of arbuscular mycorrhizal-positive segments}}{\text{Total number of segments examined}} \times 100$$

## 2.6 Statistical Analysis

The mean values obtained from the replicate readings (n) were used to calculate the standard error. Differences between means were determined by one-way analysis of variance (ANOVA) and Duncan's multiple range test was used to separate the means at P <0.05. All data were analyzed using the statistical package for social sciences (SPSS) (version 20.0 for windows).

# 3. RESULTS

## 3.1 Soil and AMF Colonization

The physicochemical analysis of the experimental soil used revealed that physically it was composed of 78.7% sand; 9.0% silt; 12.3% clay; 1.86% organic matter with a sandy loam texture and a pH of 5.69. Chemically, it was made up of 0.08mg/kg total nitrogen; 85.0mg/kg available P; 1.40cmol/kg Ca; 0.8cmol/kg and 0.11cmol/kg K.

Mycorrhizal colonization was not found in the roots of *P.mildbraedii* control seedlings. Salt stress in most cases generally increased the percentage root AMF colonization (Table 1). At 12WAE all the salt stressed seedlings had higher percentage root AMF colonization than their unstressed counterparts. Meanwhile, seedlings inoculated with *G.intraradices* under NaCl stress (*GiNaCl*) recorded the highest significant ( $P \leq 0.05$ ) AMF root colonization of 96.24% and the lowest was 66.77% in seedlings inoculated with *G.intraradices* (Table 1).

**Table 1: Effect of mycorrhizal inoculation on the AMF colonization of *Pterocarpus mildbraedii* seedlings under salt stress**

Treatment	% AMF at harvest
Control	0.00
<i>Ge</i>	86.14±2.14 <sup>bc</sup>
<i>GeNaCl</i>	89.86±0.78 <sup>bc</sup>
<i>Gi</i>	66.77±2.30 <sup>a</sup>
<i>GiNaCl</i>	96.24±0.96 <sup>c</sup>
<i>Gm</i>	69.49±7.77 <sup>a</sup>
<i>GmNaCl</i>	83.95±0.03 <sup>b</sup>
<i>Go</i>	80.83±3.60 <sup>b</sup>
<i>GoNaCl</i>	91.31±3.33 <sup>bc</sup>

\* Means of three replicates ± standard errors of mean (S.E.M). Means within each column followed by different letters are significantly different at  $P < 0.05$  according to Duncan's multiple range test.

### 3.2 Plant Growth

Generally, AMF inoculated seedlings had higher significant ( $P \leq 0.05$ ) dry matter yields than un-inoculated (control) plants regardless of the 75mM NaCl treatment (Table 2). However, salt stress significantly ( $P \leq 0.05$ ) decreased the root, stem, leaf, and total dry weight of the *P.mildbraedii* seedlings compared to the inoculated non-stressed seedlings.

Furthermore, the highest values of root, stem, leaf and total dry weight ( $P \leq 0.05$ ) values of 5.83, 3.73, 6.47 and 16.03g / plant were recorded, respectively, by seedlings inoculated with *G.intraradices*.

The root: shoot ratio did not follow any consistent pattern and un-inoculated control seedlings gave the highest significant ( $P \leq 0.05$ ) mean value of 0.64g/g and the lowest value of 0.48g/g was observed in *G.intraradices* inoculated stressed (*GiNaCl*) seedlings (Table 2). Meanwhile, the difference in root: shoot ratio between *G. intraradices* inoculated seedlings (*Gi*) and *G.intraradices* stressed seedlings (*GiNaCl*) was significant ( $P \leq 0.05$ ).

**Table 2: Effect of mycorrhizal inoculation on the dry biomass of *Pterocarpus mildbraedii* seedlings under salt stress at harvest (12 WAE).**

Treatment	Root Dry Weight (g/plant)	Stem Dry Weight (g/plant)	Leaf Dry Weight (g/plant)	Total Dry Weight (g/plant)	Root : Shoot Ratio (g/g)
Control	2.81±0.06a	1.25±0.09a	3.17±0.13a	7.07±0.19a	0.64±0.02c
<i>Ge</i>	3.93±0.14bc	2.20±0.20abc	5.13±0.18c	11.27±0.52c	0.56±0.02b
<i>Ge</i> NaCl	3.02±0.09ab	1.70±0.17ab	3.70±0.35ab	8.42±0.48ab	0.57±0.03b
<i>Gi</i>	5.83±0.29e	3.37±0.40e	6.47±0.26e	16.03±0.96e	0.57±0.01b
<i>Gi</i> NaCl	3.30±0.18ab	2.55±0.20cd	4.33±0.17b	10.18±0.54abc	0.48±0.00a
<i>Gm</i>	5.00±0.45d	3.70±0.40e	5.63±0.19cd	14.00±1.15de	0.54±0.02ab
<i>Gm</i> NaCl	3.20±0.20ab	2.15±0.09bc	4.53±0.40b	9.45±0.63abc	0.51±0.01ab
<i>Go</i>	4.51±0.44cd	3.13±0.09de	6.03±0.39de	13.68±0.93d	0.49±0.03a
<i>Go</i> NaCl	3.10±0.20ab	2.00±0.06abc	4.03±0.07b	9.13±0.33abc	0.51±0.02ab

\* Means of three replicates ± standard errors of mean (S.E.M). Means within each column followed by different letters are significantly different at P <0.05 according to Duncan's multiple range test.

#### 4. DISCUSSION

Plant productivity and growth are negatively affected by various biotic and abiotic stresses. Salinity stress is detrimental to trees and has greatly affected the structure of coastal forests and afforestation projects (Wang *et al.*, 2019) in Nigeria (Osim, 2021b). However, a possible approach to alleviate salinity includes inoculating transplants by AMF (Othman *et al.*, 2022). AMF improves salinity tolerance in plants by altering biochemical and physiological processes, such as improving nutrient availability, water uptake, ionic homeostasis and increasing photosynthetic efficiency (Khaliq *et al.*, 2022).

There was no AMF colonization in *P. mildbraedii* control seedlings due to non-mycorrhization. AMF colonization depends on the host specificity, plant genotypes, nutrient availability and alteration in nutrient uptake under diverse environmental conditions (Chandrase Karan, 2022). In the present study the result indicates that inoculated seedlings of *P. mildbraedii* under NaCl stress present higher AMF colonization percentages than their unstressed counterparts. Although colonization percentages differed among AMF strains. These findings are consistent with the reports by Osim (2021a) and Osim (2021b) that the percentage colonization of AMF in *Terminalia ivorensis* and *P. mildbraedii* seedlings increases significantly with 75mM NaCl stress, suggesting that sporulation could be stimulated under salt stress that will increase the length of the hyphal and branched absorbing structures (BAS) of Fungi (Jahromiet *et al.*, 2008). This is in contrast to the findings of Wang *et al.*, (2019), who reported a reduction in AMF colonization percentages of *Zelkova serrata* seedlings under salt stress, due to suppression of AMF hyphal growth, sporulation and spore germination.

Also, the highest significant ( $P \leq 0.05$ ) percentage root AMF colonization recorded by *P. mildbraedii* seedlings inoculated with *G. intraradices* under NaCl stress (GiNaCl) is similar to the work of Osim (2021a) who reported that *Terminalia ivorensis* seedlings inoculated with *G. intraradices* under NaCl stress gave the highest percentage root AMF colonization which could be due to the colonization capacity of *G. intraradices* and the species of trees (Wang *et al.*, 2019).

The results of the present study clearly demonstrate that inoculation with AMF can lead to a great stimulation of dry matter yield of *P. mildbraedii* seedlings, the magnitude of this improvement varying between mycorrhizal treatments, with or without NaCl stress. Inoculation resulted in significant ( $P \leq 0.05$ ) increases in dry matter content over the non-inoculated control seedlings, and there were also significant ( $P \leq 0.05$ ) differences between the different mycorrhizal fungi in their ability to enhance dry matter content of *P. mildbraedii*. This observation is consistent with the results of previous studies on the accumulation of dry biomass of *Terminalia ivorensis* (Osim, 2021a). It is also consistent with the reports of Augeet *al.*, (2014), Baltazar-Bernat *et al.* (2022) and Chandrasekaran (2022), that varying responses of different fungi suggest that symbiotic efficiency is determined not only by the species and races of the fungus but also by the host species and environmental conditions. This could favor some physiological processes for the plant like water, nutrient uptake and photosynthetic efficiency, keeping higher osmoprotectants and enzyme activities and a high  $K^+/Na^+$  ratio.

Plant growth and biomass suffered a lot under salinity stress. Salt stress inhibits the photosynthetic ability of plants leading to decrease in plant biomass accumulation and growth (Wang *et al.*, 2019). However, the significant ( $P \leq 0.05$ ) decrease in dry matter content under 75 mM NaCl stress in this work may be due to the fact that plants exposed to salinity are subject to three primary physiological stresses; the toxicity of sodium ( $Na^+$ ) and chloride ( $Cl^-$ ) ions, the danger of physiological drought, and nutrient imbalance (Wang *et al.*, 2018). In this present work, the effects of NaCl stress on dry biomass may have been alleviated by AMF colonization (Table 2). These findings are similar to the observations of Al-Karaki (2006); Giri and Mukerji (2004); Evelinet *al.*, (2012) who reported that AMF can alleviate the salt stress on plants by inhibiting the high uptake of  $Na^+$  and  $Cl^-$  and their transfer to the shoot. The effects of mycorrhizae on development depend on the type and dose of AMF in relation to the host plant species (Baltazar-Bernal *et al.*, 2022).

The biomass allocation in this work was high in the leaf tissues compared to that of stem or root tissues signifying that leaf is the primary sink in *P. mildbraedii* seedlings (Table 2). Root growth in *P. mildbraedii* appears to be less affected in spite of the negative effects of NaCl on roots, while the stem was drastically affected, so the dry weight ratio was higher in seedlings grown under NaCl stress than in the control environment (Table 2). These observations are similar to the results of Osim (2021a) on dry biomass accumulation of *Terminalia ivorensis* seedlings and also the reports of Singhet *al.*, (2012) and Giriet *al.*, (2007) on tomato. This could be attributed to the uptake of carbohydrates from the plant, AMF enhancement of the sink capacity of the root system and this in turn increase the photosynthetic performance of plant leading to improved plant growth (Al-karaki and Williams 2021). The enhanced mycorrhizal tolerance to salinity stress was closely associated to a higher accumulation of soluble sugars in plant roots (Fenget *al.*, 2022).

The highest significant root: shoot ratio ( $P < 0.05$ ) recorded by noninoculated control seedlings in this present research may have been due to the significant decrease ( $P \leq 0.05$ ) decrease in shoot (stem) dry weight due to nonarbuscular mycorrhizal colonization (absence of colonies in the roots) (Feng *et*

al., 2002 and Al-karaki, 2006). This is in contrast with the results of previous researchers (Osim, 2021a and Giri *et al.*, 2007) who observed a higher root: shoot ratio in *Terminalia ivorensis*, tomato and *Acacia nilotica* inoculated with AMF than non-mycorrhizal seedlings. This could be due to the availability of more nutrients to plant roots in direct contact with the soil as a result of the synergistic effect of the symbionts that results in an increase in root dry weight (Diagne *et al.*, 2020).

Physiological differences exist between mycorrhizal and non-mycorrhizal plants, and between mycorrhizal fungi in their effects on host species (Baltazar-Bernal *et al.*, 2022). The highest significant ( $P \leq 0.05$ ) dry biomass (root, stem, leaf and total dry weight) of *P. mildbraedii* seedlings inoculated with *G. intraradices* found in this present research is in harmony with the results of previous work on *Terminalia ivorensis* seedlings (Osim, 2021a). He reported the highest root, stem, leaf and total dry weight in *Terminalia ivorensis* seedlings inoculated with *G. occultum*. An increase in the developmental and biomass variables of plants inoculated with mycorrhizae has been observed. The effects of the mycorrhizae on development depend on the type and dose of AMF in relation to the host plant species (Baltazar-Bernal *et al.*, 2022). Also, AMF ability to absorb nutrient is found to vary with the fungal and plant species and concentration of salt in soil (Wang *et al.*, 2019; Diagne *et al.*, 2020).

## 5. CONCLUSION

AMF acts as a growth regulator in most terrestrial environments and has been used as a bioameliorator of saline soil. The result of this study has shown that inoculation with appropriate AMF can enhance biomass production and accumulation of *P. mildbraedii* seedlings in saline (75mM level) sandy loam soil of Cross River Basin in Calabar Nigeria. Therefore, the use of *G. intraradices* as the most promising candidate in the rehabilitation of afforestation and vegetation in coastal areas of Nigeria is recommended.

## REFERENCES

- Abbaspour, H., Fallahyan, F. & Fahimi, H. (2005). Effects of endo-mycorrhizal fungi and salt stress on nutrient acquisition and growth of *Pistacia avara* L. Asian Network. *Pakistan Journal of Biological Sciences*, 8 (7): 1006-1010.
- Al-karaki, G. N. (2006). Nursery inoculation of tomato with arbuscular mycorrhizal fungi and subsequent performance under irrigation with saline water. *Scientia Horticulturae*, 109: 1-7.
- Al-karaki, G. N. and Williams, M. (2021). Mycorrhizal mixture affect the growth, nutrition, and physiological responses of soybeans to water deficit. *Acta Physiologiae Plantarum*, 43, 1-9.
- A.O. A. C. Association of Official Analytical Chemist. Methods of Analyss, Washington, DC, USA. 2003.
- Auge, R. M., Toler, H. D. and Saxton, A. M. (2014). Arbuscular mycorrhizal symbiosis and osmotic adjustment in response to NaCl stress: A meta-analysis. *Frontier of Plant Science*, 5, 562.

- Babar, S., Giriraj, A., Reddy, C. S., Jentsch, A. & Sudhaka, S. (2012). Species distribution models: Ecological explanation and prediction of an endemic and endangered plant species (*Pterocarpus santalinus* L. F.) *Current Science*, 102(8): 1157-1165.
- Baltazar-Bernal, O., Spinoso-Castillo, J. L., Mancilla-Alvarez, E. and Bello-Bello, J. J. (2022). Arbuscular mycorrhizal fungi induce tolerance to salinity stress in Taro plantlets (*Colocasia esculenta* L. Scgott) during acclimatization. *Plants*, 11, 1780.
- Chandrasekaran, M. (2022). Arbuscular mycorrhizal fungi mediated enhanced biomass, root morphological traits and nutrient uptake under drought stress: A meta-analysis. *Fungi*, 8, 660.
- Choi, J., Summers, W. and Paszkowski, U. (2018). Mechanisms underlying establishment of arbuscular mycorrhizal symbioses. *Annual Review of Phytopathology*, 56, 135-160.
- Diagne, N., Ngom, M., Djighaly, P. I., Fall, D., Hoher, V. & Svistoonoff, S (2020). Roles of Arbuscular Mycorrhizal fungi on plant growth and performance: importance in biotic and abiotic stressed regulation. *Diversity*, 12 (370); 1-25.
- Evelin, H., Giri, B. and Kapoor, R. (2012). Contribution of *Glomus intraradices* inoculation to nutrient acquisition and mitigation of ionic imbalance in NaCl-stressed *Trigonella foenum-Graecum*. *Mycorrhiza*, 22, 203-217.
- Feng, G., Zhang, F.S., Li, X. L, Tiang, C.Y., Tang, C. and Rengel, Z. (2002). Improved tolerance of maize plants to salt stress by arbuscular mycorrhiza is related to higher accumulation of soluble sugars in roots. *Mycorrhiza*, 12: 185 -190
- Giovannetti, M. & Mosse, B. (1980). An evaluation of techniques for measuring vesicular arbuscular mycorrhizal infection in roots. *New Phytologist*, 84: 489-500.
- Giri, B. and Mukerji, K. G. (2004). Mycorrhizal inoculants alleviates soil stress in *Sesbania aegyptiaca* and *Sesbania grandiflora* under field conditions: evidence for reduced sodium and improved magnesium uptake. *Mycorrhizal*, 14: 307-312.
- Giri, B., Kapoor, R. and Mukerji, K. G. (2007). Improved tolerance of *Acacia nilotica* to salt stress by arbuscular mycorrhizal, *Glomus fasciculatum* may be partly related to elevated  $K^+/Na^+$  ratios in root and shoot tissues. *Microbial Ecology*, 54: 753 – 760.
- International Legume Database & Information Service (2005). Genus *Pterocarpus*. Version 10.01, November 2005. ILDIS Retrieved November, 2008 from [https://www.revolvy.com/topic/Pterocarpus&item\\_type=topic](https://www.revolvy.com/topic/Pterocarpus&item_type=topic).
- Iwena, O. A. (2008). *Essential geography*. Ibadan: Tonal Publishers.
- Jahromi, F., Aroca, R, Porcel, R. and Ruiz-Lozano, J.M. (2008). Influence of salinity on the in vitro development of *Glomus intraradices* and on the in vivo physiological and molecular responses of mycorrhizal lettuce plants. *Microbial Ecology*, 55:45-53.

- Khaliq, A., Perveen, S., Alamer, K. H., Haq, M. Z. U., Rafique, Z., Alsudays, I. M., Althobaiti, A. T., Saleh, M. A., Hussain, S. and Attia, H. (2022). Arbuscular mycorrhizal fungi symbiosis to enhance plant-soil interaction. *Sustainability*, 14, 7840.
- Koske, R. E. and Gemma, J. N. (1989). A modified procedure for staining roots to detect VA mycorrhizas. *Mycology Research*, (92): 486-488.
- Kumar, A., Dames, J. F., Gupta, A., Sharma, S., Gilbert, J. A. and Ahmad, P. (2015). Current developments in arbuscular mycorrhizal fungi research and its role in salinity stress alleviation: A biotechnological perspective. *Critical Review in Biotechnology*, 35; 461-474.
- Lemmens, R.H.M.J., Soerianegara, I. & Wong, W.C. (1995). *Timber trees: Minor commercial timbers*. Leiden: Backhuys Publishers.
- Osim, S.E., (2021a). Dry biomass accumulation of *Terminalia ivorensis* (A. Chev) seedlings inoculated with arbuscular mycorrhizal fungi under salt stress. *Journal of Science, Engineering and Technology*, 8 (1): 10-17.
- Osim, S.E., (2021b). Effects of salt stress on chlorophyll content and foliar nutrient yield of *Pterocarpus mildbraedii* (Harms) seedlings inoculated with arbuscular mycorrhizal fungi. *Journal of Science, Engineering and Technology*, 8 (1): 26-35.
- Othman, Y. A., Tahat, M., Alananbeh, K. M. and Al-Ajlouni, M. (2022). Arbuscular mycorrhizal fungi inoculation improves flower yield and postharvest quality component of Gerbera grown under different salinity levels. *Agriculture*, 12, 978.
- Rabie, G.H. & Almadini, A.M. (2005). Role of bio-inoculants in development of salt-tolerance of *Vicia faba* plants under salinity stress. King Faisal University, Saudi Arabia. *African Journal of Biotechnology*, 25: 221-242.
- Rodrigues, R. S., Mascarenhas, A. & Jagtap, T. G. (2011). An evaluation of flora from coastal sand dunes of India: Rationale for conservation and management. *Ocean Coast. Management*, 54; 1818-188.
- Salwan, R., Sharma, A. and Sharma, V. (2019). Microbes mediated plant stress tolerance in saline agricultural ecosystem. *Plant soil*, 442, 1-22.
- Singh, J., Sastry, E. V. D. and Singh, V. (2012). Effect of salinity on tomato (*Lycopersicon esculentum* Mill.) during seed germination stage. *Physiological Molecular Biology of Plant*, 18(1): 45-50.
- Thangavel, P., Anjum, N. A., Muthukumar, T., Sridevi, G., Vasudhevan, P. and Maruthupandian, A. (2022). Arbuscular mycorrhizae: Natural modulators of plant-nutrient relation and growth in stressful environments. *Arch. Microbiol.* 204, 1-22.
- Wang, Y., Wang, M., Li, Y., Wu, A. and Huang, J. (2018). Effects of arbuscular mycorrhizal fungi on growth and nitrogen uptake of *Chrysanthemum morifolium* under salt stress. *PLOS ONE*, 13, e0196408.

Wang, J., Fu, Z., Ren, Q., Zhu, L., Lin, J., Zhang, J., Cheng, X., Ma, J., and Yue, J. (2019). Effects of Arbuscular Mycorrhizal Fungi on Growth, Photosynthesis, and Nutrient Uptake of *Zelkova serrata*(Thunb) Makino Seedlings under salt stress. *Forest*, 10 (186); 1-16.

World Agro-forestry Center (2008).Agro-forestry Tree Database *Pterocarpus soyauxii*.Retrieved November, 2008.

UNDER PEER REVIEW