

## ***In vitro* Diazotrophic Bacterial Response to Herbicide Toxicity**

### **ABSTRACT**

Diazotrophic bacteria play critical role in biological nitrogen fixation (BNF). Their growth and activities may be impeded in the presence of herbicides. To investigate this, 1.0 ml of broth stock culture containing  $2.9 \times 10^5$  cfu/ml and  $2.6 \times 10^6$  cfu/ml of *Azotobacter* and *Bradyrhizobium* respectively were inoculated into 10.0 ml of their respective growth medium [ TDC broth, and Yeast extract mannitol broth (YEMB)] containing 0, 0.5, 1.0, 1.5 and 2.0 % v/v of the respective herbicides, atrazine, glyphosate, paraquat and 2,4-D. Thereafter, plate counts of the diazotrophs for each concentration was taken at intervals of 24, 48, 72 and 120 h using spread plate method on TDC agar and YEMA after incubation at room temperature ( $30 \pm 2$  °C) for 72 h. The LC<sub>50</sub> of the respective herbicides for *Azotobacter* and *Bradyrhizobium* was determined at 120 h using Probit analysis. Results showed that all tested concentrations except control, retarded diazotrophic bacterial population growth. Growth reduction increased progressively with increased concentrations of herbicides ( $P = 0.05$ ). Though, diazotrophs exhibited differential responses to tested herbicides, generally there was 29.7 – 100 % growth suppression. The LC<sub>50</sub> indicated symbiotic *Bradyrhizobium* displayed greater sensitivity to tested herbicides than free-living *Azotobacter* ( $P < 0.05$ ). Conclusively, herbicides suppressed diazotrophic bacterial growth.

**Keywords:** *Azotobacter*, *Bradyrhizobium*, diazotrophs, herbicides, nitrogen fixation, toxicity

### **1. INTRODUCTION**

Microorganisms that have the ability to fix molecular nitrogen (N<sub>2</sub>) are collectively known as diazotrophs. Nitrogen fixation in soil is critically important for substantial growth and yield of crop plants. Biological nitrogen fixation in soil which involves microbial transformation of inorganic gaseous N<sub>2</sub> to organic forms such amino acids, pyrimidines and purines [1], represent one of the most important source of nitrogen addition to soil [2]. An estimated input of over 200 million tons of nitrogen is added to Earth's ecosystem annually via nitrogen fixation [3]. Biological nitrogen fixation is the second most significant biological phenomenon on earth after photosynthesis [4].

A number of diazotrophic bacteria such as *Azotobacter* and *Bradyrhizobium* are important microbial agents that contributes to nitrogen fixation in the ecosystem. While *Azotobacter* exist in soil as free-living nitrogen fixing agent, *Bradyrhizobium* fix atmospheric nitrogen in soil symbiotically in close association with leguminous plants. In addition to fixing nitrogen in soil, *Azotobacter* has been reported to enhanced the growth and wellbeing of many crop plants such as barley, wheat and potato [1]. Furthermore, they also enhance plant growth and yield via the production of plant growth hormones, synthesis of phytopathogenic inhibitors, stimulation of rhizospheric microorganisms and modification of nutrient uptake [5]. Nitrogen fixation through nodulation of legumes accounts for about 25 % of the total nitrogen fixed yearly on earth. Nodulated legumes display remarkable ability to grow well in unfertilized soil lacking nitrogen where other group of plants grow poorly [6]. *Bradyrhizobium* is known to exhibit cosmopolitan

existence in its nodulation ability over a broad variety of legumes such as groundnut (*Arachis hypogaea*), cowpea (*Vigna unguiculata*), soybean (*Glycine max*), Bambara groundnut (*Vigna subterranean*), common bean (*Phaseolus vulgaris*) and Kersting's bean (*Macrotyloma geocarpum*) [7].

Spurred by the need to increase food production to keep pace with all-time rise in human population, the application of herbicides in agriculture has in recent time witnessed an astronomical increase globally. The increase in herbicide usage has been linked to their ease of application, availability and effectiveness in weed control. Some of the most commonly applied herbicides includes, atrazine, glyphosate, butachlor, 2,4-D, paraquat among others. However, irrespective of their spectrum of activities on weed plants, they are non-discriminatory in their effect towards other unintended organisms in the ecosystem including microorganisms. In general, microbial populations of actinomycetes, bacteria, cyanobacteria, fungi and protozoa in the ecosystem have been reported to be negatively impacted by herbicides [8, 9]. This notwithstanding, knowledge of the specific response of certain important soil microbes such as diazotrophic bacteria whose roles is critical in maintaining soil fertility is required, as nitrogen is one of the major elements limiting crop growth and yield in agriculture. It is on the basis of ascertaining the specific toxicity response of two important diazotrophs, the free-living *Azotobacter* and symbiotic *Bradyrhizobium* to some commonly applied herbicides in agriculture that this study was carried out.

## 2. MATERIALS AND METHODS

### 2.1 *Azotobacter* and *Bradyrhizobium* isolation

*Azotobacter* was isolated from agricultural soil by collecting soil samples within 0-15 cm of soil top layer. One gram of the soil sample (obtained from 10.0 g of the homogenized soil sample) was added to 9.0 ml physiological saline in a test tube. From this, ten-fold serial dilutions ( $10^{-1}$ ,  $10^{-2}$ ,  $10^{-3}$ ,  $10^{-4}$  and  $10^{-5}$ ) were prepared and 0.1 ml aliquots of respective dilutions were subsequently plated out on TDC agar medium in triplicates using the spread plate method for *Azotobacter* isolation. The composition of TDC agar medium includes, glucose, 5.0 g;  $K_2HPO_4$ , 1.0 g;  $MgSO_4$ , 1.0 g;  $CaCO_3$ , 10.0 g and agar, 20.0 g in 1000.0 ml of distilled water [10].

On the other hand, *Bradyrhizobium* for this study was isolated from root nodules of 3-months old *Arachis hypogaea* as described by Ubogu et al. [11]. Mature well-formed nodules were pulled off from roots. One gram of the nodules was washed in tap water. This was then surface-sterilized in 70 % ethanol for 2 minutes and subsequently rinsed with distilled sterile water. Further surface-sterilization was carried out for 2 minutes using 3.5 % v/v sodium hypochlorite and immediately rinsed thrice with distilled sterile water after which nodules were crushed in a few drops of distilled sterile water in Mac Cartney bottle. After thorough crushing, this was made up to 10.0 ml using physiological saline. From this, ten-fold serial dilutions ( $10^{-1}$ ,  $10^{-2}$ ,  $10^{-3}$ ,  $10^{-4}$  and  $10^{-5}$ ) were prepared and 0.1 ml aliquots of respective dilutions were subsequently plated out on yeast extract mannitol agar (YEMA) in triplicates using the spread plate method for *Bradyrhizobium* isolation. The composition of the YEMA includes  $MgSO_4$ , 0.1 g; NaCl, 0.13 g;  $K_2HPO_4$ , 2.5 g; yeast extract, 0.5 g; agar powder, 20.0 g in 1000.0 ml of distilled water.

Both TDC agar and YEMA plates were incubated at  $30 \pm 2$  °C (room temperature) for 72 h. Pure culture of isolated colonies obtained were then transferred to respective TDC agar and YEMA slant for further studies.

## 2.2 Characterization of isolated *Azotobacter* and *Bradyrhizobium*

Pure culture of bacterial isolates was identified based on cultural, morphological and biochemical properties employing Bergey's Manual of Systemic Bacteriology (volume 2, Part B and C) [12 a and b].

## 2.3 In vitro toxicity assessment

*In vitro* toxicity response of *Azotobacter* and *Bradyrhizobium* to the herbicides, atrazine, glyphosate, 2,4-D and paraquat were evaluated in their respective growth medium (TDC broth and YEMB).

Using a sterile wire loop, bacterial growth from pure culture agar slant of *Azotobacter* was aseptically scrapped into 200 ml of TDC broth medium in 500 ml conical flask. Flask was then cocked with sterile cotton wool and incubated at  $30 \pm 2$  °C for 72 hours with regular hand-shaking of flask for 30 minutes every 12 hours. This served as the stock culture from which toxicity study was conducted.

One milliliter from the stock culture containing  $2.9 \times 10^5$  cfu/ml of *Azotobacter* were respectively inoculated into 10.0 mL of fresh TDC broth in a 15 ml test tube, containing the various concentrations of the respective test herbicides at 0, 0.5, 1.0, 1.5 and 2.0 % v/v. The TDC broth medium containing the respective test herbicides were then incubated at  $30 \pm 2$  °C for 120 hours. The total *Azotobacter* counts (cfu/ml) for each tested herbicide concentration were determined in triplicates at time interval of 24, 48, 72 and 120 hours. Counts were determined by plating out 0.1 ml of serially diluted samples of each tested concentration of the tested herbicides on TDC agar medium using the spread plate method. Plates were incubation at  $30 \pm 2$  °C for 72 hours. Thereafter, the total *Azotobacter* counts were taken and estimated in cfu/ml of sample.

Similar procedure employed for *Azotobacter* were repeated for *Bradyrhizobium*, except that YEMB and YEMA were the medium of growth, and the 1.0 mL from the broth stock culture inoculated into the respective broth containing the various concentrations of the tested herbicides and the control contained  $2.6 \times 10^6$  cfu/ml of *Bradyrhizobium*.

The  $LC_{50}$  of the respective tested herbicides for *Azotobacter* and *Bradyrhizobium* in the presence of their respective growth medium were determined at 120 hours of the study.  $LC_{50}$  was determined using Finney Probit Analysis [13].


## 2.4 Statistical analysis

All the data garner from this study were analyzed using Microsoft Excel (Analysis Tool Pak). Triplicate data were analyzed using measures of central tendency and dispersion. The effect of herbicides concentration on *Azotobacter* and *Bradyrhizobium* populations were determined using analysis of variance (ANOVA), while the overall comparative sensitivity of *Azotobacter* and *Bradyrhizobium* to the test herbicides were determined using Student's *t*-test ( $P < 0.05$ ). Finney Probit Analysis was used for determining  $LC_{50}$  of tested herbicides at 95 % confident level.


### 3. RESULTS

#### 3.1 In vitro response of *Azotobacter* and *Bradyrhizobium* to herbicides


The populations of *Azotobacter* and *Bradyrhizobium* generally increased with time for the respective tested herbicides concentrations including the control. Nonetheless, there was a progressive reduction in population with increased concentrations of the herbicides ( $P < 0.05$ ) (Fig. 1). No observable growth occurred for *Azotobacter* beyond 1.5 % v/v for the herbicide atrazine within the tested period. There was no growth of *Bradyrhizobium* at 2.0 % v/v in glyphosate within the first 24 hours. Similarly, *Azotobacter* and *Bradyrhizobium* did not manifest any form of growth at all the tested concentrations of paraquat within the first 24 h of the study. However, after 24 h, there was substantial growth of *Azotobacter* and *Bradyrhizobium* at all the tested concentrations of glyphosate and paraquat within the period of study. Generally, the growth of diazotrophs were suppressed between 29.7 – 100 % by the tested herbicides (Table 1).




**Figure 1a.** Effect of different concentrations of atrazine on *Azotobacter* and *Bradyrhizobium* populations. \*Values with same superscript alphabet (a, b, c, d, e) for different concentrations at 120 hours did not differ significantly for same tested organism (n = 15, ANOVA,  $P < 0.05$ ).



**Figure 1b.** Effect of different concentrations of glyphosate on *Azotobacter* and *Bradyrhizobium* populations. \*Values with same superscript alphabet (a, b, c, d, e) for different concentrations at 120 hours did not differ significantly for same tested organism (n = 15, ANOVA,  $P < 0.05$ ).



**Figure 1c.** Effect of different concentrations of paraquat on *Azotobacter* and *Bradyrhizobium* populations. \*Values with same superscript alphabet (a, b, c, d, e) for different concentrations at 120 hours did not differ significantly for same tested organism (n = 15, ANOVA,  $P < 0.05$ ).



**Figure 1d.** Effect of different concentrations of 2,4-D on *Azotobacter* and *Bradyrhizobium* populations. \*Values with same superscript alphabet (a, b, c, d, e) for different concentrations at 120 hours did not differ significantly for same tested organism (n = 15, ANOVA,  $P < 0.05$ ).

**Table 1.** Percentage population growth suppression of *Azotobacter* and *Bradyrhizobium* by herbicides at 120 hours

		Percentage (%) population growth suppression			
		Herbicide conc. (% v/v)			
Bacteria	Herbicide	0.5	1.0	1.5	2.0
<i>Azotobacter</i>	Atrazine	45.8 <sup>a</sup>	51.5 <sup>a</sup>	100.0 <sup>a</sup>	100.0 <sup>a</sup>
<i>Bradyrhizobium</i>		29.7 <sup>b</sup>	35.7 <sup>b</sup>	36.2 <sup>b</sup>	44.0 <sup>b</sup>
<i>Azotobacter</i>	Glyphosate	53.8 <sup>a</sup>	58.1 <sup>a</sup>	58.1 <sup>a</sup>	61.3 <sup>a</sup>
<i>Bradyrhizobium</i>		29.8 <sup>b</sup>	48.0 <sup>b</sup>	54.6 <sup>b</sup>	62.1 <sup>a</sup>
<i>Azotobacter</i>	Paraquat	28.0 <sup>a</sup>	30.3 <sup>a</sup>	34.0 <sup>a</sup>	34.6 <sup>a</sup>
<i>Bradyrhizobium</i>		36.4 <sup>b</sup>	46.9 <sup>b</sup>	48.0 <sup>b</sup>	63.0 <sup>b</sup>
<i>Azotobacter</i>	2,4-D	25.8 <sup>a</sup>	35.5 <sup>a</sup>	37.5 <sup>a</sup>	44.1 <sup>a</sup>
<i>Bradyrhizobium</i>		29.0 <sup>b</sup>	50.9 <sup>b</sup>	55.0 <sup>b</sup>	57.5 <sup>b</sup>

\*Values with different superscript alphabet (a, b), along the same column for same tested herbicide and concentration differ significantly (n = 3, Student's *t* test,  $P < 0.05$ )

### 3.2 LC<sub>50</sub> of tested herbicides on *Azotobacter* and *Bradyrhizobium*

For *Azotobacter*, the LC<sub>50</sub> of all the herbicides were greater than the highest tested concentrations of 2.0 % v/v except for atrazine. Conversely, for *Bradyrhizobium* the LC<sub>50</sub> of all the herbicides were less than the highest tested concentrations except for atrazine (Table 2). implies that at the tested concentrations. Generally, *Bradyrhizobium* displayed more sensitivity to the tested herbicides than *Azotobacter* ( $P < 0.05$ ).

**Table 2.** LC<sub>50</sub> of tested herbicides on *Azotobacter* and *Bradyrhizobium* at 120 h.

Herbicide	LC <sub>50</sub> (% v/v)	
	<i>Azotobacter</i>	<i>Bradyrhizobium</i>
Atrazine	1.6	5.0
Glyphosate	4.5	1.1
Paraquat	28.7	1.3
2,4-D	3.13	1.0

\*LC<sub>50</sub> determined using Probit analysis at 95 % confident level

#### 4. DISCUSSION

In this study, though there was a progressive rise in the population of the two diazotrophic bacteria (*Azotobacter* and *Bradyrhizobium*) with time, their growth rates were substantially retarded by the four tested herbicides (atrazine, glyphosate, paraquat and 2,4-D) in comparison to the control devoid of the herbicides. This finding is in concordance with that of Nahi et al. [14], who reported that all the tested herbicides (2,4-D, pretilachlor and paraquat) significantly decreased the growth of the diazotroph *Stenotrophomonas maltophilia*. Milosevia and Govedarica [15], also reported significant reduction in the populations of *Azotobacter* spp. and *Bradyrhizobium japonicum* with the application of a number of herbicides. Herbicides application to soil have been reported to reduce the number and diversity of diazotrophic bacteria [16].

Furthermore, studies have shown that herbicides application negatively impacted nitrogen fixation at certain concentrations [17]. The growth of the diazotrophs in this study generally decreased with increased concentrations of the tested herbicides. However, the degree to which the growth decreased varied with the diazotroph and type of herbicide. Similar to the finding in this study, Mohamed et al. [18], reported increased negative impacts of glyphosate and paraquat on the symbiotic nitrogen fixing bacteria, *Rhizobium nepotum*, *Rhizobium tibeticum*, *Rhizobium radiobacter* and *Pantoea agglomerans* as the concentrations of the herbicides increased.

The free-living (*Azotobacter*) and symbiotic (*Bradyrhizobium*) diazotrophs exhibited differential response to the tested herbicides. While the growth of *Azotobacter* was completely inhibited throughout the study period at concentration of above 1.5 % v/v of atrazine, *Bradyrhizobium* growth was inhibited by glyphosate at 2.0 % v/v only for the first 24 h. In the same vein, *Azotobacter* and *Bradyrhizobium* which were inhibited by all concentrations of paraquat within the first 24 h recovered from the growth inhibition thereafter. The findings here corroborate that of Latha and Gopal [8], who reported that pyrazosulfuron, butachlor, 2,4-D and pretilachlor initially decrease the population of *Azospirillum lipoferum* in comparison to the control but subsequently witnessed increased growth after 24 hours. Nahi et al. [14], also reported the recovery of the diazotroph *Stenotrophomonas maltophilia* after 7 days following initial inhibition

at higher concentration of tested herbicides. The diazotrophic bacterial recuperation following initial inhibition of growth may be attributed to their inherent resilience and capacity to adjust to the herbicides at specific concentrations over time. Comparatively, the symbiotic diazotroph *Bradyrhizobium*, displayed more sensitivity to the tested herbicides than the free-living *Azotobacter*. A number of studies lend credence to the finding here reported that diazotrophic bacterial response differ with the type of applied agrochemical pesticides [14, 19, 20]. The differential responses of the diazotrophic bacteria to the tested herbicides may be ascribed to the dissimilarity in the chemical composition of the herbicides and genetic make-up of the microorganisms.

## 5. CONCLUSION

The results of this study showed that diazotrophic bacterial population growth were retarded by atrazine, glyphosate, paraquat and 2,4-D at all the tested concentrations. Though, the degree of herbicides toxicity varied among the two tested diazotrophs with the symbiotic *Bradyrhizobium* displaying more sensitivity to the tested herbicides than the free-living *Azotobacter*, this implies that the application of these herbicides in agriculture have the potential to significantly suppress biological nitrogen fixation, a critical element required for the maintenance of soil fertility.

## REFERENCES

- [1] Willey JM, Sherwood LM, Woolverton CJ. Prescott's Microbiology. 10<sup>th</sup> ed. New York: McGraw-Hill; 2017.
- [2] Yadav A, Singh AL, Kumar A, Singh AK, Pandey KD, Singh M. Rhizobium-legume symbiosis and effects of environmental stresses on the symbiosis. In: Choudhary KK, Dhar DW, editors. Microbes in Soil and their Agricultural Prospects. New York: Nova Science Publishers, Inc.; 2015.
- [3] Rascio N, La Rocca N. Biological nitrogen fixation. In: Jorgensen SE, Faith BD, editors. Encyclopedia of Ecology. Amsterdam: Academic Press; 2008.
- [4] Unkovich M, Herridge D, Peoples M, Cadisch G, Boddey B, Giller K, Alves B, Chalk P. Measuring plant-associated nitrogen fixation in agricultural systems. Australian center for International Agricultural Research (ACIAR) Monographs No. 136, 258pp., 2008. [www.aciar.gov.au](http://www.aciar.gov.au).
- [5] Lenart A. Occurrence, characteristics, and genetic diversity of *Azotobacter chroococcum* in various soils of Southern Poland. Pol J Environ Stud. 2012; 21: 415-424.
- [6] Madigan MT, Martinko JM, Stahl DA, Clark DP. Brock's biology of microorganisms, 13<sup>th</sup> ed. New York: Benjamin Cummings; 2012.
- [7] Jaiswal SK, Dakora FD. Widespread distribution of highly adapted *Bradyrhizobium* species nodulating diverse legumes in Africa Front Microbiol. 2019; doi:10.3389/fmicb.2019.00310.
- [8] Latha PC, Gopal H. Effect of herbicides on soil microorganisms. Ind J Weed Sci. 2010; 42:217-222.

- [9] Tyagi S, Mandal SK, Kuma R, Kuma S. Effect of different herbicides on soil microbial population dynamics in rabi maize (*Zea mays* L.) Int J Curr Microbiol Appl Sci. 2018; (Spl issue 7) 7: 3751-3758.
- [10] Atlas RM. Handbook of microbial media, 4<sup>th</sup> ed. Washington D.C: ASM Press; 2010.
- [11] Ubogu M, Akponah E, Odeh A. *Arachis hypogaea* yield analysis in rhizobial (*Bradyrhizobium* sp.) inoculated soil-solarized soil. Clin Biotechnol Microbiol. 2018; 2(2): 331-339.
- [12a] Brenner DJ, Krieg NR, Staley JT. Bergey's manual of systemic bacteriology, vol 2, part B (The *Gammaproteobacteria*), 2nd ed. USA: Springer; 2005.
- [12b] Brenner DJ, Krieg NR, Staley JT (2005) Bergey's manual of systemic bacteriology, vol 2, part C (The *Proteobacteria*), 2nd ed. USA: Springer; 2005.
- [13] Finney DJ. Probit analysis, Reissue ed. New York: Cambridge University Press; 2009.
- [15] Nahi A, Othman R, Omar D, Ebrahimi M. Effects of selected herbicides on growth and nitrogen fixing activity of *Stenotropomonas maltophilia* (Sb16). Pol J Microbiol. 2016; 65 (3): 377- 382.
- [16] Milosevia NA, Govedarica MM. Effect of herbicides on microbiological properties of soil. Proc Nat Sci, Matica Srpska Novi Sad, No. 102, 5-21; 2002.
- [17] Angelini J, Silvina G, Taurian T, Ibanez F, Tonelli M L, Valetti L, Anzuay MS, Luduena L, Munoz V, Fabra A. The effects of pesticides on bacterial nitrogen fixers in peanut-growing area. Arch Microbiol. 2013; 195:683 – 692.
- [18] Mohamed M, Aliyat FZ, Ben Messaoud B, Simone C, Marina M, Filippo G, Laila N, Jamal I. Effects of pesticides use (glyphosate and paraquat) on biological nitrogen fixation. Water Air Soil Pollut.2021; doi:10.1007/s11270-021-05367-x.
- [19] Khudhur AM, Askar KA. Effect of some pesticides on growth, nitrogen fixation and *nif* genes in *Azotobacter chroococcum* and *Azotobacter vinelandii* isolated from soil. J Toxicol Environ Health Sci.2013; 5(9):166 – 171.
- [20] Gulhane PA, Gomashe AV, Sundarkar KM. Influence of pesticides on nitrogen fixing bacteria. Int J Tech Res Appl.2015; 3(4):157 – 160.