

Suppression of Chickpea (*Cicer arietinum* L.) Fusarium Wilt by *Bacillus* spp., *Pseudomonas* sp. and Rhizobacterial isolate

Suppression of *Fusarium* Wilt of Chickpea by *Bacillus* spp., *Pseudomonas* sp. and Rhizobacterial isolate

Abstract

This experiment was conducted to evaluate antagonistic effects of some selected rhizobacteria on *Fusarium oxysporium* f.sp. *ciceris* in pot experiment. Rhizobacterial isolates (one isolate of *Pseudomonas*, eight isolates of *Bacillus* genera and one bacterium isolate) and two chickpea cultivars (Shendi and Burgeig) were arranged in factorial pot experiment in CRD with four replicates. Disease incidence and severity were detected weekly. Disease reduction percentage was estimated at the end of the study. Generally, the application of rhizobacterial isolates as biological control agent reduced disease incidence compared with the control in both cultivars. The incidence in cultivar Shendi occurred at the third week after inoculation when treated with *Pseudomonas stutzeri* strain W28 (SA3) and *Bacillus subtilis* strain CM14(SA9). For the two cultivars, Shendi and Burgeig, the *Geobacillus* sp. CRR1-HN-1(SA2) and *Bacillus* sp. (SA1), respectively, had the highest positive effect on disease incidence and severity throughout the experiment compared with the control. ³Department of Botany, Faculty of Science, University of Khartoum, Sudan, these were 45.36% and 44.82% in incidence; 55.36% and 63.89% in severity, respectively.

Keywords: *biological control, disease incidence, severity, in vivo screening*

1. Introduction

Chickpea (*Cicer arietinum* L.) is one of the important food legumes, it is cultivated in more than 57 countries, it stand third in production following dry bean and peas with the productivity of about 913 kg ha (FAO, 2012). The cultivated chickpea originated in south-eastern Turkey (Van der Maesen, 1984). In Sudan, it is a cash crop that generate income for farmers and rural communities and as significant source of protein for poor people. Despite this, production fluctuates widely and farmers face a number of debilitating constrains: the wide spread incidence of diseases, the destructive activities of pests, parasitic weeds, and limited access to quality high-yielding cultivars. The project ICARDA ~~has~~ demonstrated high-yield varieties of chickpea to farmers and other stakeholders in the Gezira region, and other areas throughout the River Nile State. In the Gezira, the varieties Salwa and Burgaig ~~have~~ performed extremely well, generating and average 4.01 and 3.84 t/ha, respectively, far higher the 1.66 t/ha average achieved by traditional crops (ICARDA, 2014). More than 50 pathogens ~~have been was~~ reported so far to infect chickpea in different parts of the world, but only a few of them ~~have had~~ the potential to devastate the crop, The important diseases are *ascochyta* blight, dry root rot, stunt (caused by bean leaf roll virus), *botrytis* gray mold, collar rot, black root rot, *phytophthora* root rot, *pythium* root and seed rot and *Fusarium* wilt (Nene *et al.*, 1991). *Fusarium* wilt caused by *Fusarium oxyspoum* f.sp. *ciceris* is a major constraint to chickpea cultivation through the world (Nikam *et al.*, 2011). Yield losses attribute *Fusarium* wilt ~~vary-varied~~ from 10-15%, but the disease span completely destroyed the crop under unfavorable conditions (Cherif *et al.*, 2007). Plant growth promoting rhizobacteria (PGPR) ~~have been was~~ proved as biocontrol agents of soil borne plant pathogens, offer an attractive alternative to chemical fertilizers, pesticides and supplements. Thus, the use of PGPR is steadily increasing in agriculture (Ashrafuzzaman *et al.*, 2009). Use of biological control agents, such as plant growth promotion rhizobacteria (PGPR), can be suitable approach in control of disease (Schmidt *et al.*, 2004). Plant growth promotion rhizobacteria (PGPR), such as *Pseudomonas* and *Bacills* strain, ~~are were~~ the major root colonizers (Joseph *et al.*, 2007), and can elicit plant defenses (Kloepper *et al.*, 2004). Different mechanisms ~~have been was~~ reported for their performance such as production of antibiotics, siderphore cyanide hydrogen, competition for nutrition and space, inducing resistance, inactivation of pathogen enzymes and enhancement of root and plant development (Intana *et al.*, 2008). *Pseudomonas* and *Bacillus* strain ~~hadve~~ great potential in control of *Fusarium* wilt disease of chickpea (Anjajah *et al.*, 2003; Hervas *et al.*, 1997; Landa *et al.*, 1997). Plant growth promoting rhizobacteria (PGPR)

Formatted: Font: Italic

~~have been~~ reported as biocontrol agents of soil borne plant pathogen, offer an attractive alternative to chemical fertilizers, pesticides and supplements. Thus, the use of PGPR is steadily increasing, plant growth. Promoting rhizobacteria are an heterogeneous group of bacteria that can be found in the rhizosphere at the root surfaces and in association with roots which can improve the extent or quality of plant growth directly or indirectly (Joseph *et al.*, 2007; Datta *et al.*, 2011). This study was conducted to assess the antagonistic effect of- rhizobacterial isolates against ~~chickpea~~ *Fusarium* of chickpea wilt. The main objective of this study ~~is to~~ determination of effective rhizobacteria to be used for biological control of FOC.

2. Materials and Methods

The study was conducted to assess the antagonistic effect of ten rhizobacterial isolates against chickpea *Fusarium* wilt using two chickpea cultivars namely, Burgeig and Shendi. Two varieties of chickpeas were used in this study, namely cv. Shendi (susceptible) and cv. Burgaig (resistant). Soil was prepared by mixing sand and clay soil at 1:1 ratio. The soil was placed into 30×40 inch plastic sacks.

2.1 Rhizobacterial inocula:

Ten isolates of rhizobacteria SA1, SA2, SA3, SA4, SA5, SA6, SA7, SA8, SA9 and SA10 which were identified as *Bacillus* sp., *Geobacillus* sp., *Pseudomonas stutzeri* strain W28, *Bacillus subtilis* strain VT03, *Bacillus subtilis* strain FBRo3. *Bacillus* sp., Bacterium MOBOSA51, *Bacillus tequilensis* strain MML2, *Bacillus subtilis* strain CM14 and *Bacillus subtilis* strain SH23, respectively.

Rhizobacteria isolates were grown in Erlenmeyer flasks (250 ml) containing 100 ml of NA broth and shake for 24 hrs at rotary shaker, The growth was diluted with an adequate amount of non-inoculated nutrient broth to obtain a bacterial suspension of 10^8 -cfu/ml, using spectrophotometer (660 nm). Chickpea seeds were surface sterilized with 70% ethanol, then immersed for 2 minute in 2% sodium hypochloride and washed four times with sterilized distilled water and left for dried. 20 seeds were impressed in Petri-dish filled with bacterial suspension for 24 hrs, then placed onto sterile filter paper moistened with sterilized distilled water in Petri plates (four plates with 20 seeds/plate) and incubated at room temperature for 5-days. Control plates were arranged in similar way, except that they were treated with non-inoculated nutrient broth only.

2.2 Pathogen inocula:

Ten ml of sterilized water were added to each culture of the pathogen isolates, and the surface of the culture was scraped with a glass spatula to dislodge the chlamyospores. The spore suspensions were transferred to 100 ml sterilized flasks. Concentration of the suspensions were determined with a haemocytometer. High suspension of 9×10^2 spore ml^{-1} was prepared from each isolate ready for soil treatment. Half ml of the spores suspension was injected gently beside each one week old seedling using sterilized insulin syringe (Fisher and Toussoun, 1983).

2.3 Pot experiment

Pot experiment was conducted to evaluate the chickpea *Fusarium* wilt disease progress (disease incidence and severity). The germinated seeds of the two chickpea cultivars were treated by rhizobacteria isolates SA1, SA2, SA3, SA4, SA5, SA6, SA7, SA8, SA9 and SA10. In addition, the control represented by germinated seeds treated with non-inoculated nutrient broth. Treated germinated seeds were transferred into 30×40 inch plastic sacks. The plastic sacks filled with soil enclosed of sand and clay soil at ratio of 1:1. Treatments and the two cultivars were arranged in a factorial experiment in Completely Randomized Design (CRD) with four replicates; with one sack per replicate and three plants per sack.

2.4 Assessment of disease reaction

Disease reactions were assessed by the incidence and severity of symptoms at 7-day intervals. Severity of symptoms in individual plants of a microplot was assessed on a 0 to 4 rating scale based on the percentage of foliage with yellowing or necrosis in acropetal progression (0 = 0%, 1 = 1 to 33%, 2 = 34 to 66%, 3 = 67 to 100%, and 4 = dead plant). Incidence of foliar symptoms, (0-to-1 scale) (Landa, 1997; and Navas-Cortes *et al.*, 1998). The plants displaying the typical symptoms of the *Fusarium* wilt disease were considered infected. Percentage of the disease incidence was calculated using the following formula:

$$\text{Wilt incidence} = \frac{\text{No. of plants wilted}}{\text{Total No. of plants}} \times 100$$

Disease reactions were assessed according to the severity of symptoms weekly, the disease severity assessed by visual estimation adopting the scale shown as a follow:

0 = No infection* on leaf,

1=1-33% of the leaf were infected

Comment [DZ1]: Include here Reference

2= 34-66% of the leaf were infected

3= 67-100% of the leaf were infected

4= Dead plant

The disease-reduction percentage (DRP) was calculated by the method described by Yun Cao *et al.* (2011) using the following formula:

$$DRP = 1 - DT/DC \times 100$$

(DT = Disease incidence percentages in treatment; DC= Disease incidence percentages in control).

2.5 Statistical analysis

Statistical analysis for factorial experiments in Completely Randomize Design using STATISTIX 8.0 Analytical Software.

Formatted: Font: Italic

3. RESULTS AND DISCUSSION

3.1 Disease incidence progress:

Fusarium wilt disease cause yellowing and drying of leaves from the base to upward and finally death of plants. The overall progress of disease incidence in each cultivar presented in Figure 1 and Table 1. Generally, the application of rhizobacterial isolates as biological control agent reduced disease incidence compared ~~with to~~ the control in both cultivars. In Shendi cultivar SA1

and SA2 the incidence occurred in the second week after inoculation. The same trend observed in Burgeig when treated with SA6. Moreover, the incidence in cultivar Shendi occurred at the third week after inoculation when treated with SA3 and SA9.

For cultivar Shendi, the ten bacterial isolates compared with the control had a positive effect on disease incidence throughout the experiment except SA4, SA6, SA7 and SA8. For cultivar Burgeig, all bacterial isolates compared with the control had a positive effect on disease incidence throughout the experiment except SA4 and SA7. Schroth *et al.* (1981) and Cook *et al.* (1993) stated that there appears to be a nongeneral correlation between the *in vitro* ability of the antagonists and their ability to suppress diseases in the field. Juhnke *et al.* (1987) noted that the biological control depends upon maintaining a threshold population of the antagonist on planting material.

3.2 Main effect of cultivars and bacterial isolates on disease incidence

With the exception of cultivars non significant differences were observed among rhizobacterial isolates and for the rhizobacterial isolates \times cultivars interaction (Table 2).

The highest disease incidence was recorded in cultivar Shendi throughout the experiment. In the second week Shendi and Burgeig scored 27 and 11% respectively, where as in the 8th week 76 and 58% disease incidence were recorded for Shendi and Burgeig, respectively. This result confirms the finding obtained by Ahmed and Adam (2014).

3.3 Disease severity progress

The overall development of disease severity in each of the two cultivars was presented in Figure 2. For cultivar Shendi, all bacterial isolates compared ~~with to~~ the control had a positive effect on disease severity from the 4th week and onwards except SA4 and SA6. However, for cultivar Burgeig, the ten isolates had a positive effect on severity, compared ~~with to~~ the control, throughout the experiment except SA4 and SA6. Anjajah *et al.* (2003); Hervas *et al.* (1997) and Landa *et al.* (1997) reported that *Pseudomonas* and *Bacillus* strain have great potential in control of *Fusarium* wilt disease of chickpea.

3.4 Main effect of cultivars and bacterial isolates on disease severity

Significant difference ~~were~~~~was~~ observed between cultivars in all weeks except the 4th and 5th ones (Table 1). However non-significant differences were detected among the bacterial isolates in all readings. Additionally, non-significant rhizobacterial isolates × chickpea cultivars were obtained throughout the experiment (Table 3). Concerning the main effect of cultivars the highest disease severity throughout the experiment was observed in cultivar Shendi. In week two disease severity was 0.26 cultivar Shendi and 0.1 for cultivar Burgeig. In week eight it was 2.84 for cultivar Shendi and 1.91 for Burgeig.

3.5 Effect of rhizobacterial antagonist on disease index

The disease index measured in terms of disease incidence and severity. The first symptom of disease was appeared 12 days after

inoculation. Table 1 reveals the simple effects of bacterial antagonist on the disease in each cultivar in week eight as reduction percentages. SA2 and SA1 showed the highest disease incidence reduction (DIR%), this was 45.36% and 44.82% in cultivar Shendi and Burgeig, respectively. The same isolates had the highest disease severity reduction (DSR%), it was 55.36% and 63.89% in cv. Shendi and cv. Burgeig, respectively. On the other hand, SA4 and SA7 had the lowest DRP% in both cultivars, it was similar to the control. Moreover, SA4 scored the lowest DSR%. Karimi *et al.* (2012) reported that *B. subtilis* and *P. aueruginosa* isolates reduced disease severity more effects in seed treatment (39.47 and 34.21%), respectively. Zaim *et al.* (2013) noted that *Bacillus* isolates reduced disease severity caused by FOC from 60 to 99% in the field trials.




Figure 1. Main effect of cultivars on disease incidence throughout the experiment




Figure 2. Main effect of cultivars on disease severity throughout the experiment

Table 1. The effects of bacterial isolates on disease index of *Fusarium* wilt of chickpea in seed treatment in two cultivars after eight weeks from inoculation-

	Shendi		Burgeig		Shendi		Burgeig	
	DI	DIR (%)	DI	DIR (%)	DS	DSR (%)	DS	DSR (%)
SA1	66.25	27.60	41.25	44.82	2.50	36.22	1.08	63.88
SA2	50.00	45.36	58.00	22.41	1.75	55.36	1.99	33.44
SA3	66.25	27.60	57.75	22.74	2.42	38.27	1.75	41.47
SA4	91.50	0.00	75.00	-0.33	3.41	13.01	2.58	13.71
SA5	83.00	9.29	49.50	33.78	3.08	21.43	2.08	30.43
SA6	83.00	9.29	66.50	11.04	3.16	19.39	2.25	24.75
SA7	91.50	0.00	75.00	-0.33	3.33	15.05	2.42	19.06
SA8	83.00	9.29	49.75	33.44	3.08	21.43	1.16	61.20
SA9	58.25	36.34	58.00	22.41	2.17	44.64	1.25	58.19
SA10	74.75	18.31	41.50	44.48	2.42	38.27	1.42	52.51
control	91.50	0.00	74.75	0.00	3.92	0.00	2.99	0.00

DI= disease incidence , DS= disease severity,

DIR%= Disease incidence reduction percentage, DSR%= Disease severity reduction percentage

Formatted: Font: Not Italic

Table 2. Mean square for disease incidence in Chickpea cultivars, rhizobacterial isolates and rhizobacterial isolates x chickpea cultivar throughout the experiment-

Source of variation	df	Mean squares						
		Week2	Week3	Week4	Week5	Week6	Week7	Week8
Rhizobacterial isolates	10	702.35	620.77	802.57	1058.16	782.03	914.70	825.66
Chickpea cultivar	1	3319.04*	4006.26*	4279.02	8027.35*	5474.40	5792.49	4959.60
		*	*	*	*	*	*	*
Rhizobacterial isolates x Chickpea cultivar	10	737.11	603.67	609.06	100.11	167.00	189.99	226.40
Error	66	421.08	534.50	886.52	909.64	926.58	917.05	941.77
CV%		133.58	120.72	107.43	53.65	52.59	51.45	51.60

* and ** denote significant at 5% and 1% level of probability.

Table 3. Mean square for disease severity in Chickpea cultivars, rhizobacterial isolates and rhizobacterial isolates x chickpea cultivar throughout the experiment-

Source	df	Mean squares						
		Week2	Week3	Week4	Week5	Week6	Week7	Week8
Rhizobacterial isolates	10	0.13	0.28515	0.51852	1.27429	2.07155	2.42	2.70
Chickpea cultivar	1	0.61**	1.20*	1.29	4.09	9.52**	15.02**	19.10***
Rhizobacterial isolates x Chickpea cultivar	10	0.09	0.17	0.46	0.23	0.32	0.44	0.54
Error	66	0.08	0.26	0.52	1.08	1.34	1.49	1.58
CV %		154.28	149.37	138.74	72.29	59.39	55.90	53.04

* and ** denote significant at 5% and 1% level of probability.

4. Conclusion

Based on the study results, the *Geobacillus sp.* CRRI-HN-1 and *Bacillus sp.* had the highest positive effect on disease incidence and severity throughout the experiment compared with the control. Thus the *Fusarium* wilt can be conquered in chickpea by using the *Geobacillus sp.* CRRI-HN-1 and *Bacillus sp.* isolate. Further studies could be carried out in order to strengthen the result obtained in this experiment

Formatted: Font: Not Italic

Formatted: Justified

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Formatted: Font: Not Italic

References

Comment [DZ2]: The editor has to check according to style of Journal.

Ahmed N, and Adam A. (2014). Evaluation of Some Chickpea Cultivars for Resistance to *Fusarium* Wilt Disease in Sudan. Tropentag, "Bridging the gap between increasing knowledge and decreasing resources". Prague, Czech Republic. 6.14 : 4.438.

Anjajah, V, Cornlis P. and Kooedam, N. (2003). Effect genotype and root colonization in biological control of *Fusarium* wilts in pigeonpea and chickpea by *Pseudomonas aeruginosa* PNAI. *Canadian Journal of Microbiology*, Vol. 49, No. 2, 85-91.

Ashrafuzzaman M, Hossen F, Razi Ismail M, AnamulHoque Md, Zahurul Islam M, Shahidullah SM and Meon S. (2009). Efficiency of plant growth-promoting rhizobacteria (PGPR) for the enhancement of rice growth. *Afr. J. Biotechnol.* 8(7): 1247-1252.

Cherif M, Arfaoui A and Rhaiem A. (2007). Phenolic compounds and their role in bio-control of chickpea to fungal pathogenic attacks. *Tunisian Journal Plant Protection* 2:7-21.

Cook RJ. (1993). Making greater use of introduced microorganisms for biological control of plant pathogens. *Annu Rev Phytopathol* 31:53-80.

Datta MR, Palit C, Kumar M and Banerjee S. (2011). Plant growth promoting rhizobacteria enhance growth and yield of chilli (*Capicum annuum* L.) under field conditions. *Aust. J. Crop Sci.* 5P531-536.

FAO (2012). Food and Agricultural Organization of the United Nation, FAO Statistical Database, <http://faostat.fao.org>.

- Hervas A, Landa BB, Jimenez-Diaz RM (1997). Influence of chickpea genotype and *Bacillus* sp. On protectin of *Fusarium* wilt by seed treatment with nonpathogenic *Fusarium oxysporum*. Eur J Plant Pathol 103:631-642.
- Hervas A, Landa BB, Jimenez-Diaz RM (1997). Influence of chickpea genotype and *Bacillus* sp. On protectin of *Fusarium* wilt by seed treatment with nonpathogenic *Fusarium oxysporum*. Eur J Plant Pathol 103:631-642.
- ICARDA (2014). Improved chickpea strengthens food security in Sudan. Whats New at ICARDA International Center for Agricultural Research in the Dry Areas. Issue No.(7).
- Intana W, Yenjit P, Suwanno T, Sattasakulchai S, Suwanno M, Chamswarnng C (2008). Efficacy of antifungal metabolites *Bacillus* spp. For controlling tomato damping-off caused by *Pythium aphanidermatum*. Walailak Journal Science and Technology 5(1):29-38.
- Joseph B, and Patra RR. (2007). characterization of plant growth promoting rhizobacteria associated with chickpea (*Cicer arietinum* L.) Int.J. Pl. Prod. 1: 141-152.
- Juhnke ME, Mathre DE, Sands DC. (1987). Identification and characterisation of rhizosphere-competent bacteria of wheat. Appl Environ Microbiol 53:2793–2799.
- Karimi, K, Amini, J, Harighi, B. and Bahramnejad, B. (2012). Evaluation of biocontrol potential of *Pseudomonas* and *Bacillus* spp. against *Fusarium* wilt of chickpea. Australian Journal of Crop Science 6(4): 695-703. ISSN:1835-2707.
- Kloepper S, Ryu C and Zhang S. (2004) induce systemic resistance and promotion of plant growth of *Bacillus* spp., phytopathology, Vol. 94 (11): 1959-1266.
- Landa BB, Hervas A, Bettiol W, Jimenez-Diaz RM. (1997). Antagonistic activity of bacteria from the chickpea rhizosphere against *Fusarium oxysporum* f.sp. *ciceris*. Phytoparasitica, 25(4):305-318.
- Landa, BB, Navas-Cortes, J.A and Jimenez-Diaz, R.M. (2004). Integrated management of *Fusarium* wilt of chickpea with sowing date, host resistance and biological control. *Phytopathology*, Vol. 94, PP. 946-960.

- Navas-Cortes, J. A., Hau, B. and Jimenes-Dias, R. M. (1998). Effect of sowing date, host cultivar and race of *Fusarium oxysporum* f.sp *ciceris* on development of *Fusarium* wilt of chickpea. *Phytopathology*, 88:1338-1346.
- Nene YL, Reddy MV, Haware M.P, Ghanekar A.M and Amin K.S (1991). Field Diagnosis of Chickpea Diseases and their Control. ICRISAT. Information Bulletin no.28, International Crops Research Institute for the Semi-Arid Tropics. Patancheru, Andhra Pradesh 502 324, India.
- Nikam PS, Jagtap GP, Sontakke P.L (2011). Survey, Surveillance and Cultural Characteristics of Chickpea Wilt Caused by *Fusarium oxysporum* f.sp *ciceris* Afr.J.Agric.Res. 6(7):1913-1917.
- Schmidt CS, Agostini F, Leifert C, Killham K, Mullins CE (2004). Influence of soil temperature and matric potential on sugar beet seedling colonization and suppression of *Pythium* damping-off by the antagonistic bacteria *Pseudomonas fluorescens* and *Bacillus subtilis*. *Phytopathology*, 94:351-363.
- Schroth MN, Hancock JG. (1981). Selected topics in biological control. *Annu Rev Microbiol* 35:453-476, 1981.
- STATISTIX 8.0 Copyright (C) 1985-2003 Analytical Software. USA
- Van der Maesen, L.J.G. (1984). Availability and use of wild Cicer germplasm. *Plant genetic resources newsletter* (IBPGR/ FOW).
- Verma JP, Singh RP, Nayak ML. (1975). Laboratory evaluation of chemicals against *Xanthomonas malvacearum*, the incitant of bacterial blight of cotton. *Indian Phytopath* 28:170-174, 1975.
- Yun Cao, Zhenhua Zhang, Ning Ling, Yujuan Yuan (2011). *Bacillus subtilis* SQR 9 can control *Fusarium* wilt in cucumber by colonization plant roots. *Biol Fertil Soils* 47:495-506.
- Zaim S, Belaid L and Bellacene M (2013). Biocontrol of chickpea *Fusarium* wilt by *Bacillus* spp. *Rhizobacteria Journal of Plant Protection Research* vo.53, No2.

UNDER PEER REVIEW

