

---

## Another Look of Rings Domination in Ladder Graph

---

### Abstract

For a nontrivial connected graph  $G$  with no isolated vertex, a nonempty subset  $D \subseteq V(G)$  is a rings dominating set if each vertex  $v \in V - D$  is adjacent to at least two vertices in  $V - D$ . Thus, the dominating set  $D$  of  $V(G)$  is a rings dominating set if for all  $v \in V - D$ ,  $|N(v) \cap (V - D)| \geq 2$ . The cardinality of minimum rings dominating set of  $G$  is the rings domination number of  $G$ , denoted by  $\gamma_{ri}$  whereas the cardinality of maximum rings dominating set is the upper rings domination number and is denoted by  $\gamma'_{ri}$ . Here, we determine how the rings dominating set is constructed in the ladder graph with the inclusion of generated conditions for this type of domination and give new approach for its parameter.

*Keywords:* rings dominating set, minimum cardinality of rings dominating set, rings domination number, maximum cardinality of rings dominating set, upper rings domination number.

## 1 Introduction

Recently, a new parameter in domination of graphs was introduced called "rings domination" which has almost similar characteristics with the usual domination. As stated in [7], [10], [12], it has been a great area of study in graph theory to deal with, considering its applications in computer networks, electronics, and more, which inspires the creation of this paper. Prior to this, it intends to disclose the idea of rings dominating set specified over the ladder graph  $L_n$ . It is a graph generated from the cartesian product of paths of order  $n$  and of order two that induces a finite, planar, and undirected graph which has  $2n$  vertices and  $3n - 2$  edges.

Additionally, rings domination put a condition on the  $V - D$  set in the graph. Hence, we will be studying the behavior of rings dominating set in the ladder graph focusing on its  $V - D$  set. At the end of this paper, fundamental results are presented showcasing the conditions of rings dominating set and its largest cardinality for this particular graph.

---

## 2 Preliminary Notes

Some definitions of the concepts covered in this study are included below. You may refer on the remaining terms and definitions in [1], [4], [5], [8].

**Definition 2.1.** [5] (**Cartesian Product of Graphs**) The Cartesian product  $G$  of two graphs  $G_1$  and  $G_2$ , commonly denoted by  $G_1 \square G_2$  or  $G_1 \times G_2$  has vertex set  $V(G) = V(G_1) \times V(G_2)$ , where two distinct vertices  $(u, v)$  and  $(x, y)$  of  $G_1 \square G_2$  are adjacent if either (1)  $u = x$  and  $vy \in E(G_2)$  or (2)  $v = y$  and  $ux \in E(G_1)$ .




Figure 1: The cartesian product of graphs  $G_1$  and  $G_2$

**Definition 2.2.** [1] (**Rings Dominating Set, Rings Domination number**) A dominating set  $D \in V(G)$  is a rings dominating set if each vertex  $v \in V(G) - D$  is adjacent to at least two vertices in  $V(G) - D$ . The minimum cardinality among the rings dominating sets of  $G$  is called the rings domination number of  $G$  and is denoted by  $\gamma_{ri}(G)$ .




Figure 2: The graph of  $G$  and subgraph  $G_1$

**Example 2.1.** Consider the graph  $G$  in Figure 2. Obviously,  $D = \{v_2, v_4\}$  is a dominating set. Now see the subgraph  $G_1$  generated upon deleting vertices  $v_2$  and  $v_4$ . Observe that for every vertex  $v \in V(G) - D$  is adjacent to at least two vertices in  $V(G) - D$ , that is,  $deg(v_1) = deg(v_3) = 2$  and  $deg(v_5) = deg(v_6) = deg(v_7) = deg(v_8) = 3$ . Hence, the dominating set  $D$  is a rings dominating set. Also, note that  $D$  is the rings dominating set with the least cardinality, so  $\gamma_{ri}(G) = 2$ .

### 3 Main Results

In this section, the characteristics of rings dominating set in the ladder graph are presented. Also, the maximum cardinality of the rings dominating set for this particular graph is identified and new approach in determining the rings domination number is determined.

#### 3.1 Characteristics of Rings Dominating Set in the Ladder Graph, $L_n$

For convenience, we consider the path graph  $G$  of order  $n \geq 4$  with a vertex set  $V(G) = \{w_0, w_1, \dots, w_n\}$  and  $V(H) = \{u, v\}$ . The following shows the characteristics of rings dominating set in the ladder graph  $L_n = P_n \square P_2$ .

**Theorem 3.1.** *Let  $G = P_n$  where  $n \geq 4$  and  $H = P_2$ . Suppose  $D \subseteq V(P_n \square P_2)$ , then  $D = \cup(\{a\} \times T_a)$  where  $a \in S$  such that  $S \subseteq V(P_2)$  and  $T_a \subseteq V(P_n)$  is a rings dominating set of  $P_n \square P_2$  if and only if the following holds.*

1.  $S = P_2$ ; and
2.  $T_a$  is a restrained dominating set of  $P_n$ .

*Proof.* Suppose  $P_2 = \{u, v\}$  and  $P_n = \{w_1, w_2, \dots, w_n\}$  so that the vertex set of the ladder graph  $V(P_n \square P_2) = \{(u, w_1), (u, w_2), \dots, (u, w_n), (v, w_1), (v, w_2), \dots, (v, w_n)\}$ . Let  $\emptyset \neq D \subseteq V(P_n \square P_2)$  be a rings dominating set. Then  $D = \bigcup_{a \in S} (\{a\} \times T_a)$  where  $S \subseteq V(P_2)$  and  $T_a \subseteq V(P_n)$ .

Suppose  $S \neq V(P_2)$ , then  $|S| = 1$ . Without loss of generality, let  $S = \{u\}$ . Since  $D$  is a rings dominating set, then  $D = \bigcup_{u \in S} (\{u\} \times T_u) = \{(u, w_1), (u, w_2), \dots, (u, w_n)\}$ . Thus,  $V(P_n \square P_2) - D = \{(v, w_1), (v, w_2), \dots, (v, w_n)\}$ . Now,  $|N(v, w_1) \cap [V(P_n \square P_2) - D]| = |N(v, w_n) \cap [V(P_n \square P_2) - D]| = 1$ . A contradiction to the assumption that  $D$  is a rings dominating set. Hence,  $S = V(P_2)$ .

Further, assume  $T_a$  is not a restrained dominating set. It is equivalent to say that our assumption is  $T_a - V(P_n)$  has an isolated vertex or  $T_a$  is not a dominating set at the first place. Again, since  $D = \bigcup_{a \in S} (\{a\} \times T_a)$ , then it is a must that whenever  $(u, w_i)$  is in  $D$ , then  $(v, w_i)$  is also in  $D$  for some natural number  $i$ . Otherwise, we find a vertex  $(u, w)$  such that  $|N((u, w)) \cap [V(P_n \square P_2) - D]| \leq 1$ . When  $T_a - V(P_n)$  has at least one isolated vertex but  $T_a$  is a dominating set, then  $D$  is a dominating set with the induced subgraph  $(P_n \square P_2)[V(P_n \square P_2) - D]$  that has at least two corresponding vertices of degree 1, that is,  $|N((u, w)) \cap [V(P_n \square P_2) - D]| = |N((v, w)) \cap [V(P_n \square P_2) - D]| = 1$ . On the other hand, when  $T_a - V(P_n)$  has no isolated vertex but  $T_a$  is not a dominating set, then for every vertex  $(u, w) \in V(P_n \square P_2) - D$ ,  $deg((u, w)) \geq 2$ . Even so, a number of vertices of multiple of two is not dominated. In any case contradicts the assumption that  $D$  is a rings dominating set. Therefore  $T_a$  is a restrained dominating set.  $\square$

**Example 3.2.** *Let  $G = P_4$  and  $H = P_2$ . Then the ladder graph  $L_4$  is shown in Figure 3 below. Note that the vertex set of  $L_n$  is  $V(P_n \square P_2) = \{(u, w_1), (u, w_2), \dots, (u, w_n), (v, w_1), (v, w_2), \dots, (v, w_n)\}$ . Here, we let  $a = (u, w_1), b = (u, w_2), c = (u, w_3), d = (u, w_4), e = (v, w_1), f = (v, w_2), g = (v, w_3), h = (v, w_4)$ . Clearly,  $S = P_2$  as we have taken vertices containing both  $u, v \in P_2$ . Now, consider the subgraphs, say  $S_k$  for  $k = 1, 2$  shown in Figure 4. For each  $T_{a_i}$  where  $i = 1, 2$  is a restrained dominating set since there exists no isolated vertex in  $L_n[V(S_k) - T_{a_i}]$ . Lastly, note  $a, e \in D$  are corresponding vertices in  $L_4$ . Similarly,  $d, g \in D$  are corresponding vertices. Hence (i), (ii), and (iii) are satisfied.*




Figure 3: The rings dominating set of  $L_4$




Figure 4: The subgraphs  $S_1$  and  $S_2$  of  $L_4$

### 3.2 Rings Domination Number of a Ladder graph via Congruence Modulo

Here, we introduce a different way of finding the rings domination number of a ladder graph via congruence modulo. Consequently, this will serve as another convenient way of determining the rings domination number of a ladder graph of a specific order which is divided into three different cases shown below.

**Theorem 3.3.** *Let  $L_n = P_n \square P_2$  be a ladder graph of order  $2n$ . For  $n \geq 4$ , the rings domination number of the ladder graph is*

$$\gamma_{ri}(L_n) = \begin{cases} 2\left(\frac{n+6}{3}\right), & \text{if } n \equiv 0 \pmod{3} \\ 2\left(\frac{n+2}{3}\right), & \text{if } n \equiv 1 \pmod{3} \\ 2\left(\frac{n+4}{3}\right), & \text{if } n \equiv 2 \pmod{3}. \end{cases}$$

*Proof.* Let  $L_n = P_n \square P_2$  be a ladder graph of order  $2n$  such that  $P_n = \{1, 2, 3, \dots, n\}$  and  $P_2 = \{a, b\}$ . Suppose  $S$  is a rings dominating set of minimum cardinality, that is,  $S$  is a  $\gamma_{ri}$ -set, then we consider the following cases.

Case 1:  $n \equiv 0 \pmod{3}$ .

Let  $R \subseteq V(L_n)$  with  $R = \{(a, 1), (a, 4), (a, 7), (a, 8), (a, 9), (a, 12), (a, 15), \dots, (a, n-3), (a, n)\} \cup \{(b, 1), (b, 4), (b, 7), (b, 8), (b, 9), (b, 12), (b, 15), \dots, (b, n-3), (b, n)\}$ . Clearly,  $R$  is a rings dominating set and  $R = V(P_2 \times P)$ , where  $P \subseteq V(P_n)$  and  $P = \{1, 4, 7, 8, 9, 12, 15, \dots, n-3, n\}$ . For this circumstance,  $|R| = 2\binom{n+6}{3}$ . Since  $S$  is a  $\gamma_{ri}$ -set,  $|R| \geq |S|$ . Hence  $|S| \leq |R| = 2\binom{n+6}{3}$ . On the other hand, since  $S$  is a  $\gamma_{ri}$ -set of  $L_n$  then  $S$  must have at least  $2\binom{n+6}{3}$  number of vertices in  $L_n$ . Hence,  $|S| \geq 2\binom{n+6}{3}$ . Therefore,  $|S| = 2\binom{n+6}{3}$ .

Case 2:  $n \equiv 1 \pmod{3}$ .

On similar manner in case 1, let  $K \subseteq V(L_n)$  with  $K = \{(a, 1), (a, 4), (a, 7), \dots, (a, n-3), (a, n)\} \cup \{(b, 1), (b, 4), (b, 7), \dots, (b, n-3), (b, n)\}$ . It can easily be seen that  $R$  is a rings dominating set and  $K = V(P_2 \times P)$ , where  $P \subseteq V(P_n)$  and  $P = \{1, 4, 7, \dots, n-3, n\}$ . With this,  $|K| = 2\binom{n+2}{3}$ . Since  $S$  is a  $\gamma_{ri}$ -set,  $|K| \geq |S|$ . Hence  $|S| \leq |K| = 2\binom{n+2}{3}$ . Meanwhile, since  $S$  is a  $\gamma_{ri}$ -set of  $L_n$ , then  $S$  must have at least  $2\binom{n+2}{3}$  number of vertices in  $L_n$ . Hence,  $|S| \geq 2\binom{n+2}{3}$ . Thus,  $|S| = 2\binom{n+2}{3}$ .

Case 3:  $n \equiv 2 \pmod{3}$ .

Comparably, let  $T \subseteq V(L_n)$  with  $T = \{(a, 1), (a, 4), (a, 7), (a, 8), (a, 11), (a, 14), \dots, (a, n-3), (a, n)\} \cup \{(b, 1), (b, 4), (b, 7), (b, 8), (b, 11), (b, 14), \dots, (b, n-3), (b, n)\}$ . Obviously,  $T$  is a rings dominating set and  $R = V(P_2 \times P)$ , where  $P \subseteq V(P_n)$  and  $P = \{1, 4, 7, 8, 11, 14, \dots, n-3, n\}$ . On this account,  $|T| = 2\binom{n+4}{3}$ . Since  $S$  is a  $\gamma_{ri}$ -set,  $|T| \geq |S|$ . Hence  $|S| \leq |T| = 2\binom{n+4}{3}$ . On the other hand, since  $S$  is a  $\gamma_{ri}$ -set of  $L_n$  then  $S$  must have at least  $2\binom{n+4}{3}$  number of vertices in  $L_n$ . Hence,  $|S| \geq 2\binom{n+4}{3}$ . Therefore,  $|S| = 2\binom{n+4}{3}$ .  $\square$

### 3.3 Maximum Cardinality of Rings Dominating Set in the Ladder Graph

Here, the maximum cardinality of rings dominating set is referred to as the upper rings domination number, denoted by  $\gamma'_{ri}$ .

**Theorem 3.4.** *Let  $D \subset L_n$ . The upper rings domination number in the ladder graph  $L_n$  is  $\gamma'_{ri} = 2n - 4$ .*

*Proof.* Let  $L_n$  be a ladder graph with order  $2n$ . Assume that the maximum cardinality of rings dominating set in the ladder graph is  $2n - 3$ . From the proof of Theorem 3.1, for every vertex  $(u, w_i) \in D$ ,  $(v, w_i)$  must also be in  $D$ . Hence the cardinality of rings dominating set in the ladder graph is an even number. Note that  $2n - 3$  is an odd number for every natural number  $n$ . Thus  $2n - 3$  is not the maximum cardinality of rings dominating set in the ladder graph  $L_n$ . To this end,  $2n - 2$  is not the maximum cardinality since it will lead to an induced subgraph  $L_n[V(L_n) - D]$  of order 2 that could either be a path or an empty graph, both of order 2.  $\square$

**Example 3.5.** *Consider the ladder graph  $L_7$  in Figure 5 below. The possible rings dominating set of different cardinality for  $L_7$  was provided. Based on the diagram, the maximum cardinality of the rings dominating set in the ladder graph  $L_7$  is  $2n - 4 = 2(7) - 4 = 10$ .*

*Remark 3.1.* Suppose  $D \subseteq L_n$ . The rings dominating set of maximum cardinality is  $L_n$  itself. Hence,  $\gamma'_{ri} = |L_n| = 2n$ .

Note that if  $\gamma'_{ri}(D) = 2n$ , then  $D = V(L_n)$ . Consequently,  $V(L_n) - D = \emptyset$ . Hence, the condition of rings dominating set is not violated since there exists no vertex  $v \in V(L_n) - D$  such that  $\deg(v) < 2$ . In fact, no vertex in  $V - D$  that should be considered and examined.




Figure 5: All possible rings dominating set of  $L_7$  of different cardinality

**Proposition 3.1.** *Let  $D$  be a rings dominating set in the ladder graph  $L_n$ . Then the order of the induced subgraph  $L_n[V(L_n) - D]$  is divisible by 4. Further, the order and size of  $L_n[V(L_n) - D]$  are equal.*

*Proof.* This is immediate from Theorem 3.1. □

**Example 3.6.** *Again, refer to Figure 5. It can easily be seen that the order  $L_7[V(L_7) - D_1]$  and  $L_7[V(L_7) - D_2]$  are  $8 = 4(2)$  and  $4 = 4(1)$  respectively, which coincides to their sizes.*

## 4 Conclusion

In this article, rings dominating sets in the ladder graph generated on the cartesian product of path graph of order  $n$  and of order 2 are studied. Further, the maximum cardinality of rings dominating set is also determined and new approach for the rings domination number is presented. Lastly, we intend to examine the rings dominating set and rings domination number for few unstudied graph families in the future.

---

## References

- [1] Abed, S. S., AL-Harere, M.N. (2022). *Rings domination in graphs*. International Journal of Nonlinear Analysis and Applications.
- [2] Cabahug Jr, I. S., Canoy Jr, S. R. (2016). *Independent  $d$ -Power Dominating Sets in Graphs*. Applied Mathematical Sciences, 10(8), 377-387.
- [3] Cabahug Jr, I. S., Canoy Jr, S. R. (2018). *Connected  $d$ -Power Dominating Sets in Graphs*. Advances and Applications in Discrete Mathematics, 19(3), 171-182.
- [4] Canoy Jr., S. R. (2014). *Restrained Domination in Graphs Under Some Binary Operations*. Applied Mathematical Sciences, 8(16), 6025 - 6031
- [5] Chartrand, G., Lesniak, L., Zhang, P., (2015). *Graphs and Digraphs* (Discrete Mathematics and Its Applications) (6th ed.), Chapman and Hall/CRC.
- [6] Consistente, L. F., Cabahug Jr, I. S. *Hinge Total Domination on Some Graph Families*.
- [7] Gayathri, A., Muneera, A., Rao, T.N., Rao, T.S. *Study of Various Dominations in Graph Theory and Its Applications*. International Journal of Scientific and Technology Research, 9, 3426 - 3429.
- [8] Guichard, D., (2017). *An introduction to combinatorics and graph theory*. Hitman College - Creative Commons
- [9] Mangubat, D. P., Cabahug Jr, I. S. *On the Restrained Cost Effective Sets of Some Special Classes of Graphs*.
- [10] Moussa M.I., Badr E.M. *Ladder and subdivision of ladder graphs with pendant edges are odd graceful*.
- [11] Tan, K. S. R., Cabahug Jr, I. S. *Safe Sets in Some Graph Families*.
- [12] Venkateswari, R. *Applications of Domination Graphs in Real Life*. A Journal of Composition Theory.