

## Original Research Article

# Lipid peroxidation status in embryo culture media: *The impact on fertilization and embryo quality during IVF cycles*

### Abstract

**Background:** The process of preimplantation embryo development in vitro represents a key phase during in vitro fertilization (IVF) cycles. It involves several regulatory signaling pathways as well as an optimized in vitro culture system. The resulting embryo quality helps to determine embryo competence prior to implantation and pregnancy outcomes. Reactive oxygen species (ROS) are known to play a major role in influencing the process of embryo development. Their role can be reflected in the regulation of signaling pathways as second messengers or in the irreversible cell alterations due to oxidative stress following an excess of ROS levels.

**Methods:** In this study, we investigated the association between morphological embryo quality (fertilization, cleavage quality and fragmentation levels) and lipid peroxidation levels (Malondialdehyde) in embryo culture media. After intracytoplasmic sperm injection (ICSI), total of 103 oocytes were evaluated on day 1 and day 3 of their development and their corresponding culture media were analyzed by estimating MDA levels using thiobarbituric acid.

**Results:** The results showed no significant association between MDA levels in culture media and fertilization rate ( $p=0.3$ ), sperm quality ( $p=0.99$ ;  $p=0.17$ ;  $p=0.46$ ;  $p=0.30$ ;  $p=0.65$ ;  $p=0.44$ ;  $p=0.09$ ;  $p=0.15$ ;  $p=0.56$ ), embryo fragmentation levels ( $p=0.79$ ;  $p=0.40$ ), AMH levels ( $p=0.31$ ;  $p=0.36$ ) and female age ( $p=0.60$ ;  $p=0.34$ ). However, we revealed a significant association between cleavage quality and MDA levels in the embryo environment ( $p=0.03$ ).

**Conclusion:** We conclude that oxidative stress in IVF culture media might be mainly associated with delayed embryonic development

**Keywords:** Embryo development, reactive oxygen species, malondialdehyde, oxidative stress

## INTRODUCTION

Embryo quality is one of the most important factors that are assessed during the process of in vitro fertilization (IVF) in order to increase implantation and pregnancy rates. The evaluation of embryo quality in IVF laboratory mostly involves morphological features such as cleavage kinetics and fragmentation rates and this process involves a diversity of connected signalization and regulation pathways. Whether occurring *in vivo* or *in vitro*, embryonic development requires several factors that maintain the homeostasis of these signaling pathways at each stage of the preimplantation process. Among the regulatory factors that have attracted the most interest due to their complexity are reactive oxygen species (ROS). These highly reactive molecules are generated from oxygen reduction by electrons during cellular metabolism and act as second messengers in physiological cell signaling and control pathways [1,2,3,4]. Embryos are known for three free radicals during their normal aerobic metabolism: hydrogen peroxide ( $H_2O_2$ ), superoxide anion ( $O_2^-$ ) and hydroxyl radical ( $OH^-$ ) [5, 6,1,7]. However, free radicals are more often associated with the principle of oxidative stress, which results from an imbalance in the level of oxidants and antioxidants, in favor of oxidants during cell metabolism [8,9,10,1]. Many studies reported a lot of interesting results about ROS being involved in delayed embryo development [12,13,14,15], playing an essential role in the regulation of some of the most important stages of embryo development such as fertilization, cleavage and embryo hatching [16,15,17,18] or having no association with embryo quality at all [19]. These conflicting findings make ROS involvement in cell signalizations complex and simple at the same time. In fact, it has been clearly showed that excess ROS has a negative impact on embryo quality in vitro [20,15,21,22, 23], as we know that any type of excess in our metabolism represents loss of balance. However, knowing that ROS is implicated in so many regulation pathways that are necessary for embryo quality, it is important to understand to what extent ROS are necessary and impactful during the early stages of embryo development in vitro and whether or not they are impactful.

The aim of this study is to give a better understanding of the dynamics of ROS production during in vitro culture of pre implantation embryos after intra cytoplasmic sperm injection (ICSI), and to explore the relationships between ROS levels in culture media (CM) and embryo quality.

## **MATERIEL AND METHODS**

### **Patients**

A total of 103 women aged between 24 and 50 years old, with primary and secondary infertility and undergoing oocyte retrieval cycles of assisted reproduction, between January 30, 2019 and March 03, 2020 were prospectively included. Couples were not selected for age, sperm parameters or causes of infertility (Table 1). This work did not involve experiments on humans or animals, 207 spent CM were collected.

### **Ovarian stimulation**

Women underwent controlled ovarian stimulation with the flexible gonadotrophin-releasing hormone (GnRH) antagonist protocol. A daily subcutaneous injection of recombinant follicle stimulating hormone (rFSH; Gonal-F, Merck-Serono) was used alone or in combination with human menopausal gonadotrophin (HMG, Menopur; Ferring). The FSH dose was based on the women's age, AMH concentration in addition to prior history of ovarian stimulation and was adjusted according to usual parameters of follicle growth, determined by serum estradiol (E2) concentration and ultra sound monitoring.

A daily dose of GnRH antagonist (Cetrotide, Merck-Serono or Orgalutran, MSD) was injected subcutaneously, starting from day 6 of FSH administration. The ovulation trigger was performed with 10 000 IU of human chorionic gonadotrophin (rHCG, Ovitrelle; Merck-Serono) and gonadotrophin releasing hormone (Decapeptyl, Ferring), after obtaining follicles that reached dimensions of 17mm or greater in diameter and adequate serum E2 levels. Oocytes were retrieved 34-36 hours after hCG administration.

### **Oocyte and sperm preparation**

The retrieved oocytes were isolated from follicular fluid, rinsed and cultured in CM (SAGE 1-Step, Origio). 2-3 hours after retrieval, the oocyte-corona-cumulus complexes were placed in a HEPES-buffered medium (Ferticult Flushing medium, Fertipro) containing hyaluronidase (Hyaluronidases in Ferticult Flushing medium, 80IU/mL, Fertipro) and were mechanically decoronated using a 20-200 $\mu$ L micropipette. The nuclear maturation grades were classified as metaphase II or non-metaphase II (Metaphase I or Prophase I) oocytes.

Sperm samples were collected from the male partner by masturbation in a sterile container, after 2-3 days of abstinence. At first, semen samples were evaluated for spermiatic parameters (concentration, motility and morphology) based on WHO (2010) recommendations, to be divided into several groups. For each parameter, sperm samples were divided into two groups. Sperm concentration included samples with normal concentration ( $> 15$  M/mL) and samples with oligospermia ( $\leq 15$  M/mL). The category of motility rate included samples with normal sperm motility rate ( $>32\%$ ) and samples with asthenospermia ( $\leq 32\%$ ). The category of morphology rate included normal sperm morphology rate ( $>20\%$ ) and teratospermia ( $\leq 20\%$ ). Motile spermatozoa were then selected using a discontinuous two-layer density gradient technique (Puresperm 80/40; SAGE) as described by Aboulmaouahib et al. (2016).

All matured oocytes underwent ICSI after decoronisation. One micro-injected oocyte per patient was then randomly selected and placed in an oil-covered single drop of 100  $\mu$ L of culture media (SAGE 1-Step, Origio), in a petri dish.

The medium was renewed on day 1 (18-20h after ICSI) and day 3 (68-70h after ICSI).

### **Assessment of fertilization and embryo quality**

A total of 103 oocytes were evaluated.

On day 1, fertilization was confirmed by the appearance of a double polar body and/or two pronuclei. At that stage, oocytes were divided into two groups: zygotes and unfertilized oocytes. On day 3, the embryonic quality was evaluated according to the number of blastomeres and the presence or absence of cell fragmentation. Initially, embryos were divided into 2 groups: 7-8 cells embryos and 4-6 cells embryos. Secondly, embryos were divided into 2 groups: fragments-free embryos and embryos with 5-10% fragmentation.

The temperature inside the incubators (IVF-Cube AD3100, ASTEC; Thermo Scientific HeraCell 150) was controlled by a certified thermometer and remained at  $37 \pm 0.2$  °C. Oxygen level inside the incubators was at 5% and the cultivating medium pH at  $7.3 \pm 0.02$  with CO<sub>2</sub> around 5.6%.

### **Malondialdehyde (MDA) measurement**

The spent CM of the corresponding oocytes and embryos (day 1, day 3) were collected for the determination of lipid peroxidation levels.

Microvolume protein concentration was measured using the Nanodrop spectrophotometer as described by Desjardins and colleagues (24). Lipid peroxidation was estimated by thiobarbituric acid reactive substances (TBARS) reaction with malondialdehyde (MDA) as described by Sanokyszyn and Marnett (1990). Briefly, 100  $\mu$ L of culture media was added to 10% trichloroacetic (TCA) and 0.67% thiobarbituric acid (TBA). The absorbance was measured at 532nm and MDA levels were expressed in  $\mu$ mole/ mg of proteins.

We are well aware that MDA is not the only compound that can react with thiobarbituric acid to produce colored adducts, which can represent a limitation to our findings. However, such a limitation in such a complex biological sample can show our results as a door opening to others research using more specific techniques.

### Statistical analysis

The results are expressed as the mean  $\pm$ Standard deviation or percentage of total. Data was obtained with the student's t-test using SPSS (Statistical Package for the Social Science). Statistical significance was defined as  $p < 0.05$ .

No. of injected oocytes	103
No. of cycles with no embryos available	18
Age of female partner (years)	37.4 $\pm$ 5.4 (25-46)
Age of male partner (years)	42.1 $\pm$ 7.2 (32-58)
Primary infertility	92
Secondary infertility	11
Duration of infertility (years)	7.18 $\pm$ 5.07 (1-20)
Male infertility	29
Endometriosis	9
OPK	5
Tubal infertility	4
Ovarian insufficiency	4
Combined factors	10
Idiopathic	42

Values are mean  $\pm$  SD

**Table 1: Demographic characteristics of ICSI cycles (103)**

## RESULTS

### Relationship between fertilization and MDA levels in culture media

As shown in table 2, on day 1, MDA levels were significantly higher in CM of embryos than the control samples (no oocytes or embryos in CM) ( $p < 0.0001$ ). However, MDA levels in CM were not associated with fertilization. In fact, there was no significant difference between MDA levels in CM of zygotes and unfertilized oocytes.

### **Relationship between MDA levels in culture media and embryo quality**

All embryos were divided into 2 groups depending on the embryo's cell number (7-8 / 4-6 cells) and fragmentation (with or without fragmentation). On day 1, we noted that MDA levels in CM of 4-6 cells embryos were significantly higher ( $p = 0.03$ ) than in CM of 7-8 cells embryos. While on day 3, there was no significant difference between MDA levels in CM of these two groups.

While comparing fluctuations of MDA levels in CM within the same group, we noted that MDA levels in CM of 4-6 cells embryos were significantly higher on day 1 than on day 3 ( $p = 0.02$ ). Contrariwise, there was no significant difference between MDA levels in day 1 and day 3 CM of 7-8 cells embryos (Table 2). Likewise, there was no significant difference between MDA levels in CM of fragmented (5-10% fragmentation) embryos and fragments-free embryos, on day 1 as well as on day 3. Also, in control samples, MDA levels significantly increased from day 1 to day 3 (Table 2).

### **Relationships between cleavage quality in day 3 embryos and fragmentation**

The group of fragments-free embryos included 42/ 53 (79.2 %) 7-8 cells embryos and 11/53 (20.7%) 4-6 cells embryos. The group of 5-10% embryo fragmentation included 16/27 (59.2%) 7-8 cells embryos and 11/27 (40.7%) 4-6 cells embryos (Figure 1).

### **Comparison between MDA levels in culture media, embryo quality and sperm quality**

In our work we have sought to link the results of MDA levels in CM of embryo quality with sperm quality. The latter was measured by concentration, mobility and morphology parameters of each patient. On day 1, there was no significant difference between MDA levels

in CM that corresponds to normal sperm concentration and oligospermia, for 7-8 cells embryos or 4-6 cells embryos , either on day 1 or on day 3. The same results were found in MDA levels that corresponds to normal sperm motility parameters / asthenospermia and normal sperm morphology parameters and teratospermia (Table 3).

### **Relationships between cleavage quality in day 3 embryos and sperm parameters**

In addition, we have sought to establish relationships between cleavage quality in day 3 embryos and sperm quality in terms of numeration, motility and morphology. In the normal sperm morphology parameters group, 78.8% (41/52) of the embryos were 7-8 cells embryos and 21.1% (11/52) were 4-6 cells embryos. In the teratospermia group, 55.8% (19/34) of embryos were 7-8 cells embryos and 44.1% (15/34) were 4-6 cells embryos. In the normal sperm concentration group, 78.2% (36/46) of embryos were 7-8 cells embryos and 21.7% (10/46) were 4-6 cells embryos. In the oligospermia group, 60% (24/40) of embryos were 7-8 cells embryos and 40% (16/40) were 4-6 cells embryos. In the normal sperm motility parameters group, 79.1% (19/24) of embryos were 7-8 cells embryos and 20.8% ( 5/24) were 4-6 cells embryos. In the asthenospermia group, 67.7% (40/59) of embryos were 7-8 cells embryos and 32.2% (19/59) were 4-6 cells embryos (Table 4).

### **Relationships between MDA levels in CM and AMH levels, women's age and type of infertility**

Furthermore no significant difference was noted between MDA levels in CM on day 1 or day3 and the women's age. In addition, no association was found between MDA levels in CM on day1/ day 3 and AMH levels as well as the type of infertility (Table 5).

	Control samples (n=36)	Embryo cultures (n=81)	Unfertilized oocytes (n=18)	7-8 cells embryos (n=42)	4-6 cells embryos (n=22)	Fragments free (n=53)	5-10% fragments (n=29)
<b>Day 1</b>	0.0001 ± 0.0002	0.0020 ± 0.0033	0.0035 ± 0.0061	0.0015 ± 0.0033	0.0035 ± 0.0036	0.0021 ±0.0036	0.0019 ±0.0025

<b>p value</b>		1.03E <sup>-06*</sup>	0.03*	0,03		0.79	
		0.3					
<b>Day 3</b>	0.0007 ± 0.0007	0.0014 ± 0.0020	0.0020 ± 0.0038	0.0013 ± 0.0020	0.0017 ± 0.0017	0.0014 ± 0.0020	0.0018 ± 0.0024
<b>p value</b>		0.009*	0.8*	0.4		0.4	
<b>P value</b>	2.35E <sup>-05**</sup>			0.58**	0.02**	0.20**	0.80**

**Table 2: Mean Values, Standard Deviations, and Significant Differences of MDA levels (µg/mg of proteins) at day 1 and day 3 in culture media (CM) of Zygotes, 4-6 cells embryos, 7-8 cells embryos, embryo fragmentation and control samples (no oocytes or embryos in CM)**

\*Comparison with control

\*\* Comparison between Days1 and 3

The results were evaluated by a nonparametric test: the Mann-Witney U for the comparison of the groups.

	<b>Normal sperm concentration parameters</b>	<b>Oligospermia</b>	<b>Normal sperm motility parameters</b>	<b>Asthenospermia</b>	<b>Normal sperm morphology parameters</b>	<b>Teratospermia</b>
<b>7-8 cells / Day 1</b>	0.0015±0.00025 (n=36)	0.0015±0.0036 (n= 24)	0.0019±0.0031 (n= 19)	0.0013 ± 0.0030 (n = 40)	0.0015±0.0025 (n= 41)	0.0006 ±0.0014 (n= 19)
<b>p value</b>	0.99		0.46		0.09	
<b>7-8 cells / Day 3</b>	0.0011±0.0016 (n=36)	0.0016±0.0026 (n=24)	0.0015±0.0020 (n=19)	0.0012 ± 0.0021 (n=40)	0.0014±0.0021 (n=41)	0.0014± 0.0021 (n=19)
<b>p value</b>	0.37		0.65		0.99	
<b>4-6 cells/ Day 1</b>	0.0045±0.0037 (n=10)	0.0024±0.0034 (n=16)	0.0052 ±0.0034 (n=5)	0.0032±0.0036 (n=19)	0.0046±0.0038 (n=11)	0.0025±0.0030 (n=15)
<b>p value</b>	0.17		0.30		0.15	
<b>4-6 cells/ Day 3</b>	0.0029± 0.0028 (n=10)	0.0013±0.0017 (n=16)	0.0032±0.0036 (n=5)	0.0018± 0.0019 (n=19)	0.0024±0.0028 (n=11)	0.0018±0.0018 (n=15)
<b>p value</b>	0.13		0.44		0.56	

**Table 3: MDA levels (µg/mg of proteins) in CM of 7-8 / 4-6 cells embryos, on day 1/ day 3, according to sperm quality**


	7-8 cells embryos	4-6 cells embryos
--	-------------------	-------------------

Normal sperm morphology parameters	78.8% (41/52)	21.1% (11/52)
Teratospermia	55.8% (19/34)	44.1% (15/34)
Normal sperm numeration parameters	78.2% (36/46)	21.7% (10/46)
Oligospermia	60% (24/40)	40% (16/40)
Normal sperm motility parameters	79.1% (19/24)	20.8% (5/24)
Asthenospermia	67.7% (40/59)	32.2% (19/59)

**Table 4: Percentage of 7-8 cells embryos and 4-6 cells embryos in each group of sperm parameters.**

	Female age (years)		AMH levels		Type of infertility	
	<38 (n=50)	> 38 (n=51)	<1 (n=42)	>1 (n=44)	Male infertility (n=29)	Idiopathic infertility (n=45)
<b>Jour 1</b>	0.0022 ± 0.0035	0.0026 ± 0.0046	0.0026 ± 0.0045	0.0018 ± 0.0032	0.0029 ± 0.0050	0.0019 ± 0.0038
<b>p value</b>	0.60		0.31		0.38	
<b>Jour 3</b>	0.0014 ± 0.0024	0.0019 ± 0.0026	0.0019 ± 0.0023	0.0014 ± 0.0029	0.0025 ± 0.0037	0.0011 ± 0.0013
<b>p value</b>	0.34		0.36		0.06	

**Table 5: Comparison of MDA levels (µg/mg of proteins) of embryo CM according to female age, AMH levels and type of infertility**



**Figure 1: Associations between cleavage quality and fragmentation in embryos on day 3**

## DISCUSSION

Based on our previous readings [25], research has demonstrated the key role of ROS in the regulation of calcium oscillations for the process of oocyte activation by the sperm; however the negative impact of oxidative stress on oocyte fertilization *in vitro* is a very well described outcome in assisted reproductive technology. In a balanced *in vivo* environment, mitochondrial ROS, such as H<sub>2</sub>O<sub>2</sub>, O<sub>2</sub><sup>-</sup> and NO<sup>-</sup> act as second messengers to release internal calcium reserves by acting on the calcium pumps of the endoplasmic reticulum (ER) and the plasma membranes [25,26,27,28,24,29]. During the process of fertilization, as calcium is a key factor of the quality of fertilization, an amount of ROS levels might be also necessary for the regulation of this process. In the other hand, excess intra/ extracellular ROS levels are also known to induce high lipid peroxidation which can disturb calcium oscillations and alter oocyte fertilization [30,31,17,22]. In fact, ROS levels in ICSI and IVF culture media on day 1 have been previously shown to be associated with low fertilization outcomes [33]. All these cellular events show us above all the importance of maintaining extra/ intracellular homeostasis of oocytes and embryos either *in vivo* or *in vitro*.

On day 1, our data showed that MDA levels in CM of fertilized oocytes were significantly higher than MDA levels in the control samples (no oocytes/ embryos in the media). We can speculate that this elevation may be due to a possible production of ROS by oocytes in parallel with the triggering of Ca<sup>++</sup> pathways during fertilization, however it cannot be excluded that it can simply be caused by a production of ROS by culture media components overtime [6,33,34]. The radical « singlet oxygen » that are present in the atmosphere can also attack lipidated components of CM and thus produce MDA.

Interestingly, MDA levels in control samples significantly increases from day 1 to day 3. This finding could confirm that *in vitro* culture conditions alone can cause moderate ROS production and represent a factor that is associated with oxidative stress in embryo culture media.

In this study, since all oocytes have been fertilized by ICSI procedure, the overproduction of ROS caused by sperm in CM during conventional IVF was widely mitigated [35,36,37] which made MDA levels in CM on day 1 appear to be more associated with the fertilization process. While comparing between zygotes and unfertilized oocytes on day1, we found no association between MDA levels in CM and our fertilization rate. A similar conclusion was reached by a recent study where ROS levels in CM were not associated with fertilization rate as fertilized,

polyspermic and unfertilized zygotes showed similar ROS levels [19]. Since zygotes generates different outcomes of embryo quality on day 3, we are tempted to speculate that ROS levels in CM are not a major indicator of fertilization success but some ROS deletions might appear later during the next phases of early development, such as cleavage and compaction. Successful oocyte fertilization (zygotes) on day 1 does not always result in successful embryo development on day 3 [19,38,39] and early cleavage stage embryos are more sensitive to oxidative stress during the early stages of development (40,41,42,43), thereby, we were interested to study the variations of MDA levels in CM of embryos according to the number of cells and fragmentation on day 3. On day 1, the oocyte is fertilized and ready to initiate cleavage [44,45,46]. At this stage, our results showed that 4-6 cells embryos produced higher MDA levels in CM than 7-8 cells embryos, which can reveal a possible association between cleavage quality and oxidative stress levels in the embryo environment. Overall, this finding is in consistence with previous research showing that oxidative stress can be involved in serious embryo damage such as DNA fragmentation and impairment of mitochondrial and enzymes function which can result in delays of embryonic cleavage or embryo arrest [12,32,47,13,20, 14,15,22,23]. By day 3, there were no significant difference between MDA levels of both 4-6 and 7-8 cells embryos which is in concordance with Lan and colleagues [19] who found that ROS levels on day 3 were not significantly associated with cleavage stage and embryo quality. These results may address the fact that by the end of cleavage, embryos might be able to use their antioxidant system to bring ROS levels down to a level that is low enough to trigger the process of compaction, regardless of cleavage quality. In fact, while some anti oxidants are known to be absent in mature oocytes, such as catalase [42] embryos acquire their antioxidant system once genome activation occurs after the 2 cell stage. Embryos are then able to recover their antioxidant concentrations and use it as an adaptative protection against oxidative stress, like shown in mouse embryos [6,49–52]. In CM of 4-6 cells embryos, we observed a fluctuation of MDA levels between day 1 and day 3, as the levels were higher on day 1 and significantly decreased by day 3. At first, this finding made us suspect that these fluctuations might be due to the initiation of cleavage on day 1, as the need of an amount of ROS by the cells during cleavage for regulation has been largely discussed previously. Several studies have highlighted the complex relationship between intracellular ROS levels and preimplantation embryo development. H<sub>2</sub>O<sub>2</sub> has been shown to increase during the first cell divisions, the 8-cell stage to increase blastulation capacity and hatching in mouse and bovine embryos [25,53–56]. However in CM of 7-8 cells embryos, no significant difference in MDA levels was found

between day 1 and day 3. These results suggest that fluctuations of oxidative stress in CM are more likely to be associated with delay in embryo development than with a positive regulation of the cleavage process. The redox state in CM as in embryos needs to stay stable from day 1 to the end of in vitro culture for a better embryo development. Nevertheless, we are aware of the limitations of our study as there is no following up of embryo development and MDA levels to day 5.

Other reports on the impact of ROS in embryo quality implied that an increase in the rate of ROS in ICSI cycles is associated with an increase in the rate of embryo fragmentation (>10%) [33,57–61]. However, our study showed no association between MDA levels in the embryo surrounding and embryo fragmentation. Embryo fragmentation is defined by the appearance of anucleated fragments in the cytoplasm during the cleavage stage [61,62,59]. Although it is a common feature of IVF derived embryos [64–67], mechanisms and causes of embryo fragmentation are still unclear. Many investigators came up with several hypotheses as some reported fragmentation to occur during the first cell division [65,68,69] and result in the loss of important volumes of blastomere's cytoplasm, reduction of embryo size and depletion of essential organelles, namely mitochondria, mRNA and proteins [70,71]. Plus, due to blastomere's movement during embryo cleavage, fragments in the cleavage cavity may disturb the cell-cell contact and distortion of division planes [21,64,65]. In fact, our data also showed that 42/ 53 (79,2%) fragment free embryos had 7-8 cells on day 3, which may suggest that fragments free embryos have more healthy mitochondria leading to a healthier cleavage, but since we found no significant difference between the rate of 7-8 (16/27; 59.2%) and 4-6 cells (11/27; 40.7%) embryos in the group of fragmented embryos, our results stay controversial. Therefore, we can only deduce that the quality of embryo cleavage can be associated to fragmentation rate and to ROS production; however there is no significant association between these two factors.

In order to deepen our research on the impact of oxidative stress levels in embryo surroundings, we also investigated on the possible relationships between sperm quality, embryo quality and MDA levels in CM. No significant difference was found in MDA levels between CM of embryos corresponding to normal sperm concentration and oligospermia, and also between CM of embryos corresponding to normal sperm motility rate and asthenospermia, whether it's 7-8 cells or 4-6 cells embryos. This data can be primarily explained by the fact that low sperm concentration and motility factors as well as excess ROS production by conventional IVF are attenuated by ICSI procedures [72,73]. Also, these

finding can demonstrate that sperm quality does not impact MDA levels in CM and that sperm quality is more related to embryonic quality than directly to oxidative stress in CM. In fact, our statistics showed that 7-8 cell embryos mainly originates from sperm with normal parameters.

It was also interesting to notice that while in the group of normal sperm morphology parameters, 68.3% (41/52) were 7-8 cells embryos and only 21,1% (11/52) were 4-6 cells embryos, there were no significant difference between the rate of 7-8 cells embryos (55,8%) and 4-6 cells embryos (44,1%) in the teratospermia group. During ICSI, morphology selection of spermatozoa is based on its normal looking shape under x200 or 400 magnification, which might miss the detection of minor head malformations. Plus, the risk of using spermatozoa with damaged DNA in case of teratospermia is greater [74–81]. In order to consolidate our study, we also investigated the possible relationships between MDA levels and AMH, women's age and some type of infertilities (Male infertilities, idiopathic). AMH is produced by granulosa cells from pre antral and antral follicles and its concentration is related to the number of small follicles and ovarian reserve [82–84]. As age decrease, AMH gradually decrease which is a sign of a lower ovarian reserve [83]. In fact, it is well established that AMH levels are strongly correlated with age, and age is strongly correlated with oxidative stress and infertility [85,86]. A recent study has shown that starting from age 37, the antioxidant pattern in FF decreases and impairs ROS scavenging system [87], which can be an indicator of infertility and a greater risk of IVF failure. Yet, our data showed no direct association between CM MDA levels and women's age AMH and idiopathic infertility.

## **Conclusion**

According to all these findings, although, embryo quality could be associated with other parameters that are not always associated with oxidative stress such as fragmentation levels and sperm quality, or parameters that are related to oxidative stress such as age and AMH levels, oxidative stress in IVF CM might be mainly associated to delayed embryonic development.

## **References**

1. Apel K, Hirt H. Reactive oxygen species: Metabolism, Oxidative Stress, and Signal Transduction. *Annu Rev Plant Biol.* 2004;55(1):373–99.

2. Murphy MP. How mitochondria produce reactive oxygen species. *Biochem J*. 2009;417(1):1–13.
3. Schieber M, Chandel NS. ROS function in redox signaling and oxidative stress. *Curr Biol* [Internet]. 2014;24(10):R453–62. Available from: <http://dx.doi.org/10.1016/j.cub.2014.03.034>
4. Han Y, Ishibashi S, Iglesias-Gonzalez J, Chen Y, Love NR, Amaya E. Ca<sup>2+</sup>-Induced Mitochondrial ROS Regulate the Early Embryonic Cell Cycle. *Cell Rep* [Internet]. 2018;22(1):218–31. Available from: <https://doi.org/10.1016/j.celrep.2017.12.042>
5. Thannickal VJ, Fanburg BL. invited review Reactive oxygen species in cell signaling. *Am J Physiol Lung Cell Mol Physiol* [Internet]. 2000;279(78):1005–28. Available from: [http://www.ajplung.org/1005downloadedfromwww.physiology.org/journal/ajplungby\\$%7BindividualUser.givenNames%7D\\$%7BindividualUser.surname%7D](http://www.ajplung.org/1005downloadedfromwww.physiology.org/journal/ajplungby$%7BindividualUser.givenNames%7D$%7BindividualUser.surname%7D)
6. Guérin P, El Moutassim S, Ménéz Y. Oxidative stress and protection against reactive oxygen species in the pre-implantation embryo and its surroundings. *Hum Reprod Update*. 2001;7(2):175–89.
7. Lijie X, Ullah O, Haixing L, Ali I, Li Z, Fang NZ. Pioglitazone mediated reduction in oxidative stress and alteration in level of PPAR $\gamma$ , Nrf2 and antioxidant enzyme genes in mouse preimplantation embryo during maternal to zygotic transition. *Pak J Zool*. 2019;51(3):1085–92.
8. Hensley K, Robinson KA, Gabbita SP, Salsman S, Floyd RA. Reactive oxygen species, cell signaling, and cell injury. *Free Radic Biol Med*. 2000;28(10):1456–62.
9. Nixon VL, McDougall A, Jones KT. Ca<sup>2+</sup> oscillations and the cell cycle at fertilisation of mammalian and ascidian eggs. *Biol Cell*. 2000;92(3–4):187–96.
10. Jones DP. Radical-free biology of oxidative stress. *Am J Physiol - Cell Physiol*. 2008;295(4).
11. Ahmadi H, Fathi F, Moeini A, Amidi F, Sobhani A. Evaluation of prooxidant-antioxidant balance in in vitro fertilization-conceived mice. *Clin Exp Reprod Med*. 2018;45(2):82–7.
12. Combelles CMH, Gupta S, Agarwal A. Could oxidative stress influence the in-vitro maturation of oocytes? *Reprod Biomed Online* [Internet]. 2009;18(6):864–80. Available from: [http://dx.doi.org/10.1016/S1472-6483\(10\)60038-7](http://dx.doi.org/10.1016/S1472-6483(10)60038-7)
13. Lee R, Margaritis M, Channon K, Antoniades C. Evaluating Oxidative Stress in Human Cardiovascular Disease: Methodological Aspects and Considerations. *Curr Med Chem*. 2012;19(16):2504–20.
14. Truong TT, Soh YM, Gardner DK. Antioxidants improve mouse preimplantation embryo development and viability. *Hum Reprod*. 2016;31(7):1445–54.
15. Truong T, Gardner DK. Antioxidants improve IVF outcome and subsequent embryo development in the mouse. *Hum Reprod*. 2017;32(12):2404–13.
16. Ollero M, Gil-Guzman E, Lopez MC, Sharma RK, Agarwal A, Larson K, et al.

- Characterization of subsets of human spermatozoa at different stages of maturation: Implications in the diagnosis and treatment of male infertility. *Hum Reprod.* 2001;16(9):1912–21.
17. Borowiecka M, Wojsiat J, Polac I, Radwan M, Radwan P, Zbikowska HM. Oxidative stress markers in follicular fluid of women undergoing in vitro fertilization and embryo transfer. *Syst Biol Reprod Med.* 2012;58(6):301–5.
  18. Ahmed W, Lingner J. Impact of oxidative stress on telomere biology. *Differentiation.* 2018;99:21–7.
  19. Lan KC, Lin YC, Chang YC, Lin HJ, Tsai YR, Kang HY. Limited relationships between reactive oxygen species levels in culture media and zygote and embryo development. *J Assist Reprod Genet.* 2019;36(2):325–34.
  20. J. Premkumar B, Agarwal A. Female Infertility and Assisted Reproduction: Impact of Oxidative Stress-- An Update. *Curr Womens Health Rev.* 2012;8(3):183–207.
  21. Kim SG, Kim YY, Park JY, Kwak SJ, Yoo CS, Park IH, et al. Early fragment removal on in vitro fertilization day 2 significantly improves the subsequent development and clinical outcomes of fragmented human embryos. *Clin Exp Reprod Med.* 2018;45(3):122–8.
  22. Soto-Heras S, Paramio MT. Impact of oxidative stress on oocyte competence for in vitro embryo production programs. *Res Vet Sci [Internet].* 2020;132(May):342–50. Available from: <https://doi.org/10.1016/j.rvsc.2020.07.013>
  23. Ribas-Maynou J, Yeste M, Salas-Huetos A. The relationship between sperm oxidative stress alterations and IVF/ICSI outcomes: A systematic review from nonhuman mammals. *Biology (Basel).* 2020;9(7):1–18.
  24. Desjardins P, Hansen JB, Allen M. Microvolume protein concentration determination using the NanoDrop 2000c Spectrophotometer. *J Vis Exp.* 2010;(33):1–4.
  25. Jamil M, Debbarh H, Aboulmaouahib S, Aniq Filali O, Mounaji K, Zarqaoui M, et al. Reactive oxygen species in reproduction: Harmful, essential or both? *Zygote.* 2020;(May).
  26. Chen Q, Vazquez EJ, Moghaddas S, Hoppel CL, Lesnefsky EJ. Production of reactive oxygen species by mitochondria: Central role of complex III. *J Biol Chem.* 2003;278(38):36027–31.
  27. Lubberda Z. The role of glutathione in mammalian gametes. *Reprod Biol.* 2005;5(1):5–17.
  28. Camello-Almaraz C, Gomez-Pinilla PJ, Pozo MJ, Camello PJ. Mitochondrial reactive oxygen species and Ca<sup>2+</sup> signaling. *Am J Physiol - Cell Physiol.* 2006;291(5):1082–8.
  29. Camello-Almaraz MC, Pozo MJ, Murphy MP, Camello PJ. Mitochondrial production of oxidants is necessary for physiological calcium oscillations. *J Cell Physiol.* 2006;206(2):487–94.
  30. Feissner RF, Skalska J, Gaum WE, Sheu SS. Crosstalk signaling between mitochondrial Ca<sup>2+</sup> and ROS. *Front Biosci.* 2009;14(4):1197–218.

31. Takahashi T, Takahashi E, Igarashi H, Tezuka N, Kurachi H. Impact of oxidative stress in aged mouse oocytes on calcium oscillations at fertilization. *Mol Reprod Dev.* 2003;66(2):143–52.
32. Jana SK, K NB, Chattopadhyay R, Chakravarty B, Chaudhury K. Upper control limit of reactive oxygen species in follicular fluid beyond which viable embryo formation is not favorable. *Reprod Toxicol* [Internet]. 2010;29(4):447–51. Available from: <http://dx.doi.org/10.1016/j.reprotox.2010.04.002>
33. Bedaiwy MA, Falcone T, Mohamed MS, Aleem AAN, Sharma RK, Worley SE, et al. Differential growth of human embryos in vitro: Role of reactive oxygen species. *Fertil Steril.* 2004;82(3):593–600.
34. Martín-Romero FJ, Miguel-Lasobras EM, Domínguez-Arroyo JA, González-Carrera E, Álvarez IS. Contribution of culture media to oxidative stress and its effect on human oocytes. *Reprod Biomed Online* [Internet]. 2008;17(5):652–61. Available from: [http://dx.doi.org/10.1016/S1472-6483\(10\)60312-4](http://dx.doi.org/10.1016/S1472-6483(10)60312-4)
35. Shih YF, Lee TH, Liu CH, Tsao HM, Huang CC, Lee MS. Effects of reactive oxygen species levels in prepared culture media on embryo development: A comparison of two media. *Taiwan J Obstet Gynecol* [Internet]. 2014;53(4):504–8. Available from: <http://dx.doi.org/10.1016/j.tjog.2013.12.009>
36. Agarwal A, Gupta S, Sikka S. The role of free radicals and antioxidants in reproduction. *Curr Opin Obstet Gynecol.* 2006;18(3):325–32.
37. Dai S, Qiao Y, Jin H, Xin Z, Su Y, Sun Y, et al. CLINICAL CORNER : COMMUNICATION Effect of coincubation time of sperm-oocytes on fertilization , embryonic development , and subsequent pregnancy outcome. 2012;(November 2011):348–53.
38. Zhang XD, Liu JX, Liu WW, Gao Y, Han W, Xiong S, et al. Time of insemination culture and outcomes of in vitro fertilization : a systematic review and meta-analysis. 2013;19(6):685–95.
39. Kong X, Yang S, Gong F, Lu C, Zhang S, Lu G, et al. The relationship between cell number, division behavior and developmental potential of cleavage stage human embryos: A time-lapse study. *PLoS One.* 2016;11(4):1–11.
40. Blank C, DeCroo I, Weyers B, van Avermaet L, Tilleman K, van Rumste M, et al. Improvement instead of stability in embryo quality between day 3-5: A possible extra predictor for blastocyst selection. *Eur J Obstet Gynecol Reprod Biol* [Internet]. 2020;253:198–205. Available from: <https://doi.org/10.1016/j.ejogrb.2020.08.027>
41. Ozawa M, Hirabayashi M, Kanai Y. Developmental competence and oxidative state of mouse zygotes heat-stressed maternally or in vitro. *Reproduction.* 2002;124(5):683–9.
42. Karagenc L, Sertkaya Z, Ciray N, Ulug U, Bahçeci M. Impact of oxygen concentration on embryonic development of mouse zygotes. *Reprod Biomed Online* [Internet]. 2004;9(4):409–17. Available from: [http://dx.doi.org/10.1016/S1472-6483\(10\)61276-X](http://dx.doi.org/10.1016/S1472-6483(10)61276-X)
43. Zhang J, Wang X, Vikash V, Ye Q, Wu D, Liu Y, et al. ROS and ROS-Mediated Cellular Signaling. *Oxid Med Cell Longev.* 2016;2016(Figure 1).

44. Amaral CS, Koch J, Correa Júnior EE, Bertolin K, Mujica LKS, Fiorenza MF, et al. Heat stress on oocyte or zygote compromises embryo development, impairs interferon tau production and increases reactive oxygen species and oxidative stress in bovine embryos produced in vitro. *Mol Reprod Dev.* 2020;87(8):899–909.
45. Paria BC, Dey SK. Preimplantation embryo development in vitro: Cooperative interactions among embryos and role of growth factors. *Proc Natl Acad Sci U S A.* 1990;87(12):4756–60.
46. Prados FJ, Debrock S, Lemmen JG, Agerholm I. The cleavage stage embryo. *Hum Reprod.* 2012;27 Suppl 1:50–71.
47. Iwata K, Yumoto K, Sugishima M, Mizoguchi C, Kai Y, Iba Y, et al. Analysis of compaction initiation in human embryos by using time-lapse cinematography. *J Assist Reprod Genet.* 2014;31(4):421–6.
48. Tatemoto H, Sakurai N, Muto N. Protection of Porcine Oocytes Against Apoptotic Cell Death Caused by Oxidative Stress During In Vitro Maturation: Role of Cumulus Cells. *Biol Reprod.* 2000;63(3):805–10.
49. Dalvit GC, Cetica PD, Pintos LN, Beconi MT. Reactive oxygen species in bovine embryo in vitro production. *Biocell.* 2005;29(2):209–12.
50. Ufer. The roles of glutathione peroxidases during embryo development. *Front Mol Neurosci.* 2011;4(July):1–14.
51. Surai PF, Fisinin VI, Karadas F. Antioxidant systems in chick embryo development. Part 1. Vitamin E, carotenoids and selenium. *Anim Nutr [Internet].* 2016;2(1):1–11. Available from: <http://dx.doi.org/10.1016/j.aninu.2016.01.001>
52. Qian D, Li Z, Zhang Y, Huang Y, Wu Q, Ru G, et al. Response of Mouse Zygotes Treated with Mild Hydrogen Peroxide as a Model to Reveal Novel Mechanisms of Oxidative Stress-Induced Injury in Early Embryos. *Oxid Med Cell Longev.* 2016;2016.
53. Nasr-Esfahani MH, Aitken JR, Johnson MH. Hydrogen peroxide levels in mouse oocytes and early cleavage stage embryos developed in vitro or in vivo. *Development.* 1990;109(2):501–7.
54. Thomas M, Jain S, Kumar GP, Laloraya M. A programmed oxyradical burst causes hatching of mouse blastocysts. *J Cell Sci.* 1997;110(14):1597–602.
55. Morales H, Tilquin P, Rees JF, Massip A, Dessy F, Van Langendonck A. Pyruvate prevents peroxide-induced injury of in vitro preimplantation bovine embryos. *Mol Reprod Dev.* 1999;52(2):149–57.
56. Lopes AS, Lane M, Thompson JG. Oxygen consumption and ROS production are increased at the time of fertilization and cell cleavage in bovine zygotes. *Hum Reprod.* 2010;25(11):2762–73.
57. Yang HW, Hwang KJ, Kwon HC, Kim HS, Choi KW, Oh KS. Detection of reactive oxygen species (ROS) and apoptosis in human fragmented embryos. *Hum Reprod.* 1998;13(4):998–1002.
58. Gupta S, Agarwal A, Banerjee J, Alvarez JG. The role of oxidative stress in

- spontaneous abortion and recurrent pregnancy loss: A systematic review. *Obstet Gynecol Surv.* 2007;62(5):335–47.
59. Bedaiwy MA, Mahfouz RZ, Goldberg JM, Sharma R, Falcone T, Abdel Hafez MF, et al. Relationship of reactive oxygen species levels in day 3 culture media to the outcome of in vitro fertilization/intracytoplasmic sperm injection cycles. *Fertil Steril* [Internet]. 2010;94(6):2037–42. Available from: <http://dx.doi.org/10.1016/j.fertnstert.2009.12.020>
  60. Fujimoto VY, Bloom MS, Huddleston HG, Shelley WB, Ocque AJ, Browne RW. Correlations of follicular fluid oxidative stress biomarkers and enzyme activities with embryo morphology parameters during in vitro fertilization. *Fertil Steril* [Internet]. 2011;96(6):1357–61. Available from: <http://dx.doi.org/10.1016/j.fertnstert.2011.09.032>
  61. Lee TH, Lee MS, Liu CH, Tsao HM, Huang CC, Yang YS. The association between microenvironmental reactive oxygen species and embryo development in assisted reproduction technology cycles. *Reprod Sci.* 2012;19(7):725–32.
  62. Dozortsev D, Ermilov A, El-Mowafi DM, Diamond M. The impact of cellular fragmentation induced experimentally at different stages of mouse preimplantation development. *Hum Reprod.* 1998;13(5):1307–11.
  63. Keltz M, Fritz R, Gonzales E, Ozensoy S, Skorupski J, Stein D. Defragmentation of low grade day 3 embryos resulted in sustained reduction in fragmentation, but did not improve compaction or blastulation rates. *Fertil Steril* [Internet]. 2010;94(6):2406–8. Available from: <http://dx.doi.org/10.1016/j.fertnstert.2010.03.014>
  64. Alikani M, Cohen J, Tomkin G, Garrisi GJ, Mack C, Scott RT. Human embryo fragmentation in vitro and its implications for pregnancy and implantation. *Fertil Steril.* 1999;71(5):836–42.
  65. Stensen MH, Tanbo TG, Storeng R, Åbyholm T, Fedorcsak P. Fragmentation of human cleavage-stage embryos is related to the progression through meiotic and mitotic cell cycles. *Fertil Steril.* 2015;103(2):374–381.e4.
  66. Zhang Y, Liu L, Yin TL, Yang J, Xiong CL. Follicular metabolic changes and effects on oocyte quality in polycystic ovary syndrome patients. *Oncotarget.* 2017;8(46):80472–80.
  67. Safari S, Khalili MA, Barekati Z, Halvaei I, Anvari M, Nottola SA. Cosmetic micromanipulation of vitrified-warmed cleavage stage embryos does not improve ART outcomes: An ultrastructural study of fragments. *Reprod Biol* [Internet]. 2017;17(3):210–7. Available from: <http://dx.doi.org/10.1016/j.repbio.2017.05.003>
  68. Stigliani S, Anserini P, Venturini PL, Scaruffi P. Mitochondrial DNA content in embryo culture medium is significantly associated with human embryo fragmentation. *Hum Reprod.* 2013;28(10):2652–60.
  69. Halvaei I, Khalili MA, Esfandiari N, Safari S, Talebi AR, Miglietta S, et al. Ultrastructure of cytoplasmic fragments in human cleavage stage embryos. *J Assist Reprod Genet* [Internet]. 2016;33(12):1677–84. Available from: <http://dx.doi.org/10.1007/s10815-016-0806-1>
  70. Johansson M, Hardarson T, Lundin K. There is a cutoff limit in diameter between a blastomere and a small anucleate fragment. *J Assist Reprod Genet.* 2003;20(8):309–13.

71. Otasevic V, Surlan L, Vucetic M, Tulic I, Buzadzic B, Stancic A, et al. Expression patterns of mitochondrial OXPHOS components, mitofusin 1 and dynamin-related protein 1 are associated with human embryo fragmentation. *Reprod Fertil Dev*. 2016;28(3):319–27.
72. Pereira N, Palermo GD. Intracytoplasmic sperm injection: History, indications, technique, and safety. *Intracytoplasmic Sperm Inject Indic Tech Appl*. 2018;9–21.
73. Esteves SC, Roque M, Bedoschi G, Haahr T, Humaidan P. Intracytoplasmic sperm injection for male infertility and consequences for offspring. *Nat Rev Urol [Internet]*. 2018;15(9):535–62. Available from: <http://dx.doi.org/10.1038/s41585-018-0051-8>
74. Danis RB, Samplaski MK. Sperm Morphology: History, Challenges, and Impact on Natural and Assisted Fertility. *Curr Urol Rep*. 2019;20(8).
75. Mangoli E, Khalili MA, Talebi AR, Agha-Rahimi A, Soleimani M, Faramarzi A, et al. IMSI procedure improves clinical outcomes and embryo morphokinetics in patients with different aetiologies of male infertility. *Andrologia*. 2019;51(8):1–9.
76. Colaco S, Sakkas D. Paternal factors contributing to embryo quality. *J Assist Reprod Genet*. 2018;35(11):1953–68.
77. Chapuis A, Gala A, Ferrières-Hoa A, Mullet T, Bringer-Deutsch S, Vintejou E, et al. Sperm quality and paternal age: Effect on blastocyst formation and pregnancy rates. *Basic Clin Androl [Internet]*. 2017;27(1):1–9. Available from: <http://dx.doi.org/10.1186/s12610-016-0045-4>
78. Aston KI, Uren PJ, Jenkins TG, Horsager A, Cairns BR, Smith AD, et al. Aberrant sperm DNA methylation predicts male fertility status and embryo quality. *Fertil Steril [Internet]*. 2015;104(6):1388–1397.e5. Available from: <http://dx.doi.org/10.1016/j.fertnstert.2015.08.019>
79. Mitchell V, Sigala J, Jumeau F, Ballot C, Peers MC, Decanter C, et al. ICSI treatment in severe asthenozoospermia. *Gynecol Obstet Fertil*. 2012;40(12):766–9.
80. Hashimoto H, Ishikawa T, Goto S, Kokeyuchi S, Fujisawa M, Shiotani M. The effects of severity of oligozoospermia on intracytoplasmic sperm injection (ICSI) cycle outcome. *Syst Biol Reprod Med*. 2010;56(1):91–5.
81. Mangoli E, Khalili MA, Talebi AR, Kalantar SM, Montazeri F, Agharahimi A, et al. Association between early embryo morphokinetics plus transcript levels of sperm apoptotic genes and clinical outcomes in IMSI and ICSI cycles of male factor patients. *J Assist Reprod Genet*. 2020;37(10):2555–67.
82. Weenen C, Laven JSE, von Bergh ARM, Cranfield M, Groome NP, Visser JA, et al. Anti-Müllerian hormone expression pattern in the human ovary: Potential implications for initial and cyclic follicle recruitment. *Mol Hum Reprod*. 2004;10(2):77–83.
83. Hsu MI. Changes in the PCOS phenotype with age. *Steroids [Internet]*. 2013;78(8):761–6. Available from: <http://dx.doi.org/10.1016/j.steroids.2013.04.005>
84. Hamza SM, Abd-Alrahman SJ, Raheem SM. Correlation between Levels of Serum Antioxidants and Numerous Hormones In Primary Infertility of Women. 2016;3(3):17–20.

85. Iliodromiti S, Anderson RA, Nelson SM. Technical and performance characteristics of anti-Müllerian hormone and antral follicle count as biomarkers of ovarian response. *Hum Reprod Update*. 2015;21(6):698–710.
86. Cui L, Qin Y, Gao X, Lu J, Geng L, Ding L, et al. Antimüllerian hormone: correlation with age and androgenic and metabolic factors in women from birth to postmenopause. *Fertil Steril*. 2016;105(2):481-485.e1.
87. Debbarh H, Louanjli N, Aboulmaouahib S, Jamil M, Ahbbas L, Kaarouch I, et al. Antioxidant activities and lipid peroxidation status in human follicular fluid: Age-dependent change. *Zygote*. 2021;

UNDER PEER REVIEW