

Original Research Article  
**Effect of Different Cooking Methods on  
the Concentration of Vitamin C in *Vigna  
unquiculata***

---

**ABSTRACT**

**Background:** Vitamin C is a water-soluble vitamin widely distributed in nature, mostly rich in fresh fruits and green leafy vegetables. The amount of Vitamin C in a given food depends on soil condition, climate in their area of growth, storage condition after harvest, and methods of preparation. Much amount of Vitamin C in vegetables is lost during cooking. It is upon this background that this study aimed to determine the concentration of Vitamin C in *Vigna unquiculata* when cooked using different cooking methods. (Authors should improve the end of the introduction; since there is not a good link between the information that was given and the objective of the work.)

**Methodology:** Vitamin C concentration in a solution of *Vigna unquiculata* cooked by different methods was determined by a redox titration using iodine in the presence of a starch indicator. This was conducted at Busitema University, Faculty of Science and Education, Biology Laboratory in Uganda. ()

**Results:** The study revealed that cooking *Vigna unquiculata* using microwave yields the highest concentration of vitamin C (154.8 mg/L), followed by steaming (142.9 mg/L) and boiling yielding the least concentration of vitamin C (130.4 mg/L).

**Conclusion:** The concentration of vitamin C obtained by the three methods of cooking *Vigna unquiculata* has shown that microwaving method of cooking yields the highest concentration of vitamin C followed by the steaming method and finally boiling. Further studies should be done to explain the variation of vitamin C concentration in *Vigna unquiculata* when cooked using the above cooking methods. Similar studies should be conducted on other leafy vegetables which are potential sources of Vitamin C.

*Keywords: Vigna unquiculata; cooking methods; boiling; steaming; microwaving; vitamin C concentrations.*

**1. INTRODUCTION the introduction must be improved and the references must be updated.**

Vitamin C, also known as ascorbic acid, is a water-soluble vitamin required for several functions in the human bodies-body [1]. Its chemical formula is C<sub>6</sub>H<sub>8</sub>O<sub>6</sub> [2]. Vitamin C plays an important role in forming collagen, a protein that gives structure to bones, muscles, and blood vessels; in blocking the formation of nitrosamines, potentially cancer-causing compounds; in increasing the absorption of iron; in stimulating the excretion of lead; and in promoting resistance to infection [3]. Also, it is a highly effective antioxidant, acting to lessen oxidative

stress [4]. The clinical importance of vitamin C relates principally to its role as a cofactor in a number of some enzymes reactions involved in collagen synthesis, the dysfunction of which disrupts connective tissue integrity, resulting in impaired wound healing and capillary bleeding [5]. Previous studies from meta-analysis shows that each 0.2 serving per day increase in green leafy vegetable intake was associated with a 13% lower risk of type 2 diabetes [6]. However, human bodies cannot synthesize vitamin C and it must be obtained either from diet or dietary supplements [7]. Vitamin C is widely distributed in nature, mostly rich in fresh fruits and green leafy vegetables such as guava, mango, tomato, cabbage, green paper, and Vigan spp., among others [8]. (Update the biological activities attributed to vitamin C) The amount of vitamin C in food of plant origin depends on the variety of the plant, soil condition, climate in which they grow, of time after the harvest, fruit storage conditions and methods of preparation [9]. Study-Studies have shown that vitamin C is also lost when cooking depending on the degree of heating, the surface area exposed to water, oxygen, pH and the presence of transitional metal [10]. This raises a lot of questions on what method of cooking vegetables preserves a good quantity of vitamin C since most vegetables are commonly cooked before being consumed. This study therefore, thought to investigate the effects of three different cooking methods (i.e., steaming, microwaving, and boiling) on the vitamin C content of *Vigna unguiculata* vegetable. In the introduction the authors should make it clear why it is important to evaluate the **Effect of Different Cooking Methods on the Concentration of Vitamin C in *Vigna unguiculata***

Formatted: Font: 12 pt

## 2. METHODOLOGY

### 2.1 Preparation of starch indicator

0.5g of soluble starch was weighed and 50 ml of nearly boiling water was added in to a 100 ml conical flask, stirred to dissolve, and then cooled before usage.

Comment [SD1]: Do not start the sentence with number.

### 2.2 Preparation of Iodine solution the standard solution


0.85g of potassium iodide (KI) and 0.5g of iodine crystal were dissolved in 50ml of distilled water in a 250ml beaker. The solution was then made to the final volume of 250ml with distilled water resulting into 0.005 molar iodine solutions.

### 2.3 Procedure for determining Vitamin C concentration in *Vigna Unguiculata*

Fifty (50)g of *Vigna Unguiculata* leaves were cut into smaller pieces and cooked by the three different methods under investigation: boiling, steaming, and microwaving for a period of 0, 5, 10, 15, 20, 25 and 30 minutes. The juice extracts were obtained by blending different cooked samples with 200 ml of distilled water using a blending machine for a period of two minutes. The blended samples were filtered using a sieve and the filtrate was put in different 250ml beakers for each cooking method each containing 200 mL of the extract. Twenty-five (25) mL of the sample were pipetted in a clean 250 mL conical flask and 20 drops of 1% starch indicator was-were added. The sample was titrated with 0.005 molar iodine solutions until the endpoint was reached (i.e., the point at which the stable traces of blue-black color is formed due to the formation of iodine-starch complex) and the volume of the iodine solution (standard solution) used in the titration was recorded. (Extraction buffer what proportion?)

### 2.4 Determination of vitamin C concentration in *Vigna unguiculata*

Iodine solution was used as the standard solution to determine the unknown concentrations of vitamin C in the different cooking methods using the reaction equation and the formula below.



$$[C_6H_8O_6] = \frac{[I_2] \times \text{Av. Vol of } I_2}{\text{Vol of } C_6H_8O_6}$$

### 2.3. Data analysis


One-way ANOVA was used to determine the significance of the statistical differences in the concentrations of vitamin C between the three different methods of cooking: boiling, steaming, and microwaving and significance was accepted at  $P \leq 0.05$ . The Pearson's product-moment correlation coefficient ( $r$ ) was used to determine the relationship between time and concentrations of Vitamin C in *Vigna unguiculata* for the three methods of cooking. The influence of time as a factor of determination of the concentration of vitamin C for the three methods of cooking were predicted using the coefficient of determination ( $r^2$ ).

Comment [SD2]: Correct spelling

## 3. RESULTS (HERE I MISSED A PROFOUND PLANT PHYSIOLOGY DISCUSSION. BESIDES, YOU ONLY DISCUSS THE RESULTS)

### 3.1 Effects of cooking methods on the concentrations of Vitamin C in *Vigna Unguiculata*


The results obtained showed that there is variation in the concentration of Vitamin C *Vigna unguiculata* when subjected to the three different cooking methods. The highest concentration of Vitamin C is found to be in the microwaving method of cooking (154.8 mg/L), followed by steaming (142.9 mg/L) and boiling (130.4 mg/L), (Figure 1 A,  $F(7) = 5.85$ ,  $p = .011$ ). The concentration of vitamin C in the control (un-cooked) was higher (163 mg/L) compared to those of the cooking methods because the extract used in the control experiment was obtained from a fresh sample of *Vigna unguiculata* without subjecting to cooking. Gradual decrease in the concentration of vitamin C were observed in all the three methods of cooking over a period of 30 minutes (Figure 1 B).



**Figure 1. Effects of cooking methods on the concentrations of Vitamin C in *Vigna unguiculata*:** Differences in concentrations of vitamin C after boiling, steaming, and microwaving the vegetable after 30 minutes (A), Effect of boiling, steaming and microwaving methods on the vitamin C concentration during the 30 minutes (B), The statistical significance of the differences between the methods of cooking was determined using One-way ANOVA ( $F(7) = 5.85, p = .011$ ).

### 3.2 Impact of cooking time on the concentration of vitamin C in *Vigna unguiculata*

There were strong negative correlations between the cooking time and concentrations of vitamin C in all the three methods of cooking: boiling,  $r(4) = -.92, p = .009$ , steaming,  $r(4) = -.85, p = .034$ , and microwaving,  $r(4) = -.89, p = .018$ , (Figure 2 A, B, and C). The cooking time significantly impacted the concentrations of vitamin C in all the three cooking methods: boiling;  $r^2 = .85, p = .009$ , steaming;  $r^2 = .72, p = .034$ ; microwaving,  $r^2 = .79, p = .018$ , (Figure 2 A, B and C).



**Figure 2. Impact of cooking time on the concentration of vitamin C in *Vigna unguiculata*:** Relationship between boiling time (A), steaming time (B) and microwaving time (C) and the concentration of vitamin C. Correlation coefficient between the cooking time and the concentration of vitamin C was determined using Pearson's product-moment correlation coefficient, (boiling,  $r(4) = -.92, p = .009$ ; steaming,  $r(4) = -.85, p = .034$ ; microwaving,  $r(4) = -.89, p = .018$ ) and the influence of time as a factor of determination of the concentration of vitamin C for the three methods of cooking were predicted using the coefficient of determination (boiling,  $r^2 = .85, p = .009$ ; steaming,  $r^2 = .72, p = .034$ ; microwaving,  $r^2 = .79, p = .018$ ).

## 4. DISCUSSION

This study shows that boiling method of cooking results in the highest loss of vitamin C content in *Vigna unguiculata* vegetables compared to steaming and microwaving methods of cooking. The high loss of vitamin C content during boiling could be attributed to the fact that vitamin C is very soluble in water and not stable at high temperatures [11]. Loss of vitamin C occurs primarily by chemical degradation that involves the oxidation of ascorbic acid to dehydroascorbic acid (DHAA), followed by hydrolysis to 2,3-diketogulonic acid and further polymerization to form other nutritionally inactive products [12]. Thus, the boiling temperature could have inactivated most of the vitamin C in the vegetable, while the water would also wash away the vitamin C during the boiling process [13]. Therefore, despite the fact that although boiling makes leafy vegetables like *Vigna unguiculata* less toxic and more palatable, it greatly affects their vitamin C contents [13]. Hence, a balance must be made between palatability and nutritional content by either reducing the boiling time or using alternative methods of cooking like steaming or microwaving that would retain more vitamin C than boiling method.

This study also demonstrated that microwaving method of cooking has the highest retention of vitamin C in the green leafy *Vigna unguiculata* vegetable. This contradicts a finding of a similar study conducted by Zeng C. ~~On~~ on some selected vegetables (broccoli, spinach, and lettuce) ~~w~~here out of the three methods of cooking, the cooking method that resulted in the least loss of vitamin C was the steaming method [3]. This difference in the findings could be ~~due to the fact~~ that since the loss of vitamin C in vegetables also depend on the particular vegetables. In different vegetables, there are varying rates of degradation of vitamin C contents by different methods of cooking [14].

*Vigna unguiculata* is one of the commonest vegetables routinely consumed to provide vitamin C requirements s in the diet. However, microwaving method of cooking cannot be applicable in some rural areas due to the expensive nature of the microwaving equipment and inadequate electric power supply. The local community can therefore use the steaming method of cooking ~~in order~~ to obtain a considerable quantity and quality of vitamin C.

## 5. CONCLUSION (REWRITE THE PARAGRAPH, AS SOME THINGS BECOME REDUNDANT.)

The study showed that cooking methods affects ~~e~~ the concentration of vitamin C in *Vigna unguiculata* with microwaving showing the highest retention of Vitamin C, followed by steaming and boiling showing the least retention of vitamin C. There is a need for extensive studies on other factors other than cooking methods that may affect the concentration of vitamin C in leafy vegetables such as *Vigna unguiculata*. There is also a need to replicate this study using other leafy vegetables which contain vitamin C.

## REFERENCES

1. May, J.M., *Vitamin C transport and its role in the central nervous system*. Water Soluble Vitamins, 2012: p. 85-103.
2. Saeed, A.M., M.J. Hamzah, and N.J.M. Ali, *Sensitive spectrophotometric method for determination of vitamin (C and A)*. International Journal of Pharmaceutical Sciences and Research, 2018. **9**(8): p. 3373-3377.
3. Zeng, C., *Effects of different cooking methods on the vitamin C content of selected vegetables*. Nutrition & Food Science, 2013.
4. Frei, B., I. Birlouez-Aragon, and J. Lykkesfeldt, *Authors' perspective: What is the optimum intake of vitamin C in humans?* Critical reviews in food science and nutrition, 2012. **52**(9): p. 815-829.
5. Penny, H., et al., *The role of macronutrients and micronutrients in wound healing: a narrative review*. Journal of Wound Care, 2022. **31**(5): p. S14-S22.
6. Li, M., et al., *Fruit and vegetable intake and risk of type 2 diabetes mellitus: meta-analysis of prospective cohort studies*. BMJ open, 2014. **4**(11): p. e005497.
7. Cha, J., et al., *Ascorbate supplementation inhibits growth and metastasis of B16FO melanoma and 4T1 breast cancer cells in vitamin C-deficient mice*. International journal of oncology, 2013. **42**(1): p. 55-64.
8. Phek, K.V.K., *RESOURCE INVENTORY OF DISTRICT PHEK-NAGALAND*.
9. Rivelli, A.R., et al., *Vitamin C content in leaves and roots of horseradish (*Armoracia rusticana*): Seasonal variation in fresh tissues and retention as affected by storage conditions*. Emirates Journal of Food and Agriculture, 2017: p. 799-806.
10. Vinishdharma, T. and K. Don, *Vitamin C as a therapeutic healing agent*. Drug Invention Today, 2019. **12**(3).
11. Dewanto, V., et al., *Thermal processing enhances the nutritional value of tomatoes by increasing total antioxidant activity*. Journal of agricultural and food chemistry, 2002. **50**(10): p. 3010-3014.
12. Gregory, J., *Vitamins in food chemistry*. Food chemistry, 1996. **3**: p. 532-590.

13. Oboh, G., *Effect of blanching on the antioxidant properties of some tropical green leafy vegetables*. LWT-Food Science and Technology, 2005. **38**(5): p. 513-517.
14. Lee, S., et al., *Effect of different cooking methods on the content of vitamins and true retention in selected vegetables*. Food science and biotechnology, 2018. **27**(2): p. 333-342.

UNDER PEER REVIEW