

Original Research Article

Influence of thermal regimes and irrigation schedules on growth, yield attributes and yield of wheat (*Triticum aestivum* L.) varieties

Abstract

A field experiment was conducted at Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh, India during the *rabi* season of 2020-2021 and 2021-2022, to study the Influence of thermal regimes and irrigation schedules on growth, yield attributes and yield of wheat (*Triticum aestivum* L.) varieties. The experiment was conducted in split plot design with three thermal regimes (3rd December, 18th December and 2nd January), four irrigation schedules (IW: CPE=1.0, IW: CPE= 0.9, IW: CPE= 0.8 and IW: CPE=0.7) and two varieties (Lok 1 and MP 3336) with three replications. Sowing time with proper application of irrigation and varieties significantly influenced yield attributes and ultimately yield of the wheat. Higher grain yield (4,735 kg ha⁻¹) was observed under **crop sown on 3rd December thermal regimes with IW: CPE=1.0, irrigation schedule and variety MP 3336** over rest of the treatments. It may be due to the maximum number of effective tillers (484.2 m⁻²), length of earhead (8.70 cm), no. of grains earhead⁻¹ (45.5) and harvest index (41.91%) as compare to other treatment combinations. Significant lowest grain yield was noted under 2nd January thermal regimes with IW: CPE= 0.7, irrigation schedule and variety Lok 1. **Wheat sown under 3rd December thermal regime with irrigation schedule (IW: CPE= 1.0) and variety MP 3336 produced significantly more yield attributes and yield.**

Keywords: Grain yield, straw yield, wheat, thermal regimes, irrigation schedules

Introduction

“Wheat (*Triticum aestivum* L.) is an essential cereal grain and staple food for millions of people worldwide, as well as an important crop for India during the green revolution and post green revolution era. It is India's second most important crop after rice, occupying approximately 29 mha and producing 109.1 mt” (USDA.2021). In

Madhya Pradesh, this crop is presently being cultivated on 5.52 mha with a production of 15.47 mt and productivity of 3198 kg ha⁻¹ (Department of Agriculture, M.P., 2020). Wheat demand in the developing world is expected to increase by 60% by 2050 (CIMMYT, 2013).

“The thermal environment is one of the most important factors governing crop phenology and the efficient conversion of biomass into economic yield. Wheat yield is highly affected by late sown environment. Wheat is sown at different times of the year for a variety of reasons, including the presence of previous crops and diversity in cropping systems. Wheat planted late usually experiences high temperatures during the grain filling period, results in a lower crop yield. Because of the longer growing period, early sowing produces a higher yield than late sowing” (Munir *et al.*, 2002; Tanveer *et al.*, 2003). “It has been realized that the average yield of wheat of this region, sown during November sown environment, is well comparable to the state average, but the declining trend in wheat yield has been noticed with delayed sowing i.e., in the months of December and January. A sudden increase in temperature for 4-5 days at any stage of the wheat crop can reduce crop yield (Spiertz *et al.*, 2006), and even a one-day abnormal increase in temperature at the grain formation and filling stages can result in significant grain losses” (Alexander *et al.* 2010). It is primarily due to the shorter growth period available to late-sown wheat, combined with high temperatures and hot winds during the reproductive growth period, tends to result in forced maturity and, ultimately, poor grain yield.

“Water is necessary at every stage of plant development, from seed germination to plant maturation, in order to harvest wheat with the highest potential yield. The optimum use of irrigation water enables the effective utilization of all other production elements and increases yield per unit of area and time. There is a positive correlation between grain yield and irrigation frequencies” (Kumar *et al.*, 2012). “Irrigation had a significant impact on dry matter accumulation, leaf area index, number of spike (m⁻²), number of grains per spike and test weight that ultimately led to high grain yield, straw yield and harvest index”. (Malik *et al.* 2012).

Varieties play an important role in obtaining higher yield of wheat crop. The varieties are losing their yield potential due to changes in various edaphic and environmental conditions. Therefore, the selection of suitable variety for the particular location is of paramount importance. Keeping these facts in mind, the current study was designed to determine the Influence of thermal regimes and irrigation schedules on growth, yield attributes and yield of wheat (*Triticum aestivum* L.) varieties.

Materials and Methods

The experiment was conducted on wheat crop during *Rabi* season 2020-2021 and 2021-2022 at Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh. The soil of experimental land was sandy clay loam with pH 7.1, EC 0.25 ds/m and 0.54 % organic carbon, medium in available N (257.18 kg/ha), P (15.83 kg/ha) and K (301.25 kg/ha). The site of experiment is located at 23°09' N latitude and 79°58' E longitude and an altitude of 411 above mean sea level. Twenty-four treatment combinations, consisting of three thermal environments (3rd December, 18th December and 2nd January) in main plots, four IW: CPE ratio-based irrigation schedules ($I_1= 1.0$, $I_2= 0.9$, $I_3= 0.8$ and $I_4= 0.7$) in sub-plots and two wheat varieties ($V_1=$ Lok 1 and $V_2=$ MP 3336) in sub-sub plots were laid out in split plot design with three replications. The field was prepared properly to carry out wheat sowing. The graded and healthy seed of wheat crop was sown in line 20 cm apart with seed rate of 125 kg/ha. Fertilizers were applied uniformly to all the plots through urea, di ammonium phosphate and muriate of potash @120-60-40 kg N-P₂O₅-K₂O per ha. Half of the nitrogen and full dose of phosphorus and potash were applied as basal application and remaining nitrogen in two split doses. Irrigations as per treatments were applied in individual plots. Depth of irrigation was 5 cm for each irrigation schedule. However, a shallow come-up Irrigation was given immediately after sowing of the wheat crop to all the treatments. Total rainfall received was 22.0 mm and 51.5 mm distributed in 2 and 7 rainy days during first and second year respectively. The IW: CPE ratios were calculated based on depth of irrigation water and the cumulative pan evaporation during the period. After crop establishment, five plants were randomly selected from each plot for recording periodical observations on growth and yield attributing parameters like plant height,

number of effective tillers (m^{-2}), earhead length, number of grains per earhead, 1000-grain weight, grain yield and straw yield.

Statistical Analysis

In the present study the data analysis for split plot design has been performed using OP stat software.

3. Results and Discussion

3.1. Effect of thermal regime

Sowing environment plays important role in the growth, development and yield of wheat. Different growth, yield attributes and yield of wheat varieties vary significantly under different thermal regimes. The crop sown on 3rd December thermal regime recorded the highest plant height (94.0 cm), number of effective tillers m^{-2} (484.2), earhead length (8.70 cm), number of grains per earhead (45.5), 1000 grain weight (45.5 g). Grain yield ($4,735 \text{ kg ha}^{-1}$), straw yield ($6,580 \text{ kg ha}^{-1}$) and harvest index (41.91%) were also maximum over rest of the thermal regimes (Table 1). It might be due to the congenial environment condition which enhances growth and nutrients uptake, resulting in delay maturity and ultimately increases the grain yield, whereas delay in crop sowing cause reduction in yield (Choudhary *et al.* (2010). The reduction in yield and lower magnitude of yield attributes in late sown crop might be due to the exposure to higher temperature during the reproductive phase of the crop. An increase in temperature tends to promote growth, which in turn dilutes the amount of carbohydrates and chlorophyll in late-sown crops. Thus, these findings confirm those of Agrawal *et al.* (2001), Throat *et al.* (2015), Prasad *et al.* (2017), Bobade *et al.* (2017) and Khande *et al.* (2021).

3.2. Effect of irrigation schedules

Water is essential at every developmental phase starting from seed germination to plant maturation for harvesting the maximum potential yield of wheat. Treatment I₁ (Irrigation at 1.0, IW/CPE ratio) significantly increased the plant height (96.4 cm), number of effective tillers m^{-2} (489.9), earhead length (8.50 cm), number of grains per earhead (47.2) and 1000 grain weight (45.8 g) over I₂, I₃ and I₄ (Table 1). Grain yield

(4,496 kg ha⁻¹), straw yield (6,335 kg ha⁻¹) and harvest index (41.52%) were also maximum as compare to rest of the irrigation schedules (Table 1). The lowest growth, yield and yield attributes were found under I₄ (IW: CPE= 0.7) irrigation schedule. This was due to fewer number of irrigations I₄ (IW: CPE= 0.7) than I₁ (IW: CPE= 1.0) ratio which received more irrigations during its whole growth period. All the growth, yield attributes and yield are significantly superior under irrigation schedule I₁ (IW: CPE= 1.0) probably because of the increased water availability in the root zone which resulted in absorption of nutrients from the soil and ultimately gives vigorous growth and reproductive capacity of the wheat. This finding is in conformity with Salunkhe *et al.* (2012), Verma *et al.* (2017), Deo *et al.* (2017), Pal *et al.* (2020) and Gajbhiye (2021) also reported similar results.

3.3. Effect of varieties

Selection of suitable crop varieties according to the agroclimatic conditions may play crucial role in realizing the optimum production of any crop commodity (Singh *et al.*, 2008). Result of present study indicated that variety MP 3336 attained significantly higher grain yield (4,341 kg ha⁻¹) and straw yield (5,836 kg ha⁻¹) as compared to Lok 1. This might be due to higher yield attributes viz., number of effective tillers m⁻² (445.9), earhead length (8.24 cm) and number of grains per earhead (44.6) and harvest index (42.69%). This could be because of genetic makeup of that particular variety as well as climatic requirement of the different varieties. These results are similar to those of Tripathi *et al.* (2013), Alam *et al.* (2013), Satyendra Kumar *et al.* (2013) and Prasad *et al.* (2017).

4. Conclusion

On the basis of present investigation, it may be concluded that the 3rd December thermal regime with irrigation schedule (IW: CPE= 1.0) and MP 3336 produced higher yield attributes viz., number of effective tillers m⁻², length of earhead, number of grains per earhead and harvest index resulting in a higher grain yield than Lok 1.

Table 1: Effect of thermal regimes and irrigation schedules on growth, yield attributes and yield of wheat varieties (mean of two years)

| Treatments | Plant height (cm) | No. of effective tillers (m ⁻²) | Length of earhead (cm) | Number of grains earhead ⁻¹ | 1000 grain weight (g) | Grain yield (kg ha ⁻¹) | Straw yield (kg ha ⁻¹) | Harvest index |
|---|-------------------|---|------------------------|--|-----------------------|------------------------------------|------------------------------------|---------------|
| Thermal regimes | | | | | | | | |
| E₁- 03 December | 94.0 | 484.2 | 8.70 | 45.5 | 45.5 | 4,735 | 6,580 | 41.91 |
| E₂- 18 December | 91.6 | 432.0 | 8.08 | 42.9 | 43.6 | 4,244 | 5,594 | 43.21 |
| E₃- 02 January | 86.8 | 374.2 | 7.41 | 40.4 | 40.6 | 3,660 | 5,031 | 42.00 |
| SEm± | 0.89 | 6.83 | 0.05 | 0.49 | 0.49 | 59.35 | 73.82 | 0.09 |
| CD (5%) | 3.46 | 26.66 | 0.21 | 1.92 | 1.90 | 231.71 | 288.21 | 0.34 |
| Irrigation schedules (IW: CPE Ratio) | | | | | | | | |
| I₁- 1.0 | 96.4 | 489.9 | 8.50 | 47.2 | 45.8 | 4,496 | 6,335 | 41.52 |
| I₂- 0.9 | 93.1 | 452.9 | 8.25 | 44.5 | 44.2 | 4,342 | 5,969 | 42.11 |
| I₃- 0.8 | 89.2 | 411.7 | 7.94 | 41.8 | 42.6 | 4,146 | 5,570 | 42.63 |
| I₄- 0.7 | 84.6 | 366.2 | 7.58 | 38.2 | 40.4 | 3,868 | 5,067 | 43.24 |
| SEm± | 0.84 | 5.48 | 0.06 | 0.56 | 0.52 | 47.69 | 46.21 | 0.18 |
| CD (5%) | 2.49 | 16.29 | 0.18 | 1.66 | 1.53 | 141.72 | 137.31 | 0.55 |
| Varieties | | | | | | | | |
| V₁- Lok1 | 93.6 | 414.4 | 7.89 | 41.2 | 45.1 | 4,085 | 5,634 | 42.06 |
| V₂- MP 3336 | 88.1 | 445.9 | 8.24 | 44.6 | 43.4 | 4,341 | 5,836 | 42.69 |
| SEm± | 0.56 | 3.82 | 0.04 | 0.39 | 0.36 | 21.59 | 27.99 | 0.10 |
| CD (5%) | 1.63 | 11.15 | 0.11 | 1.13 | 1.04 | 63.02 | 81.72 | 0.30 |




Fig. 1 Yield attributes of wheat as influenced under various treatments




Fig. 2 Grain yield, straw yield and harvest index under various treatments

References

- Agrawal KK, Shanker U, Upadhyay AP and Gupta VK. 2001. Phenological behaviour and thermal requirements of wheat genotypes under different dates of sowing. *Annals of Plant and Soil Research* 3 (1):12-16.
- Alam MP, Kumar S, Ali N, Manjhi RP, Kumari N, Lakra RK and Izhar T. 2013. Performance of wheat varieties under different sowing dates in Jharkhand. *J. Wheat Res.* 5 (2) : 61-64.
- Alexander B, Hayman PMC, Donald G, Talukder H ASM, Gill G. Characterizing the risk of heat stress on wheat in South Australia: Meteorology, climatology and the design of a field heating chamber. In: Dove, H. (Ed.), *Food Security from Sustainable Agriculture: Proceedings of 15th Agronomy Conference 2010*. Lincoln, New Zealand, 2010.
- Anonymous. 2020. Department of agriculture & farmer welfare. Madhya Pradesh
- Anonymous. 2021. World Agriculture Production by USDA, <https://ipad.fas.usda.gov/search.aspx>
- Bobade P, S.K. Chandrawanshi, Patel SR and Kaushik DK. 2017. Effect of different thermal environments on the growth and development of wheat varieties for chhattisgarh plain. *Journal of Plant Development Sciences* Vol. 8 (12) : 573-581. 2016.
- Choudhary, A.K., Singh, A. and Yadav, D.S. 2010. "On farm testing" of wheat cultivars for site-specific assessment under varied bio-physical regimes in mid-hill conditions of Mandi district of Himachal Pradesh. *Journal of Community Mobilization and Sustainable Development* 5(1): 1-6.
- CIMMYT (International Maize and Wheat Improvement Center). 2013. WHEAT: CGIAR Research Program (CRP) led by CIMMYT. [<http://wheat.org/who-we-are/about-us>]
- Deo K, Mishra SR, Singh AK, Mishra AN, Singh A and Kumar S. 2017. Determination of suitable irrigation schedule for optimum water use efficiency of wheat crop. *Journal of Medicinal Plants Studies*. 5(3): 343-347.
- Farooq, O., M. Ali, M. Naeem, S. Abdul, M. Ijaz, A. Sher, T.A. Yasir and M.M. Iqbal, 2015. Impact of sowing time and planting method on the quality traits of wheat. *J. Glob. Innov. Agric. Soc. Sci.*, 3(1): 8-11.
- Gajbhiye M. 2021. Effect of IW/CPE ratio-based irrigation scheduling on weeds, growth and yield of wheat under different sowing dates in vertisol. M. Sc. Thesis. at Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur (M.P.).

Khande D., Agrawal K.K. and Bhan M. 2021. Phenology And Heat Unit requirement of Wheat under Various Thermal Environments and Different Methods of Sowing in Tawa Command Area of Madhya Pradesh. *Anveshak*. Vol. 51, No.2 (XV) July- December 2021.

Kumar P, Pannu RK, Khokhar SK. Effect of organic sources of nutrition and irrigation levels on growth and yield of wheat (*Triticum aestivum* L.). *Int. J Life Sci. Bt. & Pharm. Res* 2012;1(4):178-186.

Kumar S, Alam P and Ali N. 2013. Response of Wheat (*Triticum aestivum* L. emend. Fori. & Paol.) varieties to sowing dates. *Journal of Research (BAU)* 25(1): 56-59.

Malik GC, Wasim Iftikar, Banerjee M, Ghosh DC. Effect of Irrigation, Variety and Nitrogen on Growth and Productivity of Wheat (*Triticum aestivum* L.) *International Journal of Bio-resource and Stress Management* 2012;3(2):158-164.

Munir AT, Rahman A, Tawaha M. Impact of seeding rate, seeding date, rate and method of phosphorus application in faba bean (*Vicia faba* L.) in the absence of moisture stress. *Biotechnol, Agron. Soc. Environ.* 2002; 6(3):171-178.

Pal S, Kumar H, Gangwar HK, Singh A and Kumar P. 2020. Effect of scheduling irrigation based on IW/CPE ratio dry matter accumulation, yield attributes, yield and Economics of Wheat crop (*Triticum aestivum* L.) *Journal of Pharmacognosy and Phytochemistry* 2020; 9(4): 1946-1949.

Pal S, Kumar S, Kumar P, Singh A and Gangwar HK. 2020. Effect of Moisture Regime on IW/CPE Ratio on Soil Properties, Yield and Water use Efficiency of Wheat Crop (*Triticum aestivum* L.). *Int.J.Curr.Microbiol.App.Sci.* 9(3): 2499-2506.

Prasad S, Agrawal KK, Kumar R and Prakash V. 2017. Heat unit requirement of wheat cultivars under varying thermal regimes at Jabalpur. *Journal of Agrometeorology* 19(3): 283–285.

Prasad S, KK Agrawal KK and Kumar R. 2019. Productivity, profitability, quality and nutrient uptake of heat tolerant wheat (*Tricium aestivum*) cultivars as influenced by staggered sowing and nutrient levels. *Indian Journal of Agricultural Sciences* 89(4): 670-7.

Singh, A.K., Manibhushan N, Chandra and Bharati, R.C.2008. Suitable crop varieties for limited irrigated conditions in different agro climatic zones of India. *Int. J. Trop Agr.* 26 (3-4): 491-496.

Spiertz JHL, Hamer RJ, Xu H, Primo-Martin C, Don C. van der Putten, PEL Heat stress in wheat (*Triticum aestivum* L.): effect on grain growth and quality traits, *European Journal of Agronomy.* 2006; 23:89-95.

Sslunkhe R, Patil M and Deshmukh MM. 2012. Irrigation Scheduling (Response of wheat to IW/CPE). Ph.D. Thesis. Farm of Wheat Research Unit, Dr. Panjabrao Deshmukh Krishi Vidyapeeth, Akola.

Tanveer SK, Hussain I, Sohail M, Kissana NS, Abbas SG. Effect of different planting methods on yield and yield components of wheat Asian J. Plant Sci. 2003; 2:811-813.

Thorat 2015. Assessment of wheat productibility using CERES-wheat model as influenced by varieties, planting dates and irrigation in vertisols. Phd. Thesis. Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur.

Tripathi SC, Chander S and Meena RP. 2013. Effect of early sowing, N levels and seed rates on yield and yield attributes of different wheat (*Triticum aestivum*) varieties. Indian Journal of Agronomy 58 (1): 63-66.

Verma HP, Sharma OP, Kumar R, Yadav SS, Shivran AC and Balwan. 2017. Yield Attributes and Yield of Wheat (*Triticum Aestivum* L.) as Influenced by Irrigation Scheduling and Organic Manures. Chem Sci Rev Lett. 6(23), 1664-1669.

UNDER PEER REVIEW