

Original Research Article

Homemakers' attitude towards solid waste management in urban area of Hubli-Dharwad

Abstract

Proper waste management is a major concern of public health. It is necessary to create positive behavioural changes towards waste management among public in order to control diseases. Objectives were to determine the attitude towards solid waste management among the households of Hubli - Dharwad city in Karnataka. A community based cross sectional study was conducted in 67 wards of Hubli- Dharwad city located in Dharwad district of Karnataka state from January 2021 to December 2021. A total of 400 households were studied. Multistage sampling was done. Data was collected by direct interview using a prequestionnaire. Data was analyzed using SPSS software. The participants responsible for the waste management in household were women. Out of the 400 participants, 83%, 11% and 6% were Hindus, Muslims and Christians respectively. Majority of them (82%) had an educational qualification of high school and above. Most of them (82.5%) were housewives. About 47% of the study population belongs to upper middle class. Their major sources of information about solid waste management were SHGs. In the study, 93.8% of the study population had above average attitude and 6.2% had below average attitude. Almost 70% had the belief that government is not doing anything to fix the garbage problem. About 97%, 88.6% and 92% were willing to do composting, segregation and recycling of waste respectively. Majority of the participants had above average attitude towards household waste management. Continuous awareness programmes have to be conducted on safe waste disposal and efforts should be made to sustain the supervision of household waste management.

Keywords: Homemaker, Solid waste management, Attitude, SHGs (Self Help Groups).

Comment [Review1]: Write in full, the in brackets

1. Introduction

Solid waste may be defined internationally as the non-liquid waste materials from domestic, trade, commercial, industrial, agricultural and mining activities and from public services. Wastes arising from human and animal activities are normally solid and are discarded as useless or unwanted. The Resource Conservation and Recovery Act of 1976 (RCRA) defines solid waste to include garbage, refuse, sludge from municipal sewage treatment plants, ash from solid waste incinerators, mining waste, waste from construction and demolition and some hazardous wastes.

Comment [Review2]: Show the source of this information, Ref

Solid waste can be classified in terms of their original use (such as packaging waste), the material (glass, paper, or plastics), their physical properties (combustible or biodegradable), their origin (domestic, commercial, industrial or agricultural) and the safety parameters (hazardous/ radioactive). The household wastes can be classified as biodegradable and non-biodegradable waste. Biodegradable wastes are wastes like food waste that can be decomposed by biological processes. This should be composted at the community level. Non-biodegradable wastes are wastes like plastics, broken glass, etc. that cannot be decomposed, that can be segregated and sold or recycled. Waste segregation is the sorting out or separating out the biodegradable and non-biodegradable waste in to separate bins.

Comment [Review3]: Reduce space between words

Waste should be properly stored before disposal. Storage is the action of accumulating rubbish before disposal. There are different kinds of wastes like kitchen waste, garden waste, leather, rubber, glass, plastics, electronic waste, etc. E-waste is a common terminology used for electronic equipment such as TV, refrigerators, telephones, air conditioners, computers, mobile phones etc. that is not in use now by the current user. Hazardous wastes are those wastes which can cause a hazard to the ecological or environmental balance. These includes batteries, cooking oil, pesticides, fertilizers etc.

Comment [Review4]: Delete 's' to read 'include'

Waste management is the process of collecting, transporting, processing or disposing, managing and monitoring of waste materials. The term usually relates to materials produced by human activity and the process is generally undertaken to reduce their effect on health, and the environment. The 2011 Census of India estimates a population of 1.21 billion which is 17.66% of the world population. About 0.1 million tonnes of MSW is generated in India every day. The

data report indicates that 366 cities in India, which represent 70% of India's urban population were generating 31.6 million tons of waste in 2001 and are currently generating 47.3 million tonnes (2011), a 50% increase in one decade. In case of Karnataka, 11085 tons of solid waste is being produced every day

Comment [Review5]: Show the source of this information

Increasing population, urbanization, industrialization and changing consumption patterns are resulting in the generation of increasing amounts and different types of waste. There is a need to practice integrated solid waste management approach such as incorporation of more environmental and economic friendly concepts of source separation; recovery of waste; legitimization of the informal systems; partial privatization and public participation.

Wastes can be reduced by manufacturing products with less packaging, encouraging customers to bring their own reusable bags for packaging, encouraging the public to choose reusable products such as cloth napkins and reusable plastic and glass containers, backyard composting and sharing and donating any unwanted items rather than discarding them.

India has undertaken several waste management programmes. In 1999, the Government of India restructured the Comprehensive Rural Sanitation Programme and launched the Total Sanitation Campaign (TSC) which was later (on 1 April 2012) renamed Nirmal Bharat Abhiyan (NBA). The Swachh Bharat Mission was launched on 2nd October 2014 by Government of India.

Comment [Review6]: Separate

The purpose of this study is to determine the attitude of solid waste management among urban people. The findings can be utilized to alter the quality of environment and thus improve public health.

Objective of the study:

To determine the attitude towards solid waste management among the homemakers of Hubli-Dharwad in North Karnataka.

2. Materials and Methods

A community based cross sectional study was conducted in 67 wards of Hubli- Dharwad city in Karnataka state for one year from January 2021- December 2021. A total of 400 homemakers were selected from urban area by using random sampling method for the study. The consecutive houses were obtained from each of the 67 wards making the sample size of 400. Data was collected by direct interview using a pre-tested semi-structured questionnaire with two parts: Part 1 includes questions on socio-demographic variables such as age, address, religion, caste, head of the family, education and employment of the respondent, family income, type of family, number of family members. Part 2 includes questions on attitude of waste disposal. The questionnaire in English has been translated by a language expert in to Kannada and re-translated the Kannada version back to English and corrections were done accordingly. The person involved in that particular solid waste management was chosen for interview.

Attitude questions were scored. Maximum score obtainable in attitude was 156 and minimum was zero. Attitude was graded as below average (0-124) and above average positive attitude (125-156). Data was analysed with software SPSS 20. All results are expressed in frequencies and percentages

In this study, socio economic status (SES) was assessed using modified B.G Prasad (2016) scale. B.G. Prasad scale is based on the per capita income of an individual. It classified the status into five classes.

3. Results and Discussion

Table 1: Age wise distribution of the study population (N=400)

Age group (in years)	Frequency	Percentage (%)
20-29	27	6.8
30-39	113	28.2
40-49	124	31.0
50-59	78	19.5
60-69	49	12.2
70-79	9	2.2
Total	400	100.0

Table 1 shows that out of 400 homemakers, majority were in the age group 40-49 years (31%). The mean age of the study population was 45.16 ± 11.61 years. This was almost similar to the study done by Mahima and Lavanya (2016), belong to the age group of 30-50 years (46.8%).

Table 2: Religion wise distribution of the study population (N=400)

Religion	Frequency	Percentage (%)
Hindu	332	83.0
Muslim	44	11.0
Christian	24	6.0
Total	400	100.0

Table 2 shows that, 83% of the study participants belong to Hindu religion. According to Mahima and Lavanya (2016), 83% belonged to Hindu religion followed by Muslim (11%) and christian religion (6%).




Figure 1: Distribution by the type of family (N=400)




Figure 2: Distribution by size of the family (N=400)

Figure 1 shows that 70% of the study population belongs to the nuclear family. Figure 2 shows that 57% of the study population belongs to a family size of less than or equal to 4 members and 41% belong to family size of 5-8 members. These findings are consistent with the NFHS 4 report, 54.8% of the households are with 1-4 numbers of family members, 45.2% with 5-8 members and 4.9% with more than 9 members [Health NF. India(2019)].

Table 3: Socio-demographic factors of the study population (N=400)

Education	Frequency	%
Education		
Primary school	30	7.5
Middle school	42	10.5
High school	191	47.8
intermediate school/ Post diploma	119	29.8
Graduate	18	4.5
Total	400	100.0
Occupation		
Unemployed/Housewife	330	82.5
Unskilled	45	11.2
Semi-skilled	11	2.8
skilled	6	1.5
Clerical/shop owner/ Farmer	5	1.2
Semi professional	3	0.8
Total	400	100.0

Table 3 shows that majority (47.8%) of the study population had High school education. There were no illiterate in the study population. Majority (82.5%) of the participants were housewives. According to NFHS-4 report of Kerala, 28.7% of the women have completed 12 or more years of schooling, 19% completed 10-11 years, 34% completed 5-9 years of schooling whereas, 4.2% of women have not done schooling. In the present study, 18% of the women were employed which is similar to the NFHS report Kerala, where 21% among the women in the age group of 15 to 49 years were employed.




Figure 3: Socio-demographic status* of the study population (N=400)

Figure 3 shows that majority (47%) of the study population belongs to upper middle class and 3.5% belong to lower class. In the study, majority of the houses have concrete ceiling (87.8%) followed by tiled (10%), kutcha ceiling (2%) and asbestos (0.2%). Around 97% of them have their own house and 3% live in rented house. Based on updated B G Prasad classification (2016) of socio-economic status, 10% of the study population belonged to upper class, 47% belonged to upper middle class and very few belonged to lower class (3.5%). In NFHS 4 report Kerala, based on wealth quintile, 48% of the households were in the highest quintile and 0.5% in the lowest wealth quintile. Health NF. India (2019), majority of the houses have concrete ceiling (87.8%) followed by tiled (10%), kutcha ceiling (2%) and asbestos (0.2%). Around 97% of them have their own house and 3% have rented house. The NFHS 4 report India found that 89% of households have pucca houses, 0.4% kachha house and 10.5 percent have semi-puccahouses.

Comment [Review7]: Show the sources of these data.

Table 4: Attitude towards solid waste management

	Strongly agree	Agree	No opinion	Disagree	Strongly disagree
A. Solid waste management	N (%)	N (%)	N (%)	N (%)	N (%)
A1. waste management awareness class should be held in community	400 (100)				
A2. Environmental education should be taught in school	300 (75)	81 (20.2)	18(4.5)	1 (0.2)	
A3. Household should be clean and free of waste	393 (98.2)	5 (1.2)	1 (0.2)		1 (0.2)
A4. I should play an important role in reducing household waste generation	380 (95)	18 (4.5)	1 (0.2)	1 (0.2)	
A5. The purchase decision I make can increase or decrease the amount of garbage my household must get rid off	238 (59.5)	134 (33.5)	7 (1.8)	4 (1)	17 (4.2)
A6. I don't think that burning garbage can be bad for my health and others health	52 (13)	15 (3.8)	8 (2)	12 (3)	313 (78.2)
A7. people throw garbage in streets as they have no other means of getting rid of garbage	47 (11.8)	106 (26.5)	22 (5.5)	27 (6.8)	198 (49.5)
A8. local self government is not doing enough to fix the garbage problem	109 (27.2)	172 (43)	25 (6.2)	36 (9)	58 (14.5)
A9. Regular collection of garbage is only solution to garbage problem	280 (70)	93 (23.2)	12 (3)	8 (2)	7 (1.8)
A10. Generated waste can be managed at source/household itself	254 (63.5)	110 (27.5)	14 (3.5)	11 (2.8)	11 (2.8)
B. Willingness to participate					
B1. composting	291 (72.8)	97 (24.2)	10 (2.5)		2(0.5)
B2. recycling	206 (51.5)	163 (40.8)	19 (4.8)	4 (1)	8(2)
B3. Willingness to segregate materials for collection	199 (49.8)	155 (38.8)	31 (7.8)	9 (2.2)	6(1.5)
B4. Willingness to pay for pick up for recyclable materials	222 (55.5)	148 (37)	20 (5)	3 (0.8)	7(1.8)
B5. willingness to give away plastic bottles for recycling	232 (58)	116 (29)	26 (6.5)	7 (1.8)	19(4.8)
B6. willingness to buy lesser amount of throwaway products	263 (65.8)	126 (31.5)	5 (1.2)		6(1.5)
B7. willingness to gather more information on reduction of garbage	320 (80)	67 (16.8)	8 (2)	1 (0.2)	4(1)
C. Composting					
C1. takes up a lot of time	175 (43.8)	192 (48)	6 (1.5)	10 (2.5)	17 (4.2)
C2. takes a lot of effort	157 (39.2)	203 (50.8)	6 (1.5)	13 (3.2)	21 (5.2)
C3. demand high degree of technical knowledge	173 (43.2)	190 (47.5)	8 (2)	14 (3.5)	15 (3.8)
C4. requires lot of space	72 (18)	195 (48.8)	12 (3)	35 (8.8)	86 (21.5)
C5. not worthwhile unless there is lot of waste	95 (23.8)	158 (39.5)	9 (2.2)	25(6.2)	113 (28.2)
C6. compost pits attracts insects and rodents	137 (34.2)	168 (42)	5 (1.2)	22 (5.5)	68 (17)
C7. compost pits are unsightly	98 (24.5)	160 (40)	11 (2.8)	36 (9)	95 (23.8)
D. Issues for concern	Concerned	Not concerned		No opinion	
	N (%)	N (%)		N (%)	
E1. health risk is related to burning/dumping garbage	400 (100)				
E2. Illegal dumping polluting water body	400 (100)				
E3. Diseases related to improper storage and disposal	400 (100)				
E4. Flooding due to garbage blocking drains	398 (99.5)	1(0.2)		1 (0.2)	

and gullies			
E5. Litters/illegal dumping	398 (99.5)	2 (0.5)	

Table 4 shows that cent per cent of homemakers strongly agreed to the statement that waste management awareness class should be held in community, only 75% strongly agreed to have environment education in school. About 95% strongly agreed and 18% agreed that they should play an important role in reducing household waste generation. Out of the total, 59.5%, 33.5% and 4.2% strongly agreed, agreed and disagreed with the statement that the purchase decision they make can increase or decrease the amount of garbage the household must get rid off. Similarly, by Davies *et al.* (2016), 78% agreed that public education was appropriate to tackle solid waste problems, 88% agreed that there should be waste management education in school curriculum. According to Banga (2013), 63.5% strongly agreed, 27.5% agreed and 2.8% strongly disagreed to the statement that generated waste could be managed at the source itself. Around 27% strongly agreed, 43% agreed and 14% strongly disagreed that local self-government was not doing enough to fix the garbage problem. Although half (49.5%) of the participants strongly disagreed that people throw garbage in streets as they have no other means, there were 26.5% who agreed that people have no other means for garbage disposal.

Table 5: Assessment of attitude towards solid waste management (N=400)

Grading	Frequency	Percentage (%)
Above average	375	93.8
Below average	25	6.2
Total	400	100.0

Table 5 shows that 93.8% of the study population had above average attitude towards household waste management and 6.2% had below average attitude. Similarly, in a study done by Duruet *al.* (2017) 55.4%, 38.6% and 6% of the participants had moderate, good and poor level of attitude towards environment sanitation respectively.

Table 6: Two-way ANOVA test for homemakers' attitude towards the solid waste management based on their age, education and occupation

Source of Variation	Sum of Squares	Degree of Freedom	Mean Squares	F-value	P-value
Age	945.25	2	472.63	3.828**	
Attitude towards SWM	3028.5	1	3028.5	24.53**	0.008
Error	246.9	2	123.45		
Total	4220.66	5			
Education	120.49	2	60.245	54.031**	
Attitude towards SWM	1432.22	1	1432.22	1284.5**	0.006
Error	2.23	2	1.115		
Total	1554.94	5			
Occupation	26261.08	2	13130.54	61.587**	
Attitude towards SWM	118244.9	1	118244.9	554.62**	0.002
Error	426.4033	2	213.2017		
Total	144932.4	5			

** Significant at $P < 0.01$

Table 6 shows the results obtained due to the application of two-way Analysis of Variance (ANOVA) test for homemakers' attitude towards the solid waste management based on their age, education and occupation. The F values computed as 3.828, 54.031 and 61.587 for homemakers' age, education and occupation respectively due to variation in their attitude towards the solid waste management are found to be highly significant (at $P < 0.01$). This indicates that there exists a significant variation among the different parameters due to variation in their attitude towards the solid waste management. Further, the F values computed as 24.53, 1284.5 and 554.62 for homemakers' attitude towards the solid waste management due to variation in their age, education and occupation respectively are also found to be highly significant (at $P < 0.01$). This also shows a significant variation in homemakers' attitude towards the solid waste management due to variation in their age, education and occupation.

4. Conclusion

A community based cross sectional study was conducted in Hubli – Dharwad city of Dharwad district in Karnataka to assess the Knowledge, Attitude and Practice of solid waste management among the homemakers and their mean age was 45.16 ± 11.61 years.

Out of the 400 participants, 83%, 11% and 6% were Hindus, Muslims and Christians respectively. Majority of them (82%) had an educational qualification of high school and above. Most of them (82.5%) were housewives. About 47% of the study population belongs to upper middle class.

In the study, 93.8% of the study population had above average attitude and 6.2% had below average attitude. All of them strongly agreed to have an awareness class on waste management in the community. Almost 70% had the belief that government is not doing anything to fix the garbage problem. About 97%, 88.6% and 92% were willing to do composting, segregation and recycling of waste respectively. They were all very much concerned about the illegal waste dumping and diseases related to improper waste storage and disposal.

References

- AP, Nkwocha IB. Assessment of Waste Management Practices among Residents of Owerri Municipal Imo State Nigeria. *J Environ Prot (Irvine, Calif)*. 2015;6(5):446–56.
- Ashalakshmi KS, Arunachalam P. Solid Waste Management: A Case Study of ArppukaraGramapanchayat Of Kottayam District, Kerala (India). *J Glob Econ*. 2010;6(1):33–63.
- Available from: shodhganga.inflibnet.ac.in/bitstream/10603/1462/14/14_chapter-5.pdf. Accessed on 15 January 2019.
- Banga M. Household Knowledge, Attitudes and Practices in Solid Waste Segregation and Recycling: The Case of Urban Kampala Recycling: The Case of Urban Kampala. *Zambia SocSci J*. 2013;2(1):27–39.
- Davies NT, Kudzai NS. The Usefulness of Including Women in Household Solid Waste Management. A Case Study of Dzivaresekwa High Density Suburb; Harare. *IOSR J HumanitSocSciVer II*. 2016;21(3):92–108.
- Devi S, Perry N, Gharde P, Yangad R, Nain S, Luckose K. A study to assess the attitude and concern regarding household solid waste management among the residents of Pune city. *Int J Recent Sci Res*. 2017;8(7).
- Dhanalakshmi T. Willingness to pay (wtp) for aerobic treatment of municipal solid waste: a study at Alappuzha, Kerala. 2015;1(14):384-9.
- Duru CB, Iwu AC, Diwe KC, Uwakwe KA, Merenu IA, Madubueze UC, et al. Environmental Sanitation Practices: A Case Study of Solid Waste Management in Semi-Urban Communities in Orlu, Imo State Nigeria. *Occup Dis Environ Med*. 2017, 05(04):88–105.
- Excreta disposal. Available from: http://www.who.int/water_sanitation_health/hygiene/settings/hvchap4.pdf. Accessed on 15 January 2019.
- Harikrishnan G. Solid Waste Management: a Comparative Study Between Kerala and Tamilnadu. *Int J Res SocSciHumanit*. 2014.
- Health NF. India. Available from: <http://rchiips.org/NFHS/nfhs4.shtml>. Accessed on 15 January 2019.

- Health NF. Kerala. 2015; Available from: <http://rchiips.org/NFHS/NFHS-4Reports/Kerala.pdf>. Accessed on 15 January 2019.
- Indhira K, Senthil J, Vadivel S, Appl A, Res S. Awareness and attitudes of people perception towards to household solid waste disposal :Kumbakonam Town, Tamil Nadu, India. Arch ApplSci Res. 2015;7(3):6–12.
- Kiran KG, Kini S, Ravi K. KAP study of solid waste disposal of households in Kuttar&ManjanadiPanchayath covered under gramashemaprogramme of K.S. Hegde Medical Academy. NUJHS Sept NitteUniv J Heal Sci. 2015;5(3):2249–7110.
- LaminouManzo O, Saidou H, Illiassou SA, Idrissa ST, LaminouManzo O. Assessment of Domestic Wastewater Management Practices in the Communal District I of Maradi City, Niger Republic Keywords Domestic Wastewater, Sanitation, Communal District I of Maradi, Management Practices. J Geosci Environ Prot. 2015;3(3):57–65.
- Licy CD, Vivek R, Saritha K, Anies TK, Josphina CT. Awareness, Attitude and Practice of School Students towards Household Waste Management. J Environ. 2013;2(6):147–50.
- Lutui V. Waste management practices, perceptions and attitudes in Tonga. 2001; Available from: <http://ro.uow.edu.au/theses/2897>. Accessed on 15 January 2019.
- Mahima S, Lavanya VL. A study on problems of solid waste management with special reference to Palakkad Municipality, Kerala, India.Nt Res J Soc Sci. 2016;5(9):1–6.
- Nava Kerala mission in Kerala - PradhanMantriYojana Schemes. Available at: <http://www.pradhanmantrijojana.co.in/nava-keralamission-kerala/>. Accessed on 15 January 2019.
- Population and Municipal Solid Waste Generation in India. Available at: http://nswaienvs.nic.in/pdf_FF/Population and Municipal Solid Waste Generation in India.pdf. Accessed on 15 January 2019.
- Preissler A, Chaves L, Bitencourt R, Silva D. Environmental diagnosis of hazardous household wastes and the family health strategy as liaison for implementation of a management program in the South of Brazil. 2015;23(2):109–17.

Swachh Bharat Abhiyan Report Stats. 2016. Available at: <http://sbm.gov.in/sbmreport/home.aspx>. Accessed on 15 January 2019.

UNICEF. Solid and liquid waste management in rural areas. A Tech note. 2012; Available from: www.ddws.gov.in. Accessed on 15 January 2019.

Vasudevan J, Mishra AK, Singh Z. An update on B. G. Prasad's socioeconomic scale: May 2016. *Int J Res Med Sci*. 2016;4(9):4183–6.

UNDER PEER REVIEW