

Original Research Article

Influence of foliar spray of Brassinosteroids (BR), Salicylic acid (SA) and Gibberellic acid (GA₃) on vegetative growth and flowering parameters of Cucumber (*Cucumis sativus* L) cv. Arpit.

Abstract

An investigation entitled “Influence of foliar spray of Brassinosteroids (BR), Salicylic acid (SA) and Gibberellic acid (GA₃) on vegetative and flowering parameters of Cucumber (*Cucumis sativus* L) cv. Arpit” was carried out at the Vegetable Research Farm, Department of Horticulture, Sam Higginbottom University of Agriculture, Technology & Science, Prayagraj, (U.P.) - 211007 during the year 2018-19 and 2019-2020. The experiment was laid out in a Randomized Block Design with 15 treatments and 3 replications and each replicated thrice. Source of variables were Brassinosteroids (BR) 0.05 PPM, Brassinosteroids (BR) 0.10 PPM, Brassinosteroids (BR) 0.5 PPM, Brassinosteroids (BR) 1.0 PPM, Brassinosteroids (BR) 2.0 PPM, Salicylic acid (SA) 0.05 mM, Salicylic acid (SA) 0.10mM, Salicylic acid (SA) 0.2mM, Salicylic acid (SA) 0.5mM, Salicylic acid (SA) 1.0mM, Gibberellic acid (GA₃) 25 PPM, Gibberellic acid(GA₃) 50 PPM, Gibberellic acid(GA₃) 100 PPM, Gibberellic acid (GA₃) 150 PPM and combination of them with control treatment taken as 15 treatments. As far as the vegetative growth and flowering is concerned, application of different treatment of plant growth regulators significantly enhanced the Length of main vine (cm), Number of branches per plant, Number of leaves per plant, Diameter of main stem (cm), Inter-nodal distance (cm), Leaf area (cm²), Days to first flower bud initiate (DAS), Days taken to 50% flowering, Number of male flower per plant, Number of female flower per plant at all successive stages. The maximum vine length (143.17 and 146.04cm), number of leaves per plant (93.74 and 94.74), number of branches per plant (19.49 and 19.82), diameter of main stem (14.70 and 16.02cm), leaf area (277.42 and 280.90 cm²), internodal distance (7.27 and 6.97cm) and as per flowering parameters lowest days taken to first flower bud initiation (36.66 and 35.73 days), days taken to 50% flowering (42.40 and 42.73 days), number of male flower per plant (91.71 and 92.05) and number of female flower per plant (47.59 and 46.92) were recorded in both successive year with T₅ (Brassinosteroids (BR) 2.0 PPM) and minimum growth and flowering were observed in T₀ (control).

Keywords- Cucumber, Vegetative growth, Flowering, Brassinosteroids, Salicylic acid, Gibberellic acid.

Introduction

Cucumber (*Cucumis sativus* L.) is a cross pollinated and widely grown vegetable crop in the Cucurbitaceae family, with chromosome number $2n=14$. It is an indigenous vegetable to India (**De Candole, 1967**). It is often a monoecious, annual, trailing or climbing vine with hirsute or scabrous stems and triangular ovate leaves with shallow and sharp sinuses (**Bailey, 1969**). At the leaf axils, unbranched lateral tendrils appeared. Flower clusters form in leaf axils as the lateral branches grow (**Ahmed et al. 2004**). It is a warm-season crop with minimal to no frost resistance. Cucumber growth and development are aided by temperatures over 20-30°C.

It is widely farmed in the Indian states of Madhya Pradesh, Tamil Nadu, Uttar Pradesh, Andhra Pradesh, Kerala, and Maharashtra. It covers an area of 41 million ha in India, produces 641 MT, and has a productivity of 15.63 t/ha (**Handbook of Horticulture Statistics 2019-20**).

Depending on the cultivar, area, and soil climate, the plant begins flowering early and produces marketable fruits within two or three months. Flowering is a critical stage in the development of cucurbits since it determines fruiting and yield. Cucumber is a monoecious plant, which means that the first flowers that develop near the base of a cucumber plant are male. A week after the male flower initiation, the female flowers develop, with the little cucumber fruit at the base (**Bantoc, 1964**). Cucumbers have a high water content while being low in calories, fat, cholesterol, and salt. The fruit is an elongated, round triangular fake berry or pepo. Its size, shape, and colour differ according on the cultivar. Fruits are beneficial to persons who have jaundice, constipation, or indigestion. It is eaten raw with salt and pepper, or as part of a salad with pickles. The fruit pulp is used to make mash cakes.

Cucumber's reaction to plant growth regulators displays a remarkable spectrum of floral morphology, including staminate, pistillate, and hermaphrodite flowers that appear in a variety of configurations and provide a variety of sexual expression. Growth regulators have a huge impact on sex expression and blooming in cucumber crops, either suppressing male flowers or increasing the amount of female flowers (**Al-Masoum & Al-Masri, 1999**) with no negative consequences for the environment or human health. Plant growth

regulators are also used to control cucumber plant vegetative development, which increases plant population per unit area in terms of production (Latimer, 1991).

Brassinosteroids are present in a variety of species ranging from lower plants to higher plants. They are steroidal plant hormones that help plants grow and develop. BR may show a keen interest in the role of horticulture crops. It was characterised by a decrease in the quantity of male flowers in the early stages of growth and a promotion of female flower initiation in the main stalk. Another study on cucumber found that BS is critical during early fruit development (Fu et al., 2008). Salicylic acid (SA) is a phytohormone that has recently been added to the family of phytohormones for appropriate plant growth development and induction of tolerance to both biotic and abiotic stresses. SA is a phenolic endogenous growth regulator that regulates various physiological processes in agricultural plants, including stomatal closure, ion uptake, ethylene biosynthesis inhibition, and transpiration (Khan et al., 2003 and Shakirova et al., 2003).

Gibberellic acid is important in seed germination, endosperm mobilisation, stem elongation, leaf expansion, shortening maturation time, and boosting flower and fruit set and composition (Roy & Nasiruddin 2011). GA₃ slows senescence, promotes chloroplast growth and development, and increases photosynthetic efficiency, perhaps leading to enhanced production (Yuan & Xu 2001).

Materials and Methods

The experiment entitled “Influence of foliar spray of Brassinosteroids (BR), Salicylic acid (SA) and Gibberellic acid (GA₃) on vegetative and flowering parameters of Cucumber (*Cucumis sativus* L) cv. Arpit” was carried out during the summer season of the year 2018-2019 and 2019-20. The experiment was laid out in a Randomized Block Design with 15 treatments and 3 replications and each replicated thrice. Department of Horticulture, Naini Agricultural Institute, Sam Higginbottom University of Agriculture Technology and Sciences, Prayagraj, provided all of the necessary equipment for the experiment. It reaching a height of 98 meters above the sea level and is located at 25°8'N latitude and 81°50' E longitude. Prayagraj district is situated in Uttar Pradesh's subtropical zone, which has extreme heat summers and mild winters. The area's peak temperature is 46°C–48°C, and it seldom goes below 4°C–5°C. The relative humidity level varies between 20 and 94 percent. The annual average rainfall in this area is 1013.4 mm.

Comment [RY1]: THIS PART SHOULD BE EXPLORED MORE because BE, SA, and GA₃ are the main object in this experiment. Give more example of the important of these substances

Comment [RY2]: How about the statistical analysis (ANOVA, mean separation test after anova LSD, HSD or others>)

Source of variable were T₀ (Control), T₁ Brassinosteroids 0.05 PPM, T₂ Brassinosteroids 0.10PPM, T₃ Brassinosteroids 0.5 PPM, T₄ Brassinosteroids 1.0 PPM, T₅ Brassinosteroids 2.0PPM, T₆ Salicylic acid 0.05 mM, T₇ Salicylic acid 0.10 mM, T₈ Salicylic acid 0.2 mM, T₉ Salicylic acid 0.5 mM, T₁₀ Salicylic acid 1.0mM, T₁₁ Gibberellic acid 25 PPM, T₁₂ Gibberellic acid 50 PPM, T₁₃ Gibberellic acid 100 PPM, T₁₄ Gibberellic acid 150 PPM. The observations were recorded on vegetative growth and flowering parameters of cucumber as Vine length (cm), No. of leaves /plant, No. of branches /plant, Diameter of main stem (mm), Internodel distance (cm), Leaf area (cm²), Days to first flower bud initiation, Days taken to 50% flowering, No. of male flower/plant, No. of female flower/plant were studied in the investigation.

Result and Discussion

The data on various observations recorded during experimentation were subjected to statistical analysis in Randomized Block Design in order to find out the significance of different treatments by using the analysis of variance. The results have been integrated along with the corresponding tables. In terms of vegetative parameters, it was found that Vine length, number of leaves, number of branches, diameter of main stem(mm), Internodal distance(cm) and Leaf area(cm²) were significantly increased during the both year by different treatments of plant growth regulators at all successive stage of growth.

It is clear from the table 1 that the maximum vine length at 90 days (143.17 cm and 146.04 cm in 2018-19 and 2019-20 respectively and 144.60 cm as pooled) closely conformity to the **Ramani et al.,(2016), kumar & kaur (2019) and Hirpara et al., (2014)** Maximum Number of leaves at 90 days (93.74 and 94.74 in 2018-19 and 2019-20 respectively and 94.24 as pooled) closely associated with **Irfan et al., (2017) and Vishnu ritti et al., (2019)**; Number of branches (19.49 and 19.82 in 2018-19 and 2019-20 respectively and 94.24 as pooled) associated with **Ghosh et al.,(2020)**; maximum Diameter of main stem (14.70 mm and 16.02 mm in 2018-19 and 2019-20 respectively and 15.36 mm when pooled) closely with **Vishnu ritti et al., (2019)**; and maximum leaf area (277.42 cm² and 280.90 cm² in 2018-19 and 2019-20 respectively and 279.16 cm² when pooled) close conformity to the **karuppaiah et al., (2019) and khatoon et al., (2020)** and maximum Internodal distance (7.27 and 6.97 in 2018-19 and 2019-20 respectively and 7.12 when pooled) closely associated with **khatoon et al., (2021) and Acharya et al., (2020)** were recorded in T₅ (Brassinosteroids

(BR) 2.0 PPM). While minimum Vine length at 90 days (126.62cm, 128.95cm and 127.79cm in 2018-19; 2019-20 and as pooled, respectively), number of leaves (74.53, 75.19 and 74.86 in 2018-19; 2019-20 and as pooled, respectively), Number of branches (12.35, 13.68 and 13.02 in 2018-19, 2019-20 and pooled, respectively), Diameter of main stem (11.05 mm, 11.71mm and 11.38 mm in 2018-19, 2019-20 and pooled, respectively), leaf area (258.07cm², 266.07 cm² and 262.07 cm² in 2018-19, 2019-20 and pooled, respectively), Internodal distance (4.79, 4.39 and 4.59 in 2018-19, 2019-20 and pooled, respectively) were observed with T₀ (Control).

In term of Flowering parameters, it was found that Days to first flower bud initiate (DAS), Days taken to 50 % flowering, Number of male flower per plant, Number of female flower per plant were significantly increased by different treatments of plant growth regulator at all successive stages of growth and flowering in both the individual years and when pooled.

Minimum Days to first flower bud initiate (days) (36.66 days and 35.73 days in 2018-19 and 2019-20 respectively and 36.20 days as pooled) closely associated with **Abdul et al., (2018)**. Minimum Days to 50% flowering (days) (42.40 days and 42.73 days in 2018-19 and 2019-20 respectively and 42.57 days as pooled) closely associated with **karuppaiah et al., (2019) and khatoon et al., (2020)** and Maximum Number of male flower per plant (91.71 and 92.05 in 2018-19 and 2019-20 respectively and 91.88 as pooled) close conformity with the **Khan et al., (2006) & Kadi et al., (2018)** Maximum Number of female flower per plant (47.59 and 46.92 in 2018-19 and 2019-20 respectively and 47.26 as pooled) closely associated with **khatoon et al., (2021), singh et al.,(2010) and pawar et al.,(2019)** were recorded in T₅ (Brassinosteroids (BR) 2.0 PPM). While maximum days to first flower bud initiate (days) (41.40 days and 41.26 days in 2018-19 and 2019-20 respectively and 41.33 days as pooled), maximum days to 50% flowering (days) (46.26 days and 46.53 days in 2018-19 and 2019-20 respectively and 46.40 days as pooled) associated with **Ghorbani et al., (2017)** and minimum Number of male flower per plant (76.41 and 72.07 in 2018-19 and 2019-20 respectively and 74.24 as pooled), minimum Number of female flower per plant (36.32 and 34.65 in 2018-19 and 2019-20 respectively and 35.49 as pooled) were observed with T₀ (Control).

It is determined that the treatment T₅ (Brassinosteroids (BR) 2.0 PPM) was found optimal in terms of Vegetative development, flowering parameters, and cucumber production parameters based on the current experiment in both subsequent years

2018-19 and 2019-20. Following a few more conjunctive studies, farmers may be advised to use those Plant growth regulator combinations.

Table 1:- Influence of foliar spray of Brassinosteroids (BR), Salicylic acid (SA) and Gibberellic acid (GA₃) on vegetative parameters of Cucumber cv. F₁ hybrid Arpit.

Comment [RY3]: Are there any statistical analysis?

Treatment	Vine length 90 days			Number of leaves per plant 90 days			Number of branches per plant 90 days			Diameter of main stem (mm) 90 days			Leaf area (cm ²) 90 days		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled
T0	126.62	128.95	127.79	74.53	75.19	74.86	12.35	13.68	13.02	11.05	11.71	11.38	258.07	266.07	262.07
T1	133.91	136.57	135.24	84.57	85.49	85.03	14.81	15.92	15.37	12.64	13.40	13.02	264.43	272.70	268.57
T2	134.57	137.19	135.88	85.28	87.30	86.29	15.24	16.66	15.95	12.99	13.77	13.38	265.60	273.35	269.48
T3	139.39	141.31	140.35	90.19	90.89	90.54	17.33	18.53	17.93	13.47	14.28	13.87	270.87	277.93	274.40
T4	142.00	144.36	143.18	91.34	93.34	92.34	18.10	18.89	18.50	13.85	15.10	14.47	274.23	279.22	276.73
T5	143.17	146.04	144.60	93.74	94.74	94.24	19.49	19.82	19.66	14.70	16.02	15.36	277.42	280.90	279.16
T6	129.65	131.99	130.82	79.79	81.45	80.62	13.74	14.46	14.10	11.73	12.44	12.09	260.36	269.68	265.02
T7	133.56	135.58	134.57	81.15	83.93	82.54	14.39	15.51	14.95	12.40	13.14	12.77	262.35	270.76	266.56
T8	137.08	139.83	138.45	87.65	89.15	88.40	16.08	17.76	16.92	13.15	13.94	13.54	268.35	276.60	272.48
T9	138.10	140.74	139.42	89.99	90.67	90.33	16.84	18.47	17.66	13.33	14.13	13.73	269.43	277.75	273.59
T10	139.96	143.15	141.56	90.33	91.10	90.72	17.58	18.64	18.11	13.56	14.64	14.10	272.73	278.24	275.49
T11	136.52	139.08	137.80	86.01	88.89	87.45	15.76	17.17	16.47	13.07	13.85	13.46	267.68	274.43	271.06
T12	137.78	140.15	138.97	89.25	90.33	89.79	16.33	18.33	17.33	13.25	14.05	13.65	269.22	277.38	273.30
T13	140.31	143.54	141.93	90.60	91.52	91.06	17.75	18.76	18.26	13.67	14.76	14.22	273.38	279.02	276.20
T14	142.49	145.12	143.80	92.43	94.01	93.22	18.69	19.17	18.93	14.13	15.26	14.70	274.91	280.75	277.83
CD value	6.10	6.22	6.16	3.88	3.94	3.91	0.73	0.78	0.75	0.58	0.62	0.60	8.59	5.50	6.58
S.Ed (±)	2.98	3.03	3.01	1.89	1.92	1.91	0.35	0.38	0.37	0.29	0.30	0.30	4.19	2.68	3.21

Table 2:- Influence of foliar spray of Brassinosteroids (BR), Salicylic acid (SA) and Gibberellic acid (GA₃) on flowering parameters of Cucumber cv. F₁ hybrid Arpit.

Comment [RY4]: Statistical analysis?

Treatment	Internodal distance (mm)			Days taken to first flower bud initiation (days)			Days taken to 50% flowering (days)			Number of male flower per plant			Number of female flower per plant		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled
T0	4.79	4.39	4.59	41.40	41.26	41.33	46.26	46.53	46.40	76.41	72.07	74.24	36.32	34.65	35.49
T1	5.18	4.96	5.07	40.33	40.00	40.17	45.86	45.38	45.62	84.65	84.27	84.46	41.65	40.20	40.93
T2	5.29	5.01	5.15	40.26	39.80	40.03	45.72	45.13	45.43	85.08	84.91	85.00	41.75	40.41	41.08
T3	6.20	6.11	6.16	38.86	38.53	38.70	44.73	44.63	44.68	87.90	86.18	87.04	43.38	42.38	42.88
T4	6.78	6.38	6.58	38.43	38.30	38.36	44.46	43.73	44.10	89.89	88.36	89.13	45.01	42.74	43.88
T5	7.27	6.97	7.12	36.66	35.73	36.20	42.40	42.73	42.57	91.71	92.05	91.88	47.59	46.92	47.26
T6	5.12	4.78	4.95	41.00	40.33	40.67	46.20	46.20	46.20	81.48	80.48	80.98	40.64	38.64	39.64
T7	5.13	4.85	4.99	40.66	40.23	40.45	45.86	46.09	45.98	83.27	83.74	83.51	40.87	40.18	40.53
T8	5.38	5.13	5.26	39.93	39.06	39.50	45.06	44.86	44.96	86.18	85.65	85.92	42.75	41.40	42.08
T9	6.11	5.78	5.95	38.61	38.68	38.65	44.86	44.73	44.80	87.72	85.85	86.79	43.07	42.30	42.69
T10	6.50	6.13	6.32	38.43	38.53	38.48	44.63	44.33	44.48	88.85	87.72	88.29	44.63	42.41	43.52
T11	5.35	5.05	5.20	40.06	39.33	39.70	45.53	44.94	45.24	85.30	85.64	85.47	42.31	40.65	41.48
T12	5.78	5.53	5.66	39.53	38.93	39.23	44.94	44.75	44.85	86.99	85.66	86.33	42.85	41.65	42.25
T13	6.71	6.27	6.49	38.42	38.19	38.31	44.53	44.27	44.40	89.69	87.90	88.80	44.97	42.63	43.80
T14	6.97	6.50	6.74	38.13	37.86	38.00	44.00	43.65	43.83	90.24	90.23	90.24	47.07	43.43	45.25
CD value	0.26	0.25	0.26	1.87	1.75	1.76	2.01	2.00	2.01	3.85	3.80	3.82	1.91	1.84	1.88
S.Ed (±)	0.13	0.12	0.13	0.91	0.85	0.86	0.98	0.98	0.98	1.88	1.86	1.87	0.93	0.90	0.92

Comment [RY5]: No C

Comment [RY6]: NO CONCLUSION?

References

- Al-Masoum, A.A. and A.A. Al-Masri. 1999.** Effect of ethephon on flowering and yield of monoecious cucumber. *Egyptian J. Horti.*, 26: 229–236.
- Ahmed, M., Hamid, A. and Akbar, Z. (2004).** Growth and Yield Performance of Six Cucumber (*Cucumis sativus* L.) Cultivars Under Agro-Climatic Conditions of Rawalakot, Azad Jammu and Kashmir. *Int. J. Agri. Biol.*, 6(2):396- 399.
- Ajay S. Kadi, K.P. Asati, Swati Barche and Tulasigeri R. G. (2018)** Effect of Different Plant Growth Regulators on Growth, Yield and Quality Parameters in Cucumber (*Cucumis sativus* L.) under Polyhouse Condition. *Int. J. Curr. Microbiol. App. Sci.* (2018) 7(4): 3339-3352
- Abdul Baqi, R Krishna Manohar and AG Shankar (2018)** Effect of GA3 and NAA with pruning levels on growth, sex expression and yield attributes of cucumber (*Cucumis sativus* L. Malini F₁) under protected condition. *International Journal of Chemical Studies* 2018; 6(4): 1991-1996.
- Acharya S. K., Chirag Thakar, Brahmhatt J. H. and Nikunj Joshi (2020)** Effect of plant growth regulators on cucurbits: A review. *Journal of Pharmacognosy and Phytochemistry* 2020; 9(4): 540-544.
- Bailey, L.H. (1969).** Manual of Cultivated Plants. Macmillan Company, New York., : 1116.

- Bantoc, G.B. JR. (1964).** Further studies on sex expression and sex ratio in cucumber as affected by plant regular spray. *Indian Journal. Hon.* 17:210- 216.
- De Candole, A. (1967).**Origin of Cultivated Plants. Hafnar Publishing Co., New York., : 264.
- Feng Qing Fu, Wei Hua Mao, Kai Shi, Yan Hong Zhou, Tadao Asami and Jing Quan Yu (2008)** A role of brassinosteroids in early fruit development in Cucumber. *Journal of Experimental Botany*, Vol. 59, No. 9, pp. 2299–2308.
- Ghorbani, P., Eshghi, S. and Haghi, H. (2017)** Effects of brassinosteroid (24- epibrassinolide) on yield and quality of grape (*Vitis vinifera* L.) 'Thompson Seedless'. *Vitis* **56**, 113–117 (2017).
- Ghosh, A., Panda, D., & Mondal, S. (2020).** Effect of micronutrients and growth regulators on pigment content, nitrate reductase activity and relative leaf water content of chilli (*Capsicum annuum* L). *Plant cell biotechnology and molecular biology*, 21(13-14), 37-51.
- Hipara AJ, Vaddoria MA, Jivanni LL, Patel L JB and Polara AM (2014).** Seed yield and quality as influence by plant growth regulators and stages of spray in bitter gourd (*Momordica charantia* l.) An international ejournal, 3(3): 282-287.
- Irfan, M., J. Alam, I. Ahmad, I. Ali and H. Gul. (2017)** Effects of exogenous and foliar applications of Brassinosteroid (BRs) and salt stress on the growth, yield and physiological parameters of *Lycopersicon esculentum* (Mill.). *Plant Science Today* **4**(3): 88-101.
- Khan, W., Prithiviraj, B., Smith, D.L.,(2003)** Photosynthetic responses of corn and soybean to foliar application of salicylates. *J. Plant Physiol.* 160: 485-492.
- Khan , A.S. and Chaudhry, N.Y. (2006).** GA3 improves flower yield in some cucurbits treated with lead and mercury.*African J. Biotech.*, 5 (2) :149-153.

Kumar, S. and Kaur, G. (2019) Effect of pre harvest application of salicylic acid on plant growth parameters and yield of strawberry cv. *Chandler*. *Gurpinder*.

Khatoon, F., Kundu, M., Mir, H., Nandita, K. and Kumar, D. (2020) Foliar Feeding of Brassinosteroid: A Potential Tool to Improve Growth, Yield and Fruit Quality of Strawberry (*Fragaria × ananassa Duch.*) under Non-Conventional Area. *International Journal of Current Microbiology and Applied Sciences*, ISSN: 2319-7706 Volume 9 Number 3 (2020).

Khatoon, F., Kundu, M., Mir, H., & Nahakpam, S. (2021) Efficacy of Foliar Feeding of Brassinosteroid to Improve Growth, Yield and Fruit Quality of Strawberry (*Fragaria × Ananassa Duch.*) Grown under Subtropical Plain. *Communications in Soil Science and Plant Analysis*. Volume 52.

Karuppaiah P. (2019) Effect of bioregulators and Feso4 on growth, yield and quality of crossandra (*Crossandra infundibuliformis L.*). *Horticulture International Journal*. 2019;3(1):1–5.

Latimer, J.G. (1991). Growth retardants affect landscape performance of Zinnia, Impatiens and Marigold. *Hort Sci* 26: 557–560.

Pawar, V., Meena, M.K., Jadia, M., Basediya, S. S., (2019) Study of Influence of Plant Growth Regulators (PGRs) on Yield Attributing Characters in Cucumber, *Int. J. Pure App. Biosci.* 7(3): 318-324 (2019).

Shakirova, M.F.; Sakhabutdinova, A.R.; Bezrukova, M.V.; Fathutdinova, R.A. and Fathutdinova, D.R., (2003). Change in the hormonal status of wheat seedling induced by salicylic acid and salinity. *Plant Sci.* **164** (3): 317 – 322.

Singh, Pramod Kumar A, Chaturvedi, Varun Kumar A, Bose, Bandana (2010) Effects Of Salicylic Acid On Seedling Growth And Nitrogen Metabolism In Cucumber (*Cucumis Sativus L.*) *Journal of Stress Physiology & Biochemistry*, Vol. 6 No. 3 2010, pp. 102-113.

Roy R., Nasiruddin K.M. (2011) Effect of different level of GA3 on growth and yield of cabbage. *Journal of Environmental Science and Natural Resources* 4: 79-82.

Ramani, M.M., Patel, C.R., Sharma, K.M., Amarcholi, J.J and Patel, A.H(2016) Effect of 28-homobrassinolode and Shoot Thinning on Vegetative Growth, Flowering and Yield of Mango cv. Kesar *Advances in Life Sciences* 5 (17) ISSN 2278-3849, 7097-7099.

Vishnu ritti, s. R. Doddagoudar, s.n.vasudevan, basavegowda and mallikarjun kenganal (2019) Influence of foliar spray of boron and plant growth regulators on hybrid seed yield and quality of bottle gourd. *J. Farm Sci.*, 32(3): (274-277).

Yuan L., Xu D.Q. (2001) Stimulation effect of gibberellic acid short-term treatment on the photosynthesis re-lated to the increase in Rubisco content in broad bean and soybean. *Photosynthesis Research* 68: 39-47.

UNDER PEER REVIEW