

Original Research Article

Response of different organic manures and inorganic fertilizers on vegetative growth of turnip (*Brassica rapa* L.) cv Purple Top White Globe

ABSTRACT

The experiment was laid out in a Randomized Block Design (RBD) with 17 treatment along with combinations of organic manure and inorganic fertilizer including control and each treatment replicated thrice. The turnip (*Brassica rapa* L.) crop in Brassicaceae family with Purple Top White Globe variety was grown with each comprising of treatments replicated thrice. results revealed that the application of T₁₂(50 % NPK + 50% N through Poultry manure) influenced most of the characteristics significantly and recorded the highest values of plant height (58.94cm, 59.85cm and 59.40cm), number of leaves per plant (19.24, 21.03 and 20.14), leaf length (52.80cm, 55.59cm and 54.20cm), leaf width (20.51cm, 22.16cm and 21.34cm), fresh weight of leaves (247.14, 250.84 and 248.99 g/plant) and dry weight of leaves (20.08, 22.01 and 21.05 g/plant) for the year 2018-19, 2019-20 and as pooled data, respectively.

Key words:- Organic manure, inorganic fertilizers, vegetative growth, and Turnip

INTRODUCTION

Brassica is a large genus including species used for oilseeds, leafy or root vegetables, and sauces. Turnip (*Brassica rapa* var. *rapifera* L.) (2n=2x=20) is a biennial root vegetable farmed globally as a vegetable and fodder (Rakow, 2004; Hammer et al., 2013). Turnip (*Brassica rapa*) is a vegetable of the cruciferous family. *Brassica rapa*, sometimes known as field mustard or turnip mustard, is a plant that is extensively grown as a leaf vegetable, root vegetable, and oilseed. Plants for the Future cannot accept responsibility for any negative consequences caused by the usage of plants. Before utilising a plant medicinally, always seek the counsel of an expert. Cancer is treated using a decoction of the leaves or stems. When cooked with fat, the root is used to treat breast cancers. Skin cancer is claimed to be helped by a salve made from the flowers. Furthermore, turnip extract can help decrease uric acid and remove kidney stones. It improves vision and is used to cure night blindness. Turnip syrup improves memory (Khashayar, 2007). Allardice (1993) is a natural pesticide found in turnip root peelings. Turnip agriculture in India is largely limited to the northwestern states of Punjab, Haryana, Rajasthan, and Western Uttar

Comment [E1]: I recommend: Influence of different organic manures and inorganic fertilizers on vegetative growth of turnip (*Brassica rapa* L.) cv Purple Top White Globe

Comment [E2]: ...with each comprising of treatments replicated thrice. It will be deleted. This has been mentioned before.

Comment [E3]: Results revealed that the application of T₁₂ (50 % Nitrogen:Phosphorus:Potassium - N:P:K + 50% N through poultry manure) significantly influenced most of the characteristics and recorded.....

Comment [E4]: This is from.....Lou et al. (2017) Turnip [*B. rapa* L. em. Metzg. subsp. *rapa* syn *B. campestris* L. spp. *rapifera* Sinsk, (AA, 2n = 20)] is an annual or biennial plant that is cultivated worldwide as vegetable and fodder (Rakow, 2004, Hammer et al., 2013).

So, I suggest to write...at the end [1, 2], cited by [3].

At the Reference section it will be:

1. Rakow, 2004...
2. Hammer et al., 2013....
3. Lou et al., 2017...

Please respect Instructions fro authors:

References must be listed at the end of the manuscript and numbered in the order that they appear in the text. Every reference referred in the text must also present in the reference list and vice versa. In the text, citations should be indicated by the reference number in brackets [3].

Comment [E5]: italic

Comment [E6]: italic

Comment [E7]: Please review and reformulate the paragraph. Some aspects are repeated.

Comment [E8]: Please be clear. Please use a phrase with a view to present the medicinal importance of the studied plant. For citation....[]

Comment [E9]: Please be clear. Allardice (1993) is a natural pesticide? Or Allardice si an authors? For citation....[]

Pradesh as an early winter season crop. Green Top, Purple Top, and 'Kenshin-Kaba' are all essential turnip cultivars for fodder production. **Yadav and colleagues (2021).**

Comment [E10]: ...[].

The thick base and young leaves are edible portions that are not as nutritionally deficient as is usually assumed. A kilogramme of edible roots contains 1.4 g protein, 6.2 g carbohydrate, minimal fat, 0.6 g mineral salt, 0.03 mg vitamin B-1, 0.02 mg vitamin B-2, 15 mg vitamin-C, 24 mg calcium, 0.4 mg iron, and 21 KJ. A hundred gramme of edible leaf contains: 4 g protein, 9.4 g carbohydrate, 1.5 g fat, 2.2 g minerals salt, 0.31 mg vitamin B-1, 0.57 mg vitamin B-2, 180 mg vitamin C, 710 mg calcium, 28.4 mg iron, 9396 µg calcium, 28.4 mg carotene, and 67 kJ energy (**Purseglove, 1988**). It improves visual acuity and is used to cure night blindness. Turnip syrup improves memory (**Khashayar, 2007**). Turnip root peelings include a natural pesticide (**Allardice, 1993**). The Mediterranean region is supposed to be the major centre of European kinds, whereas Eastern Afghanistan and the neighbouring areas of Pakistan are thought to represent secondary centres. It is a significant root vegetable planted as a summer crop in temperate climates and as a winter food in subtropical climates where winters are not harsh. It can be grown up to an elevation of 1500m above mean sea level or higher, however it is not appropriate for growing in damp equatorial lowlands (**Thamburaj and Singh 2018**). It is grown on 2500 acres in India, with an annual yield of 50,000 tonnes. Organic farming is a method that integrates links between soil, plant, water, soil microflora and fauna. Organic farming seeks to create a healthy soil, aids in correct energy flows in the soil, crop, water, and environment, while the plant systems maintain the biological life cycle alive and aids in the maintenance of significant yield levels (**Lampkin, 1990**). In recent years, the use of organic manures for increasing agricultural yield and sustaining soil fertility and productivity has gained popularity. Organic manure improves soil structure and water retention capacity (**Kale, 1991**).

Comment [E11]: Now, again details as regard as the plant medicinally importance. But, its composition is not the topic of this study. So, please do only a short information on this aspect. The documentation needs to focus on the fertilizers impact on plant growth (if there are previously studies).

MATERIALS AND METHODS

A investigation entitled "Response of different Organic Manures and Inorganic Fertilizers on Vegetative Growth, Root Yield and Quality of Turnip (*Brassica Rapa* L.) cv. Purple Top White Globe", was carried out during (2018-19 & 2019-20) at Horticulture Research Farm, Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (Allahabad), U.P., India-211007. The experiment was laid out in a Randomized Block Design (RBD) with 17 treatment along with combinations of organic manure

Comment [E12]: The INTRODUCTION section needs to be careful review and information clearly organised. Moreover, at the end you must to write the objectives of your work.

Comment [E13]: MATERIALS AND METHODS

Please, do not write the article title here.

For example:

It is necessary to present the experiment site and duration of the study, the biological material used, the experimental variants and design, the parameters analysed (each with the specific expression unit), the statistical analyses.

and inorganic fertilizer including control and each treatment replicated thrice. The turnip (*Brassica rapa* L.) crop in Brassicaceae family with Purple Top White Globe variety was grown with each comprising of treatments replicated thrice. Source of variables were NPK, FYM, Vermicompost, Poultry Manure and Goat manure (at 25%, 50%, 75% and 100%) and combination of them with control treatment taken as seventeen treatments. To observe the effect of organic manure and inorganic fertilizer, vegetative growth parameters viz. plant height (cm), number of leaves plant⁻¹, leaf length (cm), leaf width (cm), fresh weight (g plant⁻¹) of leaves and dry weight (g plant⁻¹) of leaves, were studied in the investigation.

RESULTS AND DISCUSSION

The results regarding vegetative growth of turnip were statistically analyzed and have been presented in table: 1. A study of the table shows that the application of organic manures and inorganic fertilizers had significant effect on the plant height (cm), number of leaves plant⁻¹, leaf length (cm) and leaf width (cm) of turnip in 1st year, 2nd year and when pooled. Revealed that the significant maximum plant height 58.94cm, 59.85cm and 59.40cm for the year 2018-19, 2019-20 and when pooled, respectively, was recorded with treatment T₁₂ (50 % NPK + 50% N through Poultry manure) and followed by treatments T₁₃ (50 % NPK + 50% N through Goat manure) 57.58cm, 58.20cm and 57.89cm at 55 DAS for the year 2018-19, 2019-20 and when pooled respectively, which were significantly at par with each other and were significantly superior over T₁ (control) 35.86cm, 37.14cm and 36.50cm at 55 DAS for the year 2018-19, 2019-20 and pooled data, respectively. The results of the present investigation agreed with the finding of Kumar *et al.* (2014), Mbatha *et al.* (2014), Verma *et al.* (2016). Okokoh and Bisong (2011) reported similarly that application of 10 to 15 t/ha of poultry manure resulted in increased height of amaranthus plants. The increased growth parameters attributed to beneficial effect of PM has been reported by Uddain *et al.* (2010) and Subedi *et al.* (2018) in radish, Sylvester *et al.* (2015) in carrot and Jagadeesh *et al.* (2018) in beetroot. Revealed that the significant maximum number of leaves per plant 19.24, 21.03 and 20.14 for the year 2018-19, 2019-20 and when pooled respectively, was recorded with treatment T₁₂ (50 % NPK + 50% N through Poultry manure) and followed by treatments T₁₃ (50 % NPK + 50% N through Goat manure) 18.68, 20.25 and 19.47 at 55 DAS for the year 2018-19, 2019-20 and when pooled respectively, which were significantly at par with each other and were significantly superior over T₁ (control) 10.83,

Comment [E14]: ...with each comprising of treatments replicated thrice. It has been mentioned before.

Comment [E15]: I appreciate that each variant should be clear described and all presented in a Table. We need to know what does it mean....T2, or T3...an so on. Also, below the results Tables, the Legend for these is necessary.

Comment [E16]: Source of variables were NPK, FYM..... At the first apparition in text, the abbreviations must be explained

Comment [E17]: Please specify also how many treatments have been done and when. When the growth parameters have been analysed. Athow many days after sowing, for instance.... These data are necessary in Materials and Methods section.

Comment [E18]: At the end, please specify the statistical analyses....

Comment [E19]: RESULTS AND DISCUSSION

Comment [E20]:statistically analyzed and are presented in Table 1.

In fact, as we can see, the vegetative growth characteristics are shown....in Table 1 and Table 2, too.

Or, as we can see later, you can use Table 1a and Table 1b, taking into consideration that there are presented vegetative growth parameters.

Then, when these parameters are presented, I recommend to have their better organisation.

For example:

Plant height (cm)

.....

Number of leaves plant⁻¹

.....

And so on...

Comment [E21]: The subject is missing...

Comment [E22]: DAS.....? ...

Comment [E23]:which were statistically at par with each other and were significantly superior over T₁

Comment [E24]: ... kumar *et al.* (2014), Mbatha *et al.* (2014), Verma *et al.* (2016). Okokoh and Bisong (2011) ...

Comment [E25]: PM
The abbreviation PM has not used before. Now, at its first apparition please explained it.

Comment [E26]: ...[]....

Comment [E27]: The subject is missing...

Comment [E28]: poultry manure)
Now, you can use only...PM)

Comment [E29]: ..were statistically at par with each other

11.13 and 10.98 at 55 DAS for the year 2018-19, 2019-20 and pooled data, respectively. These results agree with the findings. who was found significant increase in number of leaves of radish with the sole application of NPK. The results of the present investigation agreed with the finding of [Soheir et al. \(2012\)](#), [Vijaya kumara et al. \(2012\)](#), [Aisha et al. \(2014\)](#), [Kumar et al. \(2014\)](#). The increased growth parameters attributed to beneficial effect of PM has been reported by [Uddain et al. \(2010\)](#) and [Subedi et al. \(2018\)](#) in radish, [Sylvester et al. \(2015\)](#) in carrot and [Jagadeesh et al. \(2018\)](#) in beetroot. Since, poultry manure contains high amount of major nutrients comparative to the other organic nutrients used as treatments probably which helps for the proper growth and development of vegetative structures (i.e., leaves). The results were in conformity with findings of [Tiamiyu et al., 2012](#) in okra. Among the treatments applied, treatment T₁₂(50 % NPK + 50% N through Poultry manure) recorded significantly maximum leaf length (cm) 52.80cm, 55.59cm and 54.20cm for the year 2018-19, 2019-20 and as pooled data respectively, and closely followed by treatments T₁₃ (50 % NPK + 50% N through Goat manure) 51.15cm, 53.91cm and 52.53cm for the year 2018-19, 2019-20 and when pooled, respectively, which were significantly at par with each other and were significantly superior over T₁ (control) 28.43cm, 31.15cm and 29.79cm for the year 2018-19, 2019-20 and pooled data respectively. These results coincide with the previous findings of [Islam et al. \(2011\)](#) who found maximum leaf length in radish when NPK was used, as the plants received more readily available applied nutrients, which might had increased the vegetative growth and leaf length in radish. Similar results were also reported by stating an increase in cabbage leaf length, when inorganic fertilizers were used. Similar finding for the leaf length per plant were reported by [Mbatha et al. \(2014\)](#), [Zeid et al. \(2015\)](#), [Kiran et al. \(2016\)](#), [Wahocho et al. \(2016\)](#) and [Zhou-Dongmei., \(2005\)](#) reported that the effect of application of poultry manures on growth of radish (*Raphanus sativus* L.) and pakchoi (*Brassica chinensis* L.). The increased growth parameters attributed to beneficial effect of PM has been reported by [Uddain et al. \(2010\)](#) and [Subedi et al. \(2018\)](#) in radish, [Sylvester et al. \(2015\)](#) in carrot and [Jagadeesh et al. \(2018\)](#) in beetroot.

Among the treatments applied, treatment T₁₂(50 % NPK + 50% N through Poultry manure) exhibited significantly maximum leaf width (cm) (20.51cm, 22.16cm and 21.34cm) for the year 2018-19, 2019-20 and as pooled data, respectively, and closely followed by treatments T₁₃ (50 % NPK + 50% N through Goat manure) 19.87cm, 21.54cm and 20.71cm for the year 2018-19, 2019-20 and when pooled respectively, which were statistically significantly at par with

Comment [E30]: []..

Comment [E31]: []

Comment [E32]: Please attention to quote, when necessary.
Since, poultry manure contains high amount of major nutrients comparative to the other organic nutrients used as treatments probably which helps for the proper growth and development of vegetative structures (i.e., leaves).The results were in conformity with findings of [Tiamiyu et al., 2012](#) in okra.

Since, poultry manure (100%) contains high amount of major nutrients comparative to the other organic nutrients used as treatments probably which helps for the proper growth and development of vegetative structures (i.e., leaves).The results were in conformity with findings of [Tiamiyu et al., 2012](#) in okra.....These are from.....[Jagadeesh et al. 2018](#)

Comment [E33]:were significantly at par... Again, please...were statistically at par....

Comment [E34]: [Mbatha et al. \(2014\)](#), ... Again, attention...[]

each other and were significantly superior over T₁ (control) 8.81cm, 9.25cm and 9.03cm for the year 2018-19, 2019-20 and pooled data, respectively. Similar finding for the leaf width were reported by **Chitti et al. (2018)**, **Dhital et al. (2018)**, **Ingole et al. (2018)** and **Zhou-Dongmei., (2005)** reported that the effect of application of poultry manures on growth of radish (*Raphanus sativus* L.) and pakchoi (*Brassica chinensis* L.). The increased growth parameters attributed to beneficial effect of PM has been reported by **Uddain et al. (2010)** and **Subedi et al. (2018)** in radish, **Sylvester et al. (2015)** in carrot and **Jagadeesh et al. (2018)** in beetroot.

Comment [E35]: Chitti et al. (2018)...
| |...

Comment [E36]: Uddain et al. (2010) ...
| |...

The results regarding vegetative growth of turnip were statistically analyzed and have been presented in table: 2. A study of the table shows that the application of organic manures and inorganic fertilizers had significant effect on the fresh weight of leaves (g/plant) and dry weight of leaves (g/plant) of turnip in 1st year, 2nd year and when pooled. Among the treatments applied, treatment T₁₂(50 % NPK + 50% N through Poultry manure)exhibited significantly maximum fresh weight of leaves (g/plant)247.14 g/plant, 250.84 g/plant and 248.99 g/plant for the year 2018-19, 2019-20 and as pooled data respectively, and closely followed by treatments T₁₃ (50 % NPK + 50% N through Goat manure)241.97 g/plant, 243.62 g/plant and 242.80 g/plantfor the year 2018-19, 2019-20 and when pooled respectively, which were significantly at par with each other and were significantly superior over T₁ (control) 146.61 g/plant, 149.54 g/plant and 148.08 g/plant for the year 2018-19, 2019-20 and pooled data respectively. The presence of adequate amount of NPK might be the major cause of enhancing the soil fertility level which promoted plant growth thus causing an increased leaf weight plant. The enhancements in growth parameters of radish were also reported by the application of NPK. similar finding for the fresh weight of leaves per plant were reported by **Kumar et al. (2018)**, **Subedi et al. (2018)**, **Messele (2016)** and **Zhou-Dongmei., (2005)** reported that the effect of application of poultry manures on growth of radish (*Raphanus sativus* L.) and pakchoi (*Brassica chinensis* L.). The increased growth parameters attributed to beneficial effect of PM has been reported by **Uddain et al. (2010)** and **Subedi et al. (2018)** in radish, **Sylvester et al. (2015)** in carrot and **Jagadeesh et al. (2018)** in beetroot. Among the treatments applied, treatment T₁₂(50 % NPK + 50% N through Poultry manure)exhibited significantly maximum dry weight of leaves (g/plant)20.08 g/plant, 22.01 g/plant and 21.05 g/plant for the year 2018-19, 2019-20 and as pooled data respectively, and closely followed by treatments T₁₃ (50 % NPK + 50% N through Goat manure)19.67 g/plant, 21.52 g/plant and 20.60g/plantfor the year 2018-19, 2019-20 and when pooled respectively,

Comment [E37]: The results regarding vegetative growth of turnip were statistically analyzed and have been presented in table: 2. As I said at the Results section beginning, in both Tables (1 and 2) there are shown vegetative growth parameters!
So, pat attention, please.

Please consider my suggestions.
More clearly presentation is needed for the following data, too.

which were significantly at par with each other and were significantly superior over T₁ (control) 8.87 g/plant, 9.55 g/plant and 9.21 g/plant for the year 2018-19, 2019-20 and pooled data respectively. The effectiveness of NPK was obvious due to greater nutrients content and their readily availability. Our results agree with the findings a significant increase in the weight of leaves plant with the application of manures and chemical fertilizers in radish. Similar results were observed by **Vijayakumari et al. (2012)**, **Zakir et al. (2012)**, **Aisha et al. (2014)** and **Zhou-Dongmei., (2005)** reported that the effect of application of poultry manures on growth of radish (*Raphanus sativus* L.) and pakchoi (*Brassica chinensis* L.). The increased growth parameters attributed to beneficial effect of PM has been reported by **Uddain et al. (2010)** and **Subedi et al. (2018)** in radish, **Sylvester et al. (2015)** in carrot and **Jagadeesh et al. (2018)** in beetroot.

CONCLUSION

Based on the current inquiry in both the 2018-19 and 2019-20 fiscal years. The treatment T12 (50% NPK + 50% N through Poultry manure) was determined to be the most effective in terms of turnip vegetative development. Following a few more conjunctive experiments, producers can be advised to apply this organic manure and inorganic fertilizer mix.

Comment [E38]: CONCLUSION

Based on the current study, in both years (2018-19 and 2019-20), the treatment T12 (50% NPK + 50% N applied as poultry manure) was determined to be the most effective in terms of turnip vegetative development. Following a few more added experiments, producers can be advised to apply this organic manure and inorganic fertilizer mix.

REFERENCES

- Allardice P. (1993)**. A - Z of Companion Planting. Cassell Publishers Ltd. 1993, ISBN 0-304-34324-2 A well produced and very readable book 1993.
- Aisha H., Ali M.R., Mahmoud S., Asmaa R. and El- Desuki M. (2014)**. Effect of Various Levels of Organic Fertilizer and Humic Acid on the Growth and Roots Quality of Turnip Plants (*Brassica rapa*). *Current Science International*.3(1): 7-14.
- Chitti J., Madhavi M., Siva Prasad M. and Padmaja, V.V. (2018)**. Effect of Organic Manures on Growth and Yield attributes of Beet Root Cv. Crimson Globe. *Int.J.Curr.Microbiol.App.Sci*. 7(11): 3538-3553.
- Dhital B., Khanal A., Sharma G. and Sharma A. (2018)**. Effect of organic and chemical sources of nitrogen and seedling age at transplant on growth and yield of Kohlrabi. *Advances in Plants & Agriculture Research*.8(5): 357-361.

Comment [E39]: REFERENCES Please respect Instructions for authors.

References must be listed at the end of the manuscript and numbered in the order that they appear in the text. Every reference referred in the text must also present in the reference list and vice versa. In the text, citations should be indicated by the reference number in brackets [3].

Example:

Reference to a journal:

For Published paper:

1. Hilly M, Adams ML, Nelson SC. A study of digit fusion in the mouse embryo. *Clin Exp Allergy*. 2002;32(4):489-98.

- Hammer, K., Gladis, T. H., Laghetti, G. and Pignone, D. (2013).** The wild and the grown-remarks on *Brassica*. *International Journal of Agricultural Science* 3: 453-480.
- Ingole, VS, Wagh, AP, Nagre, PK and Bharad, SG (2018).**Effect of combination of organic manure and biofertilizer for better growth and yield of beetroot (*Beta vulgaris* L.). *International Journal of Chemical Studies*; 6(5): 1222-1225
- Jagadeesh, C., Madhavi, M., Prasad, M.S. and Padmaja, V.V. (2018).** effect of organic manures on growth and yield attributes of beet root cv. crimson globe. *International Journal of Current Microbiology and Applied Sciences*, 7(11):3538-3553.
- Kale, R. N., Bano, K. and Satyavati, G. P., 1991,** Influence of vermicompost application on growth and yield of cereals, vegetables and ornamental plants. *Final Rep. of KSCST Project N 67-04/vermin/34B*, p. 3478.
- Khashayar P.(2007)** Alternative Medicine, Turnip. Press TV, Tehran 2007.
- Kumar Sandeep, Sutanu Maji, Sanjay Kumar and Harsh Deep Singh(2014)** Efficacy of organic manures on growth and yield of radish (*Raphanus sativus* L.) cv. JAPANESE WHITE. *International Journal of Plant Sciences*. 1(9) 57-60.
- Kumar P., Meghwal P. R. and Painuli D. K. (2014).**Effect of organic and inorganic nutrient sources on soil health and quality of turnip.*Indian J. Hort.* 71(2): 222-226.
- Kiran M., Jilani M. S., Waseem K. and Sohail M. (2016).**effect of organic manures and inorganic fertilizers on growth and yield of radish (*raphanussativus*l). *Pakistan J.Agric. Res.* Vol. 29 No.4.
- Kumar S., Dev P., Kumar J. and Kumar H. (2018).**Impact of organic manures and biofertilizers on the performance of radish (*Raphanus sativus* L.) *HortFlora Res. Spectrum*, 7(2): 137-140.
- Lampkin, N.(1990).** In: Organic farming, Ipswich, W.K., Farming Press Book, pp. 801-910.
- Mbatha, A.N., Ceronio, G.M &Coetzer, G.M. (2014).**Response of carrot (*Daucuscarrota* L.) yield and quality to organic fertilizer. *South African Journal of Plant and Soil*, 31(1): 1-6.

- Messele B. (2016).** Effects of Combined Application of Organic –P and Inorganic N Fertilizers on Yield of Carrot (*DaucusCarrota L.*). *Agri Res & Tech: Open Access Journal*.2(2): 35-39.
- Okokoh, S. J. and Bisong, B. W. (2011)** Effect of Poultry Manure and Urea- N on Flowering Occurrence and Leaf Productivity of *Amranthus cruentus* in Calabar. *Journ.of Apl. Sic. Environmental Management*. 15. (1) 13- 15.
- Purseglove JW, Brown EG, Green CL, Robbins SRJ. (1988).** Spices; 7(8):447-462.
- Rakow. G (2004).** Species origin and economic importance of *Brassica*. Biotechnology in Agriculture and Forestry, Vol. 54. New York Springer-Verlag Berlin Heidelberg pp. 3-11.
- Soheir E. E., Hendawy S.F., Youssef A.A., Naguib N.Y. and Hussein M.S. (2012).**Response of Turnip (*Brassica rapa*) Plants to Minerals or Organic Fertilizers Treatments. *Journal of Applied Sciences Research*.8(2): pp-628-634.
- Sylvestre, H., Constance, M., Esdras, N. and Athanase, N. (2015).** Effect of poultry manure and NPK on growth and yield of carrot in Rulindo district, Rwanda. *International Journal of Novel Research in Life Sciences*, 2(1): 42-48.
- Subedi S., Srivastava A., Sharma M.D.& ShahS.C. (2018).** Effect of organic and inorganic nutrient sources on growth, yield and quality of radish (*raphanussativusL.*) varieties in chitwan, nepal. *SAARC J. Agri.*, 16(1): 61-69.
- Tiamiyu (2012).** Effect of Sources of Organic Manure on Growth and Yields of Okra (*Abelmoschus esculentus L.*) Agriculture Usmanu Danfodiyo University, Sokoto, Nigeria.
- Thamburaj, S and Singh, N. (2018).** Textbook of Vegetable, Tuber crops and Spices. ICAR, New Delhi, pp 161
- Uddain, J., Chowdhury, S. and Rahman, M.J. (2010).** Efficacy of different organic manures on growth and productivity of radish (*Raphanus sativus L.*). *International Journal of Agriculture, Environment and Biotechnology*, 3(2):189-193.
- Vijayakumari B., Sasikala V. and Poornima C. P. (2012).**Effect of organic and inorganic manures on biometric and yield parameters of radish (*Raphanussavitus L.*) cv. Pusaphepki. *International journal of plant sciences*.7(1): pp-130-134.

- Verma U.K., Kumar R., Kumar S. and Kumar A. (2016)** Integrated effect of organic manures and inorganic fertilizers on growth, yield and yield attributes of Radish cv. KalyanpurSafed. *International Journal of Agricultural Invention*.1(2): pp-158 – 161.
- Wahocho N. A., Wahocho S. A., Memon N., Leghari M. H. and Baloch Q. B. (2016).** Growth and yield response of turnip to various nitrogen application rates. *Pak. J. Agri., Agril. ngg., Vet. Sci.*32 (2): pp-143-149.
- Yadav Chandraprakash, Kumar Sudhir Mishra, Kumar Manish Singh, Roy Sapna and Tiwari Prashant (2021).** Effect of integrated nutrient management on growth, yield and shelf life of turnip (*Brassica rapa* L.) cv. purple top white. *The Pharma Innovation Journal* 2021; 10(2): 100-103
- Zhou Dong-Mei., (2005)** “Copper and Zn uptake by radish and pakchoi as affected by application of livestock and poultry manures”. *Chemosphere* 59.2 (2005): 167-175.
- Zakir H.M., Sultana M.N. and Saha K.C. (2012).** Influence of commercially available organic vs inorganic fertilizers on growth yield and quality of turnip. *J. Environ. Sci. Nat. Res.* 5 (1): 39 – 45.
- Zeid H.A., Wafaa H. M., Abou EI Seoud I. I. and Alhadad W. A. A. (2015).** Effect of organic materials and inorganic fertilizers on the growth, mineral composition and soil fertility of turnip plants (*Raphanussativus*L.) grown in sandy soil. *Middle East J. Agric. Res.* 4(1): 77-87.

Table 1 Response of different organic manures and inorganic fertilizers on vegetative growth of turnip (*Brassica rapa* L.) cv Purple Top White Globe

Treatment	Vegetative growth											
	Plant height (cm)			Number of leaves plant ⁻¹			Leaf length (cm)			Leaf width (cm)		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled
T ₁	35.86	37.14	36.50	10.83	11.13	10.98	28.43	31.15	29.79	8.81	9.25	9.03
T ₂	41.99	42.28	42.14	12.94	13.26	13.10	34.21	38.85	36.53	11.47	13.00	12.24
T ₃	43.05	44.34	43.70	13.57	14.14	13.86	36.26	40.47	38.37	12.64	14.23	13.44
T ₄	46.56	47.90	47.23	14.19	15.21	14.70	39.11	42.84	40.98	14.32	15.20	14.76
T ₅	44.92	45.22	45.07	13.96	14.67	14.32	38.79	41.28	40.04	13.74	15.10	14.42
T ₆	51.80	53.08	52.44	15.86	17.44	16.65	44.32	47.56	45.94	16.69	17.84	17.27
T ₇	52.72	54.12	53.42	16.20	18.15	17.18	45.68	49.90	47.79	17.26	18.81	18.04
T ₈	53.96	55.04	54.50	17.16	18.89	18.03	47.90	51.5	49.70	18.34	19.50	18.92
T ₉	53.21	54.69	53.95	16.51	18.37	17.44	46.81	50.67	48.74	17.59	19.22	18.41
T ₁₀	55.43	56.02	55.73	17.71	19.33	18.52	48.57	52.72	50.65	19.17	19.94	19.56
T ₁₁	56.81	57.39	57.10	18.10	19.69	18.90	49.95	53.13	51.54	19.48	21.12	20.30
T ₁₂	58.94	59.85	59.40	19.24	21.03	20.14	52.80	55.59	54.20	20.51	22.16	21.34
T ₁₃	57.58	58.20	57.89	18.68	20.25	19.47	51.15	53.91	52.53	19.87	21.54	20.71
T ₁₄	47.16	48.03	47.60	14.42	15.58	15.00	39.67	43.17	41.42	14.78	15.87	15.33
T ₁₅	48.00	50.29	49.15	14.68	16.05	15.37	40.84	44.63	42.74	15.29	16.75	16.02
T ₁₆	50.62	52.79	51.71	15.36	17.03	16.20	43.41	46.96	45.19	16.27	17.64	16.96
T ₁₇	49.76	51.64	50.70	14.96	16.47	15.72	41.64	45.57	43.61	15.70	17.13	16.42
C.D. at 5%	2.25	2.30	2.27	0.70	0.76	0.73	1.94	2.10	2.02	0.73	0.79	0.76
SEd (±)	1.10	1.13	1.12	0.34	0.37	0.36	0.95	1.03	0.99	0.36	0.39	0.37
F-Test	S	S	S	S	S	S	S	S	S	S	S	S

Comment [E40]: I suggest: Table 1a. Influence of different organic manures and inorganic fertilizers on vegetative growth of turnip (*Brassica rapa* L.) cv Purple Top White Globe

Comment [E41]: Legend for the applied treatments is necessary.

Table 2 Response of different organic manures and inorganic fertilizers on vegetative growth of turnip (*Brassica rapa* L.) cv Purple Top White Globe

Treatment	Vegetative growth					
	Fresh weight of leaves (g/plant)			Dry weight of leaves (g/plant)		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled
T ₁	146.61	149.54	148.08	8.87	9.55	9.21
T ₂	152.74	156.25	154.50	11.56	12.33	11.95
T ₃	163.70	166.97	165.34	12.24	13.30	12.77
T ₄	171.01	174.94	172.98	13.70	15.08	14.39
T ₅	168.98	170.84	169.91	12.92	14.17	13.55
T ₆	191.87	197.81	194.84	16.68	17.81	17.25
T ₇	201.86	205.89	203.88	17.12	18.04	17.58
T ₈	211.97	210.29	211.13	18.51	19.62	19.07
T ₉	205.97	207.73	206.85	17.84	18.59	18.22
T ₁₀	216.14	222.62	219.38	18.94	19.96	19.45
T ₁₁	238.01	239.55	238.78	19.26	20.73	20.00
T ₁₂	247.14	250.84	248.99	20.08	22.01	21.05
T ₁₃	241.97	243.62	242.80	19.67	21.52	20.60
T ₁₄	172.10	178.98	175.54	13.86	15.41	14.64
T ₁₅	176.86	180.39	178.63	14.67	16.15	15.41
T ₁₆	186.93	192.35	189.64	16.34	17.74	17.04
T ₁₇	181.08	183.79	182.44	15.38	16.93	16.16
C.D. at 5%	3.52	2.41	2.56	0.72	0.77	0.75
SEd (\pm)	1.73	1.19	1.26	0.35	0.38	0.37
F-Test	S	S	S	S	S	S

Comment [E42]: I suggest: Table 1b. Influence of different organic manures and inorganic fertilizers on vegetative growth of turnip (*Brassica rapa* L.) cv Purple Top White Globe

Comment [E43]: Legend for the applied treatments is necessary.