

## Original Research Article

### **Physiological traits associated with seed yield in chickpea (*Cicer arietinum* L.) genotypes under different environmental conditions**

#### **Abstract**

A field experiment was conducted during Rabi, 2019-20 at Sehore, Madhya Pradesh to investigate the association of yield and its related components of physiological traits in Chickpea under three sown environments. Seed yield per plant was showed significant positive association with LA & LAI at 60 days after sowing in E-I, CGR in E-III. This suggested that attributes might be utilised primarily for selection of high yielding genotypes, with yield contributing characters potentially playing a key role as selection parameters for isolating a high yield in eligible genotypes.

**Keywords:** Chickpea, Correlation, Seed yield

#### **INTRODUCTION**

The chickpea, (*Cicer arietinum* L.) is a significant pulse legume that is grown and eaten all over the world. The world's biggest producer and consumer of chickpeas is India. It is abundant in linoleic and oleic acid, two essential unsaturated fatty acids for nutrition. Important sterols found in chickpea oil include stigma, campesterol, and  $\beta$ -sitosterol. Chickpea seeds also include calcium, magnesium, phosphorus, and especially potassium. Important vitamins like riboflavin, niacin, thiamin, folate, and the precursor of vitamin A,  $\beta$ -carotene, can all be found in chickpeas. High temperatures have an impact on the physiological processes, growth, and development of the crop. Heat stress shortens the duration of plant developmental stages and the carbon assimilation process within the plant, resulting in low biomass production and reduced source-sink activity in most crops (Chen et al. 1982). Heat stress also affects crop growth and development by altering the morpho-anatomical, physiological, and biochemical makeup of most crops (Harding et

al., 1990). The most sensitive genotype of chickpeas had reduced chlorophyll content and symptoms of chlorosis leaves at 40/30°C. Heat waves are frequently encountered when growing chickpea, which has an impact on crop growth and productivity (Jumrani and Bhatia, 2014). After harvesting rice, early potatoes, or cotton, there is considerable potential for late planting of chickpea in North India. In such cases, the crop must be planted up until the end of December or the beginning of January. Low temperatures are seen during sowing time for this late-planted chickpea crop, and high temperatures are experienced at the conclusion of the growing season. Poor and slow vegetative development is caused by low temperatures at the beginning of crop growth, whereas forced maturation and problems are caused by high temperatures towards the end of cropping season. For these reasons, present study was undertaken to assess the associations for seed yield and related physiological traits in chickpea in three sown conditions.

#### **Materials and Methods**

The research was conducted at the Rafi Ahmed Kidwai College of Agriculture, Sehore, in three environments (normal November 28, 2019, mid-late December 28, 2019, and very late January 28, 2020) to correlate excessive heat stress with pollination time, reproductive time, and grain filling period (Madhya Pradesh). The minimum and maximum temperatures of the atmosphere ranged from 7.4°C in November

and 45.0°C in April. Under the All India Coordinated Research Project on Chickpea, the experiment was set up in Rabi 2019–20 with two replications and a randomized complete block design. Two rows of each genotype, measuring 3 m each, were sowed. The row to row spacing distance was kept 30 cm and plant to plant spacing was 30 x 10 cm. The recommended packages of practices were

followed to raise a healthy crop. Data were accumulated on Leaf Area (LA) and Leaf Area Index (LAI) at 45 days, LA and LAI at 60 days, Total Dry Matter (TDM) at 45 days, TDM at 60 days, Crop growth rate (CGR), Net assimilation rate (NAR), Chlorophyll index (SPAD) were recorded each of three

## Result and Discussion

The values of correlation coefficient for seed yield per plant with physiological traits were given in Table-I, II, III for each environment, separately. In the current investigation, a correlation between physiological traits and seed yield was recorded separately in each of the three environmental conditions. The seed yield per plant was highly influenced by physiological traits of the plants, but different environmental conditions plant was not similar interrelationship due to environment fluctuations. Seed yield per plant was showed significant positive association with LA & LAI

environments. These were estimated from five randomly selected plants. The genotypes were organized in randomized complete block design with two replications. According to, Miller *et al.* 1958 were computed utilizing variance and co-variance of phenotypic correlation coefficients among characters.

at 60 days after sowing in E-I, CGR except in E-I&E-II. This suggested that attributes might be utilized primarily for selection of high yielding genotypes, with yield contributing characters potentially playing a key role as selection parameters for isolating a high yield in eligible genotypes. In E-I, E-II, and E-III settings, intercorrelations between physiological traits showed a similar statistically significant trend because environmental factors including temperature, humidity, and daylight hours had no influence on the transmission of these traits.

**Table I. Correlation coefficient analysis among physiological traits and seed yield in Environmental-1**

Traits	LA at 45 DAS	LA at 60 DAS	LAI at 45 DAS	LAI at 60 DAS	TDM at 45 DAS	TDM at 60 DAS	NAR	CGR	Chlorophyll Index	Seed yield per plant
LA at 45 DAS	1									
LA at 60 DAS	0.476*	1								
LAI at 45 DAS	0.999**	0.475*	1							
LAI at 60 DAS	0.476*	0.999**	0.475*	1						
TDM at 45 DAS	0.746**	0.648**	0.746**	0.647**	1					
TDM at 60 DAS	0.432*	0.660**	0.432*	0.660**	0.632**	1				
NAR	0.468*	0.073	0.467*	0.072	0.534**	-0.293	1			
CGR	0.253	0.556**	0.253	0.556**	0.400*	0.963**	-0.532**	1		
Chlorophyll index	0.272	0.347	0.274	0.346	0.249	0.389	-0.026	0.374	1	
Seed Yield per Plant (g)	0.172	-0.05	0.173	-0.049	0.252	0.001	0.286	-0.087	0.079	1

Note \* and\*\* significant at 5% and 1% level of probability, respectively (DAS = Days or not ????????)  
**What is DAS?????**

The positive and significant association was noticed between LA and LAI at 45 DAS with LA & LAI at 60 DAS, total dry matter at 45 days after sowing & 60 DAS. LA and LAI at 60 DAS with LA & LAI at 45 DAS, total dry matter at 45 days after sowing & 60 DAS. TDM at 45 days after sowing with LA & LAI at 45 days after sowing & 60 DAS. TDM at 60 days after sowing with LA & LAI at 45 days after sowing & 60 DAS, crop growth rate. Intercorrelations among the traits and with seed yield did not follow similar significant trend of most of the traits in all the environmental conditions because the environmental factors viz., temperature, humidity, sunshine hours,

had much influenced in transmission of these traits. The dissimilar positive significant trend was reported between LA and LAI at 45 DAS with CGR in E-I, NAR in E-II, chlorophyll index in E-III. LA and LAI at 60 DAS with Seed yield per plant except in E-II&III, CGR except in E-III. TDM at 45 days after sowing with total dry matter at 60 DAS in E-I&E-II, CGR in E-I&E-II, NAR in E-II, Chlorophyll index in E-I, seed yield in E-I. TDM at 60 days after sowing with Crop growth rate except in E-II & E-III, total dry matter at 45 DAS in E-I&E-II, chlorophyll index in E-II. NAR with LA and LAI at 45 DAS in E-II, total dry matter at 45 DAS in E-II, CGR in E-

III. CGR with total dry matter at 60 DAS except in E-II, LA&LAI at 45 days after sowing & 60 DAS in E-I, total dry matter at 45 days after sowing in E-I&E-II, LA&LAI at 60DAS in E-II, seed yield in E-III. Chlorophyll index with total dry matter at 45 DAS except in E-II & E-III, total dry matter at 60DAS in E-II, CGR in E-II. The above results were in agreement with findings of

Tejpalet *et al.* (2004) for chlorophyll content was highest in early sowing in temperature tolerance in chickpea, Singh *et al.* (2005) for chlorophyll content was closely associated with NAR and LAI under different planting dates in chickpea, Singh *et al.* (2011) for seed yield was positive correlated with LAI, total chlorophyll content, CGR for heat tolerance in chickpea.

**Table II. Correlation coefficient analysis among physiological traits and seed yield in Environmental-II**

Traits	LA at 45 DAS	LA at 60 DAS	LAI at 45 DAS	LAI at 60 DAS	TDM at 45 DAS	TDM at 60 DAS	NAR	CGR	Chlorophyll index	Seed Yield per Plant (g)
LA at 45 DAS	1									
LA at 60 DAS	0.493*	1								
LAI at 45 DAS	0.999**	0.491*	1							
LAI at 60 DAS	0.493*	0.999**	0.491*	1						
TDM at 45 DAS	0.489*	0.595**	0.486*	0.595**	1					
TDM at 60 DAS	0.549**	0.483*	0.548**	0.482*	0.755**	1				
NAR	0.176	-0.558**	0.177	-0.558**	-0.246	0.091	1			
CGR	0.515**	0.402*	0.514**	0.402*	0.607**	0.979**	0.185	1		
Chlorophyll index	-0.066	0.069	-0.064	0.069	0.375	0.191	0.062	0.117	1	
Seed Yield per Plant (g)	0.295	0.475**	0.289	0.477*	0.384	0.213	-0.181	0.14	-0.038	1

Note \* and\*\* significant at 5% and 1% level of probability, respectively

**Table III. Correlation coefficient analysis among physiological traits and seed yield in Environmental-III**

Traits	LA at 45 DAS	LA at 60 DAS	LAI at 45 DAS	LAI at 60 DAS	TDM at 45 DAS	TDM at 60 DAS	NAR	CGR	Chlorophyll index	Seed Yield per plant (g)
LA at 45 DAS	1									
LA at 60 DAS	0.810**	1								
LAI at 45 DAS	0.999**	0.811**	1							
LAI at 60 DAS	0.810**	0.999**	0.810**	1						
TDM at 45 DAS	0.415*	0.517**	0.415*	0.517**	1					
TDM at 60 DAS	0.321	0.513**	0.321	0.513**	0.239	1				
NAR	-0.296	-0.598**	-0.297	-0.598**	-0.481*	0.148	1			
CGR	0.126	0.27	0.125	0.271	-0.232	0.888**	0.375	1		
Chlorophyll index	0.364	0.012	0.363	0.011	-0.151	-0.29	-0.017	-0.22	1	
Seed Yield per Plant (g)	-0.054	0.023	-0.054	0.024	-0.34	0.277	0.063	0.437*	-0.067	1

Note \* and\*\* significant at 5% and 1% level of probability, respectively

### What's the conclusion?

### References

- Chen YM, Backman K, Magasanik B. Characterization of a gene, *glnL*, the product of which is involved in the regulation of nitrogen utilization in *Escherichia coli*, *J Bacteriol* 1982; 150;214-220.
- Chaturvedi SK, Kumar Sand Dua RP. Chickpea breeding. In: Ali M et al (eds) Chickpea research in India. Indian Institute of Pulses Research, Kanpur, 2003; pp 69-98.
- Harding SA, Guikema JA and Paulsen GM. Photosynthetic decline from high temperature stress during maturation of wheat: II. Interaction with source and sink processes. *Plant Physiol.*, 1990; 92(3):654-658.
- Jumrani, K and Bhatia, VS. Impact of elevated temperatures on growth and yield of chickpea (*Cicer arietinum* L.). *Field Crop Res.*, 2014; 164:90-97.
- Miller PA, William C, Robinson HF and Comstock RE. Estimates of genotypic and environmental variances and covariance in upland cotton and their implications in selection. *Agron. J.* 1958; 50; 126-131.

6. Singh TP, Deshmukh PS, Srivastava GC, Kushwaha SR and Mishra SK. Growth rate of chickpea genotypes under different planting dates. Indian J. Plant physiol. 2005; 10: 254-259.
7. Singh TP, Deshmukh PS, Dutta M, Singh M, Kumari J, and Pragyaa. Morpho-physiological traits for tolerance of moisture and temperature stress chickpea (*Cicer arietinum* L.) genotypes. A correlation analysis. In A.K. Tripathi and H Pathak(Eds.), climate change and food security in india. The society of agricultural professionals 2011; (pp. 241-248). Kanpur: CSAUAT.
8. Tejpal- singh, deshmukh PS, Kushwaha SR. Physiological studies on temperature tolerance in chickpea (*Cicer arietinum* L.) genotypes. Indian J. plant physio. 2004; 9(3); 294-301.

UNDER PEER REVIEW