

## Original Research Article

### **“Efficacy of some insecticides, biopesticides and botanicals against shoot and fruit borer (*Leucinodes orbonalis* Guenee) on brinjal (*Solanum melongena* Linnaeus)”**

#### **Abstract**

A field experiment was conducted during *kharif* 2020 at Sakri Dist. Dhule Maharashtra (India). To determine ~~“Efficacy—efficacy of soame~~ insecticides, biopesticides and botanicals against shoot and fruit borer (*Leucinodes orbonalis* Guenee.) on brinjal (*Solanum melongena* Linnaeus.).<sup>2</sup> The efficacy result showed that the Emamectin benzoate 5% SG (10.35% and 15.91%) found most effective followed by Cypermethrin 25% EC (11.49 and 17.32%), Spinosad 45% SC (14.06 and 18.77%), NSKE 5% (15.43 and 20.31%), *Bacillus thuringiensis var kurstaki* 54% (15.96 and 22.30%) and *Beauveria bassiana*  $1 \times 10^8$  CFU (16.53% and 23.45%) was next effective treatment. Among all the treatments Garlic Bulb extract 3% (17.15% and 24.69%) is found be least effective but comparatively superior over the control. The best and most economical treatment was Emamectin benzoate 5% SG (1:3.3), followed by Cypermethrin 25% EC (1:3.1), Spinosad 45% SC (1:2.6), NSKE 5% (1:2.5), *Bacillus thuringiensis var kurstaki* (1:2.4), *Beauveria bassiana*  $1 \times 10^8$  CFU (1:2.1), Garlic Bulb extract 3% (1:1.9) as compared to control (1:1.1). Emamectin benzoate 5% SG (179q/ha) followed by Cypermethrin 25% EC (163q/ha).

**Keywords:** Biopesticides, botanicals, Emamectin benzoate, *Leucinodes orbonalis* Guenee and *Solanum melongena*.

#### **Introduction**

Vegetable cultivation is one of the most profitable and dynamic branch of agriculture. It has become an important source of income for both farmers and field labours, serving as a vehicle for reducing poverty in rural areas. Brinjal (*Solanum melongena* Linnaeus.) also known as eggplant is referred as the “King of vegetables” originated from India and now grown as a vegetable throughout the tropical, subtropical and warm temperate areas of the world. Brinjal is one of the most important common vegetable grown around the year under intensive cultivation

Formatted: Font: Not Bold

Comment [h1]: Add a concluding sentence

Formatted: Font: Not Bold

Comment [h2]: Introduction is too lengthy. Reduce its length to 1/3 or 1/4.

practices. The major constraint in cultivation of brinjal is infestation of shoot and fruit borer. (*Leucinodes orbonalis* Guenee.)

It is an important vegetable grown in all the seasons. Due to its nutritive value, consisting of minerals like iron, phosphorous, calcium and vitamins like A, B and C, unripe fruits are used primarily as vegetable in the country. It is also used as a raw material in pickle making and as an excellent remedy for those suffering from liver complaints. It has been reported as Ayurvedic medicine for curing the diabetes. In addition, it is used as a good appetizer, good aphrodisiac, cardio tonic, laxative and reliever of inflammation (Nawale et al., 2018).

Brinjal (*Solanum melongena* L.) is one of the popular vegetables favored by the people of many countries viz., Central, South and South East Asia, some parts of Africa and Central America (Harish et al., 2011). Apart from India, the other major brinjal growing countries are China, Turkey, Japan, Italy, Indonesia, Iraq, Syria, Spain, and Philippines. Brinjal is one of the most commonly grown vegetable crop in the country (Yadav and Tayde, 2018).

Amongst the solanaceous vegetables, brinjal is one of the most common, popular, principle annual crop grown in all the three seasons and economically important vegetable among small-scale farmers as it is a source of cash income for resource poor farmers. The major brinjal growing states in India are, Andhra Pradesh, Karnataka, West Bengal, Tamil Nadu, Maharashtra, Orissa, Uttar Pradesh, Bihar and Rajasthan. In India, West Bengal contributes highest area 181.5 million hectare and production 2877 million tonnes, Karnataka has highest productivity 25.4 million tonnes per hectare. In Uttar Pradesh, the area under cultivation of brinjal is 3430 hectare producing 111.70 MT and the productivity is 8 MT/ha (Yadav et al., 2015).

Though brinjal is a summer crop, it is being grown throughout the year under irrigated condition. Hence, it is subjected to attack by number of insect pests right from nursery stage till harvesting (Regupathy et al., 1997). Among the insect pests infesting brinjal, the major ones are shoot and fruit borer, *Leucinodes orbonalis* (Guenee), whitefly, *Bemisia tabaci* (Genn.), leaf hopper, *Amrasca biguttula* (Ishida), and non-insect pest, red spider mite, *Tetranychus macfurlanei*. Of these, *L. orbonalis* is considered the main constraint as it damages the

Formatted: Font: 12 pt, Not Bold

Formatted: Font: Not Bold

Formatted: Font: 12 pt, Not Bold

Formatted: Font: 12 pt, Not Bold

Formatted: Font: Not Bold

Formatted: Font: 12 pt, Not Bold

Formatted: Font: Not Bold

Formatted: Font: Not Bold

crop throughout the year. It is known to damage shoot and fruit of brinjal in all stages of its growth. The yield loss due to the pest is to the extent of 70-92 per cent (Reddy and Srinivas, 2004; Chakraborti and Sarkar, 2011; Jagginavar *etal* 2009). The infested fruits become unfit for consumption due to loss of quality and hence, lose their market value. It is also reported that there will be reduction in vitamin C content to an extent of 68 percent in the infested fruits (Hemi, 1955).

Formatted: Font: Not Bold

Damage to fruits particularly in autumn, is very severe and the whole crop can be destroyed. It alone causes damage as high as 85.90% and even up to 100% damage is also recorded. The larvae bore into tender shoots and cause wilting and dead heart and in later stage, they bore the tender fruits rendering them unfit for human consumption. So far, *L. orbonalis* Guenee is considered a major pest of brinjal as shoot and fruit borer in established crop in main field. It is also reported that there will be reduction in vitamin C content to an extent of 68 per cent in the infested fruits. It was reported that the shoot and fruit borer (on shoot) were more prevalent during vegetative phase of crop. The yield loss by this pest varied from 0.08-1.11 q/ha on the basis of inconsumable pest of damaged fruits and 0.46- 3.80 q/ha when whole of the damaged fruits were taken into consideration. It was reported that the borer infestation was 78.66% on top shoots in vegetative phase and then shifted to flowers and fruits with infestation reaching 66.66% in fruiting phase. (Nawaleet *al.*, 2018).

Comment [h3]: Too lengthy Introduction of the Brinjal. Just one paragraph is enough

Formatted: Font: Not Bold

Formatted: Font: Not Bold

The major pest of this economically important crop, farmers always apply the potent toxicant with unnecessarily high frequency during the total cropping duration. Thus, the insecticides which were potent to the pest at the beginning become ineffective due to the resistance development among the pest population. This has promoted the necessity for the development and use of new safer and more ecologically acceptable biorational insecticides as a part of IPM programs affecting specifically harmful pests, while sparing beneficial insect species and other organisms. Keeping this in view, the present study was carried on the field evaluation of effectiveness of biorational insecticides against *Leucinodes orbonalis* (Guenee) on brinjal and on yield (Devi *etal.*, (2015). (Alam *et al.*, 2003), pollution and health hazards (FAO, 2003). Again the question of residual toxicity of pesticides in brinjal is another big threat to our vegetable exports in the foreign markets (Islam *etal.*, 1999). Therefore, environment friendly

Comment [h4]: Add more references to support various statements.

Comment [h5]: Write correctly

management approaches to manage pests and diseases in agriculture using botanicals (Javaid *et al.*, 2020; Khan *et al.*, 2020; Jabeen *et al.*, 2021) and biocontrol agents (Javaid *et al.*, 2018; Khan and Javaid, 2022) are ~~must~~ highly necessary to reduce environmental and health hazards and to manage this pest eco-friendly. Various non-chemical approaches like biopesticides, botanicals, clean cultivation, mechanical control like hand picking and destroying of infested plant parts particularly shoots and fruits are common practices used for suppressing the insect pests (Hassan, 1994). Due to lack of knowledge and unavailability of non-chemical pest management approaches, growers of Bangladesh mostly depend on insecticides to keep the crop production steady. Appropriate knowledge and availability of botanical pest management approaches and their integration with selective chemicals may give better results against brinjal shoot and fruit borer. Considering the circumstances, the present study was conducted to evaluate the effectiveness of some eco-friendly management approaches against *L. orbonalis* in brinjal field. (Chowdhury *et al.*, 2017).

Formatted: Font: 12 pt, Italic

New generation biorational pesticide molecules have been claimed to be effective as well as safer for non-target organisms (Sontakke *et al.*, 2007; Misra, 2008). Journal of Entomology and Zoology Studies With a view on the climate change projections for India, an attempt has been made here to study the impact of the likely changes in weather factors in relation to shoot and fruit borer on brinjal crop under Western UP agroclimatic conditions. (Saran *et al.*, 2018).

Comment [h6]: Add objectives of the study

## Materials and methods

Comment [h7]: Divide into sub-headings

The experiment was conducted during the *khari*f season 2020. The sites selected uniform, cultivable with typical sandy loam soil having good drainage. The Brinjal seedlings of variety 'Gallan' were transplanted after 35 days at 60 cm x 60 cm spacing. The experiment was laid down in randomized block design (RBD) with eight treatments replicated thrice with each plot size of 2m x 2m and proper irrigation was provided. The treatments comprising of Emamectin benzoate 5% SG, Cypermethrin 25% EC, Spinosad 45% SC, NSKE 5%, *Bacillus thuringiensis var kurstaki* 54% and *Beauveria bassiana*  $1 \times 10^8$  CFU/Garlic Bulb extract 3% and were applied two times using knapsack sprayer in 15 days of interval. From each plot five plants were selected randomly and labeled for recording observations. As soon as

he infestation of pest was initiated, the observations on total number of larvae on brinjal on five observational plants from each treatment replication wise were recorded at 3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> days after imposing treatments. The data recorded in the different treatments were subjected to statistical analysis after suitable transformation by following standard procedures of RBD experiment. After harvesting of brinjal from each individual plots produce were calculated to work out the yield of the treatments. Yield of healthy heads was converted into quintal per hectare.

The observations on the larval population were recorded visually per plant from five randomly selected and tagged plants in each plot. The insecticides were sprayed at recommended doses when larval population reaches its ETL level. Larvae count was taken 24 hours before spraying at tagged plant at 5 tagged plants per treatment, which was further converted into per plant population and subsequent observation was recorded at 3<sup>rd</sup>, 7<sup>th</sup> and 14<sup>th</sup> days after spraying on same plants.

The healthy marketable yield obtained from different treatments was collected separately and weighted. The cost of insecticides and biopesticides used in this experiment was obtained from nearby market. The total cost of plant protection consisted of cost of treatment, sprayer, rent and labour charges for the spray. There are two sprays throughout the research period and the overall plant protection expenses was calculated. Total income was realized by multiplying the total yield per hectare by the prevailing market price, while the net benefit is obtained by subtracting the total cost of plant protection from the total income. Benefit over the control for each sprayed treatment was obtained by subtracting the income of the control treatment from that of each sprayed treatment.

## Results and discussion

The results after 1<sup>st</sup> and 2<sup>nd</sup> spray revealed that all the treatments were significantly superior to control in managing the pest infestation of *Leucinodes orbonalis* Guenee.) on brinjal. The data on the Percent (%) infestation of shoot borer on third, seventh and fourteenth day after spray revealed that all the treatments were significantly superior over control. Among all the treatments lowest Per cent (%) shoot, infestation was recorded in Emamectin benzoate 5% SG followed by Cypermethrin 25% EC followed by

**Comment [h8]:** 1-Improve language and formatting  
2- Add more Discussion points.

**Formatted:** Right: -0.03"

**Formatted:** Superscript

**Formatted:** Superscript

Spinosad 4.5% SC and NSKE 5%, then treatments *Bacillus thuringiensis* var. *kurstaki* 54% and *Beauveria bassiana*  $1 \times 10^8$  CFU followed by treatments Garlic bulb extract 3%, were found to be least effective but significantly superior over the control. Treatments T6 and T7 are non-significant to each other, Treatments T7 and T5 are non-significant to each other, Treatments T5, T1, T3 and T4 are non-significant to each other, T1, T3, T4 and T2 are non-significant to each other.

The data on the Percent (%) infestation of shoot borer on third, seventh and fourteenth day after second spray revealed that all the treatments were significantly superior over control. Among all the treatments lowest Per cent (%) shoot, infestation was recorded in Emamectin benzoate 5% SG followed by Cypermethrin 25% EC followed by Spinosad 4.5% SC and NSKE 5%, then treatments *Bacillus thuringiensis* var. *kurstaki* 54% and *Beauveria bassiana*  $1 \times 10^8$  CFU followed by treatments Garlic bulb extract 3%, were found to be least effective but significantly superior over the control. Treatments T6 and T7 are non-significant with each other, treatments T7 and T5 are non-significant with each other, treatments T5 and T1 are non-significant with each other and Treatments T3, T4 and T2 are non-significant with each other.

**Table 1 : Percent shoots and fruit infestation of *Earias vittella***

Comment [h9]: Add stats

Treatments		Percent shoots and fruit infestation of <i>Earias vittella</i>							
		First spray				Second spray			
		3 <sup>rd</sup> DAS	7 <sup>th</sup> DAS	14 <sup>th</sup> DAS	Mean	3 <sup>rd</sup> DAS	7 <sup>th</sup> DAS	14 <sup>th</sup> DAS	Mean
T <sub>0</sub>	Untreated	20.35	22.65	29.91	24.30	25.00	28.14	33.31	28.82
T <sub>1</sub>	NSKE 5%	15.94	14.20	16.14	15.43	20.70	18.00	22.23	20.31
T <sub>2</sub>	Garlic Bulb extract 3%	17.68	15.26	18.52	17.15	24.20	23.09	26.79	24.69
T <sub>3</sub>	<i>Bacillus thuringiensis</i> var. <i>kurstaki</i> 54%	16.69	14.52	16.68	15.96	21.91	20.85	24.15	22.30

T <sub>4</sub>	<i>Beauveria bassiana</i> 1×10 <sup>8</sup> CFU	17.31	14.54	17.76	16.53	22.65	21.65	26.04	23.45
T <sub>5</sub>	Spinosad 45% SC	14.46	12.84	14.88	14.06	19.68	16.63	20.00	18.77
T <sub>6</sub>	Emamectin benzoate 5% SG	11.20	8.84	11.01	10.35	17.11	12.88	17.73	15.91
T <sub>7</sub>	Cypermethrin 25% EC	11.54	10.01	12.92	11.49	18.28	14.82	18.85	17.32
Overall Mean		15.64	14.11	17.23	15.66	21.19	19.51	23.64	21.44
F-test		S	S	S	S	S	S	S	S
C.D		1.678	1.452	1.833	2.915	1.644	1.440	1.835	2.558
SE(m)		0.548	0.474	0.598	0.952	0.537	0.470	0.599	0.835
SE(d)		0.775	0.671	0.846	1.346	0.759	0.665	0.847	1.181
C.V		6.065	5.821	6.015	10.525	4.387	4.173	4.389	6.744

## CONCLUSION

Treatment Emamectin benzoate was found to be most effective when considered from the point of view of percent shoot and fruit damage and benefit: cost ratio BCR (if all the treatments produced were sold at equal price). Emamectin benzoate belongs to the chemical group semi-synthetic Avermectin. The specific mode of action of Emamectin benzoate is to inhibit muscle contraction, causing a continuous flow of chloride ions in the GABA and H-Glutamate receptor sites. The resistance risk is assessed to be low to moderate because of the current pest situation and the intended use of Emamectin benzoate in India and it was one of the insecticides that showed negligible effect on predators, pollinators and parasitoids. This selective beneficial to arthropods makes Emamectin benzoate a strong tool for

Comment [h10]: Can be improved

Integrated Pest Management programs.

## References:

Alam, S. N., Rashid, M. A., Rouf, F. M. A., Jhala, R. C., Patel, J. R., Satpathy, S., Shivalingaswamy, T. M., Rai, S., Wahundeniya, I., Cork, A., Ammaranan, C. and Talekar N.S. (2003). Development of an integrated pest management strategy for eggplant fruit and shoot borer in South Asia. Shanhua, Taiwan. *Asian Vegetable Research and Development Center Publication*, 28 (3), 548-56.

**Chakraborti, S. and Sarkar, P. K. (2011).** Management of *Leucinodes orbonalis* Guenee on eggplants during the rainy season in India. *Journal of Plant Protection Research*. 51 (4): 325-328.

**Chowdhury, M. A. H., Alam, M. Z., Hossain, M. S., Suh, S. J., and Kwon, Y. J. (2017).** Management of brinjal shoot and fruit borer *Leucinodes orbonalis* Guenee using different approaches. *Bangladesh Journal of Agricultural Research*, 42(3): 539-548.

**Devi, L. L., Ghule, T. M., Chatterjee, M. L., and Senapati, A. K. (2015).** Effectiveness of biorational insecticides for the management of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee and on yield. *Ecology, Environment and Conservation*, 21(2): 783-788.

## FAO-

**Food and Agricultural Organization (2003).** Inter-country programme for integrated pest management in vegetables in South and South-East Asia. Eggplant integrated pest management: An ecological guide. Rome, Italy. 177.

**Harish, D. K., Agasimani, A. D., Imamsaheb, S. J., and Patil, S. S. (2011).** Growth and yield parameters in brinjal as influenced by organic nutrient management and plant protection conditions. *Research Journal of Agricultural Sciences*. 2(2): 221-225.

**Comment [h11]:** Format uniformly and also add a few latest references of 2021 and 2022

**Formatted:** Font: Not Bold

**Hemi, M. A. (1955).** Effect of borer attach on the vitamin 'C' content of brinjal. *Pakistan Journal of Health*.4: 223-224.

**Hassan, S. A. (1994).** Strategies to select *Trichogramma* species for use in biological control. In: Biological control with the gregarious parasitoids, (Eds.): E. Wajin bergand S.A.Hassan. Oxon, U.K.: CAB International. 557.

**Islam, M. S., Choudhury, A. R., Maleque, M. D., Mondal, H. K. and Khan, A. U. (2019).** Management of brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee.) using selected bio-rational insecticides. *Fundamental and Applied Agriculture*.4(4): 1025–1031.

**Jagginavar, S.B., Sunitha, N.D. and Biradar, A.P. (2009).** Bioefficacy of flubendiamide 480 SC against brinjal fruit and shoot borer, *Leucinodes orbonalis* Guenee. *Karnataka Journal of Agricultural Sciences*. 22(3):712-713.

[Jabeen, N., A. Javaid, A. Shoaib and I.H. Khan \(2021\). Management of southern blight of bell pepper by soil amendment with dry biomass of \*Datura metel\*. \*Journal of Plant Pathology\* 103\(3\): 901-913.](#)

[Javaid, A., I.H. Khan and A. Shoaib \(2018\). Management of charcoal rot of mungbean by two \*Trichoderma\* Javaid A. species and dry biomass of \*Coronopus didymus\*. \*Planta Daninha\* 36: e018182795.](#)

[Javaid, A. R. Munir, I.H. Khan and A. Shoaib \(2020\). Control of the chickpea blight, \*Ascochyta rabiei\*, with the weed plant, \*Withania somnifera\*. \*Egyptian Journal of Biological Pest Control\* 30:114.](#)

[Khan IH, Javaid A, Al-Taie AH, Ahmed D \(2020\). Use of neem leaves as soil amendment for the control of collar rot disease of chickpea. \*Egyptian Journal of Biological Pest Control\* 30:98.](#)

[Khan, I.H. and A. Javaid \(2022\). DNA cleavage of the fungal pathogen and production of antifungal compounds are the possible mechanisms of action of biocontrol agent \*Penicillium italicum\* against \*Macrophomina phaseolina\*. \*Mycologia\* 114\(1\): 24-34.](#)

**Misra, H. P. (2008).** New promising insecticides for the management of brinjal shoot and fruit borer, *Leucinodes orbonalis* Guenee. *Pest Management*

Formatted: Font: 11 pt, Not Bold

Formatted: Font: 11 pt, Not Bold

Formatted: Font: 11 pt, Not Bold

Formatted: Font: 11 pt, Not Bold

Formatted: Font: 11 pt, Font color: Black

Formatted: Font: 11 pt, Not Bold

Formatted: Font: 11 pt, Font color: Black

Formatted: Indent: Left: 0.1", Hanging: 0.1", Right: 0.07", Space Before: 8.25 pt, Line spacing: 1.5 lines, No bullets or numbering, No widow/orphan control, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers, Tab stops: Not at 0.38"

Formatted: Font: Not Bold

Formatted: Portuguese (Brazil)

Formatted: Font: 11 pt, Portuguese

Formatted: Right: 0.07"

*Horticultural.Ecosystems*.14(2):140-147.

- Nawale, J. S., Kumar, A. Patil, A. A. and Narode, M. K. (2018).** Efficacy of certain insecticides and bio-pesticides against brinjal shoot and fruit borer (*Leucinodes orbonalis* Guenee). *Journal of Entomology and Zoology Studies*, 6(5):292-295.
- Regupathy, A., Palanisamy, S., Chandramohan, N. and Gunathilagaraj, K. (1997).** A guide on crop pests. Sooriya desktop publishers, Coimbatore, 264.
- Reddy, E. and Srinivasa, S. G. (2004).** Management of shoot and fruit borer, (*Leucindoesorbonalis* Guenee) in brinjal using botanicals/oils. *Pestology*, 28: 50-52.
- Sontakke, B. K., Das, N., Panda, P. K., Swain, L. K. (2007).** Bio-efficacy of Emamectin benzoate 5% SG against fruit and shoot borer in okra. *Journal of Plant Protection and Environment*. 4(2):30-33.
- Saran, S., Singh, D. V., Singh, A., Kumar, S. and Kumar, U. (2018).** Bio-efficacy of selective eco-friendly insecticides and biopesticides against shoot and fruit borer, (*Leucinodes orbonalis* Guenee) in brinjal. *Journal of Pharmacognosy and Phytochemistry*, 7(6): 1690-1694.
- Yadav, R.V. and Tayde, A.R. (2018).** Comparative efficacy of cypermethrin with combination of different botanicals against brinjal shoot and fruit borer (*Leucinodes orbonalis*) on Brinjal. *Journal of Entomology and Zoology Studies*. 6(3): 1675-1677.
- Yadav, R., Lyall, H., Kumar, S. And Sanp, R. K. (2015).** Efficacy of certain botanical insecticides against shoot and fruit borer, *Leucinodes orbonalis* (Guenee) on brinjal (*Solanum melongena* L.). *An International Quarterly Journal of Life Sciences*. 10(2):987-990.

UNDER PEER REVIEW

