

Vermiwash : A Potential tool for crop production in organic agriculture

ABSTRACT

In India, the first green revolution enormously enhanced ~~the~~ crop production, but on the other hand, massive application of chemical fertilizers and pesticides over a period has resulted in poor soil health, pesticide toxicity, heavy disease and pest incidences, and environmental pollution. Scientists and policy planners are, therefore, reassessing agricultural practices which relied more on biological inputs rather than heavy usage of chemical fertilizers and pesticides. Organic farming can provide quality food without adversely affecting ~~the~~ soil health and the environment. Vermiwash is emerging as an important potential tool for promotion of organic agriculture.

Vermiwash is a brown coloured, odorless, liquid extract produced from vermicompost in a medium where earthworms are richly populated. It contains mucus, excretory products of worms, and various concentrations of macro, micro, and beneficial nutrients along with beneficial microorganisms, growth hormones, vitamins, enzymes, and amino acids, therefore is a good source for plant nutrition used as foliar spray in sustainable agriculture.

Comment [11]: the

Comment [12]: -

Comment [13]: a

Keywords: Vermiwash, vermicompost, potential tool, vermiculture, ~~and~~ organic agriculture

1. Introduction

Conversion of biodegradable waste into enriched compost is possible through the technology known as vermicomposting. Earthworms consume raw materials and excrete it in a digested form called worm cast which is rich in nutrients, growth promoting substances, beneficial soil microflora, etc. (Das et al., 2014). Accumulation of wastes is a serious problem which needs to be managed to make environment free from pollution. Vermiwash helps to manage market wastes in a profitable manner. Vermiwash is an eco-friendly liquid organic fertilizer obtained from units of vermiculture/vermicompost in the form of drainage. The vermiwash is formed due to the movement of water in the vermicompost unit from the increased moisture content due to the heat generated during the vermicompost process. This water absorbs the earthworm body secretions, enzymes, plant growth hormones, and other nutrients from the organic waste to form the vermiwash which can be easily absorbed by plants (Sundaravadivelan et al., 2011). Presently, in organic farming systems, vermiwash plays a key potential for sustainable crop production. This vermiwash contains enzymes, secreted from earthworms which would stimulate the growth and yield of agricultural crops

Comment [14]: In the introduction section there is no subtitle, but it is written based on an explanation that describes the importance of the problems described in the title and the problems to be solved.

Comment [15]: I.

Comment [16]: them

Comment [17]: -

Comment [18]: the

Comment [19]: in

and even develop resistance in crops. Vermiwash also contains the soluble plant nutrients apart from some organic acids and mucus of earthworms and microbes (Sundararasu, 2016). It is a very good plant tonic which can be used for foliar spray. There is no special device required to collect the vermiwash except for a tap which is fitted at the bottom of the containers where earthworms are cultured. Even during the period of normal management of the vermiculture, water is sprayed regularly to maintain adequate moisture and excess water is drained which contains some essential plant nutrients. The composition of vermicompost and vermiwash prepared from the same organic matter are essentially the same (Gudeta *et al.*, 2021). Vermiwash plays an important role in the plant growth and development, contribute to the initiation of rooting, root growth, enhancement of crop production, improvement in soil organic matter and facilitate nutrient availability to crops, resulting in good crop yield. Vermiwash can be used for crop production as both foliar spary and soil application. Presently, foliar application is the current trend of sustainable crop production to fulfill the target of production with a low quantity of manure (Rajasooriya and Karunarathna, 2020). It was found that spraying of vermiwash on vegetables markedly improved both the quality and quantity of the yield (Sundararasu, 2016). Further, its production cost is less expensive as compared to chemical fertilizer and very ease to produce. Besides its application as organic fertilizer to enhance crop productivity, it can also be applied in disease suppression and pest control due to the presence of essential antimicrobial and anti-pest chemicals (Kanchan *et al.*, 2013; Thakur and Sood, 2019).

Comment [I10]: that

Comment [I11]: the

Comment [I12]: the

Comment [I13]: ing

Comment [I14]: the

Comment [I15]: ing

Comment [I16]: y

2. Vermiwash and its composition

Comment [I17]: II.

Vermiwash is a brown colored, odorless, liquid extract and a by-products parts of vermiculture/vermicompost in the form of drainage that is collected after the channel of water through a column of worm action of vermiculture and vermicomposting and is very beneficial as a foliar spray (Jaysawal *et al.*, 2020). It is a worm coelomic fluid extract containing several enzymes (cocktails of *proteases*, *amylases*, *urease* and *phosphatase*), plant growth hormones (cytokinin, auxin), vitamins, amino acids, macro and micronutrients (Nayak *et al.*, 2019) along with excretory substances and mucus secretion of earthworms, therefore is a good source for plant nutrition.

It contains total solids (2448 mg/L), volatile solids (738 mg/L), silica (8 mg/L), auxin (0.98 µg/L) and cytokinin (0.68 µg/L) (Patil *et al.*, 2007). Microbial study of vermiwash found the presence of many useful micro-organisms such as heterotrophic fungi,

actinomycetes, bacteria including nitrogen-fixing bacteria like *Azotobacter* spp., *Agrobacterium* spp., and *Rhizobium* spp. and phosphate solubilising bacteria (Zambare *et al.*, 2008). The nutrient composition of vermiwash is presented at Table 1 (Nayak *et al.*, 2019).

Comment [I18]: z

Comment [I19]: in

Table 1. Nutrient analysis of vermiwash

Sl. No.	Parameters	Value
1	pH	7.39-7.5
2	EC	0.008±0.001
3	Organic carbon	0.25 ± 0.03%
4	Nitrogen	0.01-0.001%
5	Phosphorus	1.70%
6	Potassium	26 ppm
7	Sodium	8 ppm
8	Calcium	3 ppm
9	Copper	0.01 ppm
10	Iron	0.06 ppm
11	Magnesium	160 ppm
12	Manganese	0.60 ppm
13	Zinc	0.02 ppm
14	Total heterotrophs (cfu/ml)	1.79×10^3
15	Nitrosomonas (cfu/ml)	1.01×10^3
16	Nitrobacter (cfu/ml)	1.12×10^3
17	Total fungi (cfu/ml)	1.46×10^3

3. Vermiwash preparation

Vermiwash is the liquid extracted from vermicompost, which is prepared by feeding earthworms with raw materials like leaf litter or cow dung or other organic materials. Two exotic species *Eisenia fetida* and *Eudrilus eugeniae* are widely used in India for production of vermicompost/vermiwash.

Comment [I20]: 2.1 Vermiwash Preparation and Production

3.1 Vermiwash production from vermicompost pit

Comment [I21]: delete

The vermipits were prepared with (2m x1m x0.75m) (length x breadth x height) with the arrangement of layers performed from bottom to top as shown in fig. 1. The pebbles or coconut shell filled on the bottom layer to absorb excess water from the composting pit. The second layer filled with sandy soil to prevent the accumulation of extra water in the medium and the third layer contained organic soil and old compost inoculated with earthworms. Cow dung and leaf litter blended at the ratio of 1:2 and added to the pit in the fourth layer. Finally, on the top layer, the gunny bags spread on the top of the pit to prevent from direct sunlight and keep the medium moist enough. The contents of the pit turned once in 2 or 3 days to enhance aeration in the course of vermicomposting. After 30 days of the composting process, vermiwash started to collect at the bottom of the vermicompost pit. The water that passes

Comment [I22]: was

through the vermiculture/vermicompost, resulting in washing of the live and dead earthworms, soil microorganisms and decomposed organic matter, carries all these dissolved substances. As the water can dissolve some vermi casts which contains lots of nutrients, they find their way into vermiwash.

Comment [I23]: the




Fig 1: Model of vermicompost pit

Comment [I24]: include the reference or image source

3.2 Setting up of a vermiwash unit for commercial production

Comment [I25]: 2.2

(ECO – SCIENCE research foundation method)

Vermiwash units can be set up either in barrels or in buckets or even in small earthen pots. The procedure explained here was standardised by Ismail (1997).

Materials needed

- A plastic barrel (250 lt) or small bucket
- broken stones
- coarse sand
- garden soil
- earthworms
- cattle dung
- straw and



- water

Production process

Fig 2: Vermiwash unit (in buckets) for commercial production




Fig 3: Vermiwash unit (in small earthen pots) for commercial production

Comment [I26]: include the reference or image source

The empty plastic barrel with one side open is taken and the base of the barrel is fitted with a tap to collect vermiwash. The barrel is filled with different successive layers (Fig. 2). First base layer is filled with medium sized bricks or stones up to a height of 10-15 cm or 10-15% of the container. Above the base layer, a layer of coarse sand (up to 6 inches) and fine sand (5 inches) are spread. After that, a layer of partially decomposed cow dung (20-25 cm) is filled. Introduce of 1000 -1500 juvenile or adult earthworms (*Eisenia foetida* / *Eudrilus eugeniae*) with fertile soil into the container. Then decomposed organic residues of 40-45 cm of the container were loosely placed in the plastic barrel. On the top layer dry leaves, straw, etc were placed as additional raw material that mainly prevents direct exposure from sunlight. All the layers in the container is moistened by sprinkling water. Container is sprinkled with approximately 2 lit of water per day. After 16 to 20 days vermiwash will be produced in the barrel. Every day about 3-4 lit of vermiwash will be collected from this unit. Hang one pot with a bottom hole over the plastic barrel in a way so that waterfall drop by drop. Every day 4-5lt of water is poured into the hanging pot. Keep another pot under the tap and collect 3-4lt of vermiwash every day. The tap of the unit is opened the next day morning and the vermiwash is collected. The entire setup may be emptied and reset between 10 and 12 months of use. The collected vermiwash must be stored in a cool and dry place. For a nutrient

supplement, vermiwash (1:10) i.e @ 1 litre is generally mixed with 10 litres of water and the solution is sprayed on entire parts of the growing crops. However, to control various pest and diseases of crop, vermiwash @ 1 litre thoroughly mixed with 1 litre of cow urine and 10 liters of water and is kept overnight and sprayed on the affected crops (Nayak *et al.*, 2019).

4. Application methods of vermiwash

Comment [I27]: 2.3

Vermiwash alone or mixed with cow urine (One liter vermiwash + one liter cow urine+ eight liters water) is also an excellent growth promoter. Before the application of vermiwash as foliar application on any plants, it should be diluted with water (10 percent). Das *et al.*, 2014 reported that before transplanting, seedlings are dipped in vermiwash solution for about half an hour after diluting it with water (5 times). When vermiwash liquid is added to compost pits, it enhances the degradation process.

5. Benefits of vermiwash

Comment [I28]: 2.4

Being an excellent nutrient supplement, it enhances soil the physico-chemical properties and maintains soil fertility in a long run. It improves soil aeration, soil tilth and texture thereby reducing the compaction. It also enhances water holding capacity of soil and improves soil fertility status. It can also be added in the compost pit to hasten the degradation process. Vermiwash is known to play a major role in the plant growth and development, contribute to root initiation, root growth, plant development, promotion of growth rate and improvement in crop production, increases nutrient uptake by crops resulting in good crop yield (Nayak *et al.*, 2019). Besides serving as a plant growth promoter, vermiwash has tremendous biopesticide properties. It is highly toxic against insect-pest survival and increases the disease resistance power of the crop (Sayyad, 2017). Now a days, it is being popularised as liquid manure. It increases the rate of photo synthesis in crops/plants. It also increases the number of micro-organisms in the soil which helps in decomposing soil organic matter (Tripathi *et al.*, 2005).

6. Effect of vermiwash on plant growth and yield

Comment [I29]: 2.5

It was found that foliar spray of vermiwash significantly increased the growth and yield parameters of brinjal (Sundararasu and Jeyasankar, 2014). Verma *et al.*, [2017] reported that application of 100% RDF along with vermiwash @100 lt ha⁻¹ produced distinctly higher plant height, number of primary and secondary branches, leaf area index and dry matter accumulation. The plant height, leaf area index, pod length and girth, fresh weight of the pods, 100 seed weight and yield per plant of greengram were recorded highest due to application of 50% RDF along with foliar spray of vermiwash @75% (Rajasooriya and

Comment [I30]: parenthesis

Karunaratna 2020). Yassen, 2020 reported that the growth parameters of lettuce were significantly higher due to application of vermicompost @ 4t/ha with foliar spray of vermiwash @ 150 ml/lit .

Comment [I31]: parenthesis

It was found that application of vermiwash at Low concentration is effective in bringing about seed germination and vigour seedling growth (Fathima and Sekar, 2014). This could be attributed due to presence of nutrients and growth promoting substances in vermiwash which showed its potentiality in seed germination and seedling vigour (Chattopadhyay, 2015).

7. Effect of vermiwash on soil properties

Comment [I32]: 2.6

Vermiwash contains micro and macro nutrients, hormones which promote plant growth and yield and improves soil fertility status (Sharma *et al.*, 2005). Use of Organic formulations in agriculture could be a dynamic source to move forward soil fertility (Verma *et al.*, 2017). Combination of vermicompost and vermiwash recorded a significant influence on the biochemical characteristics of the soil with marked improvement in soil micronutrients and better qualitative improvement in the physical and chemical properties of the soil (Nayak *et al.*, 2019). Tripathi *et al.* (2005) suggested that vermiwash increases the number of microorganisms in the soil which helps in the decomposition of organic matter.

Comment [I33]: or 2004? see reference and check the original article

Sundararasu and Jeyasankar, 2014 reported that application of vermiwash significantly increases the macro and micro nutrient status of soil.

8. Conclusion

Comment [I34]: III.

Vermiwash seems to possess an inherent property which acts not only as a liquid organic biofertilizer which promot growth and yield of plants but also acts as a mild biopesticide. So, it can be used as a potent input in organic farming and sustainable crop production for both soil health and insect, pest and disease management. Vermiwash is now effectively used in organic agriculture for both soil health and disease management for sustainable crop production and to prevent environment and human health.

Comment [I35]: that

Comment [I36]: promotes the

Comment [I37]: environmental

References

- Chattopadhyay, A. (2015) Effect of vermiwash of *Eisenia foetida* produced by different methods on seed germination of green mung, *Vignaradiata*. *International Journal of Recycle Organic Waste Agriculture*. 4 (4): 233-237.
- Das SK, Avasthe RK and Gopi R. (2014) Vermiwash: Use in Organic Agriculture for Improved Crop Production, *Popular Kheti*, 2 (4): 45-46.

- Fathima M and Sekar M. (2014) Studies on Growth Promoting effects of Vermiwash on the Germination of Vegetable Crops. *International Journal of Current Microbiology and Applied Sciences*. 3(6):564-570.
- Gudeta K, Julka JM, Kumar A, Bhagat A and Kumar A. (2021) Vermiwash: An agent of disease and pest control in soil, a review, *Heliyon*, 7: e06434.
- Ismail, S. A. (1997) *Vermiculture: Biology of Earthworm*. Orient Longman, Hyderabad, India.
- Jaysawal, P.K., Arun, D.P., Verma, S.K., Rakshit, A., Chattopadhyay, A. and Pandey, P. (2020) Vermiwash: Key for the Crop Production in Organic Agriculture, *International Journal of Bioresource Science*, 7(1): 21-23.
- Kanchan, M., Keshav, S., Tripathi, C.P.M. (2013) Management of pod borer (*Helicoverpa armigera*) infestation and productivity enhancement of gram crop (*Cicer aritenium*) through vermiwash with biopesticides. *World J. Agric. Res.* 9 (5), 401–408.
- Nayak H, Rai S, Mahto R, Rani P, Yadav S, Prasad SK and Singh RK. (2019) Vermiwash: A potential tool for sustainable agriculture, *Journal of Pharmacognosy and Phytochemistry*, SP5: 308-312.
- Patil SS, Kengar SB, Sathe TV. (2007) New vermiwash model for sustainable agriculture in India, *Nature Environment and Pollution Technology*, 6(2):281-284.
- Rajasooriya AP Sonali and Karunarathna B. (2020) Application of vermiwash on growth and yield of green gram (*Vigna radiata*) in sandy regosol, *AGRIEAST*, 14(2): 31-42
- Sayyad NR. (2017) Utilization of vermiwash potential against insect pests of tomato. *International Research Journal of biological Sciences*, 6(1):44-46.
- Sharma AK. (2004) *A Handbook of Organic Farming*. Agrobios, Jodhpur.
- Sundararasu K (2016) Vermiwash: Effect of vermiwash on growth and yielding pattern of selected vegetable crop Chilli, *Capsicum annum*, *International Journal of Advanced Research in Biological Sciences*, 3(9): 155-160.
- Sundararasu, K. and Jeyasankar, A. (2014) Effect of vermiwash on growth and yield of brinjal, solanum melongena (eggplant or aubergine) *Asian Journal of Science and Technology*, 5(3), 171-173
- Sundaravadivelan C, Isaiarasu L, Manimuthu M, Kumar P, Kuberan T and Anburaj, J. (2011) Impact analysis and confirmative study of physico-chemical, nutritional and biochemical parameters of vermiwash produced from different leaf litters by

Comment [138]: in manuscript 2005, check the original article

using two earthworm species”, *Journal of Agricultural Technology*, 7 (5), 1443-1457.

Thakur, S., Sood, A.K. (2019) Lethal and inhibitory activities of natural products and biopesticide formulations against *Tetranychus urticae* Koch (*Acarina: Tetranychidae*). *Int. J. Acarol* 45 (6-7), 381–390.

~~Tripathi YC, Hazarika P and Pandey BK. (2005) Vermicomposting: An ecofriendly approach to sustainable agriculture. In: Arvind Kumar (eds), *Verms and vermitechnology*. APH Publishing Corporation, New Delhi. pp23-39.~~

Tripathi YC, Hazarika P, Pandey B.K. (2005) Vermicomposting: An ecofriendly approach to sustainable agriculture, In: Arvind Kumar (eds), *Verms and vermitechnology*, APH Publishing Corporation, New Delhi, 23-39.

Verma S, Singh A, Pradhan SS, Singh RK, Singh JP. (2017) Bio-efficacy of Organic Formulations on Crop Production-A Review, *Int. J. Cur. Microbiol. App. Sci*, 6(5):648-665

Yassen A.A, Entsar M Essa, Neama M. Marzouk and Sahar M. Zaghoul (2020) Impact of vermicompost and foliar spray of vermiwash on growth, yield and nutritional status of lettuce plants, *Plant Archives*, 20 (1) :449-455

Zambare VP, Padul MV, Yadav AA, Shete T. B. (2008) Biochemical and Biological approach as eco-friendly soil conditioner. *ARPJ Journal of Agricultural and Biological Sciences*. 3(4):28-37.