

**Identification of new four races of barley stem rust (*Puccinia graminis* f. sp. *tritici*) for
the first time in Egypt**

ABSTRACT

Four pathogenic races belonging to *Puccinia graminis* f.sp. *tritici* Eriks and Henn, were genetically identified in Egypt for the first time on barley, using the molecular biology method, using specific ITS primers (PCR) and Fingerprinting using (RAPD) markers, these races were recorded, during the present study, in the gene bank under accession numbers MW 931757, 931758, 931759 and 931760. Also, five Egyptian barley varieties, *i.e.* Giza 123, Giza 124, Giza 125, Giza 126 and Giza 2000 were evaluated for their resistance to stem rust and some vegetative traits under field conditions in Sids, Giza, Nubariya and Sakha agricultural research stations during growing season 2020/2021. All tested barely cultivars were resistant to the pathogen races. Cultivar Giza 125 exhibited an earliness for heading and maturity, while cv. 2000 exhibited the highest values for spike length, grain numbers/spike, grain weight, biological and grain yield.

Keywords: Barley, stem rust, PCR, RAPD markers, yield components.

20

INTRODUCTION

Barley is one of the most abundantly utilized cereal crops over the world. It accounts for 23% of total global cereal production, ranking fourth after wheat, rice, and maize. In Egypt, barley grows in the Northern Coastal Regions and new reclaimed lands.


Barley stem rust disease caused by *Puccinia graminis* f.sp. *tritici* Eriks and can affect wheat, triticale and many other related grasses. It is found wherever temperate climate that favors growing of cereals [1]. The alternate hosts, out of Egypt, are *Berberis* and *Mahonia* species [2].

29 Rusts, are the most important common diseases of barley and wheat because of the
ability to spread for long distance, in addition to their ability to produce new races that can
attack resistant varieties and their potential to develop rusts rapidly high under optimal
environmental conditions and cause serious losses [3,4]. Stem rust considered as one of the
most dangerous diseases among the other three barley rust diseases. The causal organism
under the suitable conditions may destroy the whole plants with no yield seeds. In Egypt, high
losses in wheat grain yield can be expected due to the suitable environmental conditions for
disease infection. On the contrary, barley varieties have a high degree of resistance [5]. Stem
rust epidemics in Minnesota, North Dakota and South Dakota caused average yield losses
over 20% [6,7]. A dangerous epidemic of stem rust disease damaged wheat crops in the
Southern states of Australia in 1974 [8]. In this study, identifying the races of stem rust
caused by *Puccinia graminis* f.sp. *tritici* that infects barley using molecular biology methods
is of great importance. Also, definition of appropriate locations for the pathogen spread,
definition of barley resistant varieties and using them in breeding programs with excluding
the susceptible lines reduces the losses and spread of the disease in the future.

MATERIALS AND METHODS

Isolation, purification and identification of the physiological races:

46 Symptoms of stem rust disease were observed on barley plants (*Hordeum vulgare* L.)
grown in El-Alameen County–Marsa Matrouh Governorate, during March/2019 growing
season, specifically, on line which its pedigree is (Lignee 527/Gerbel/3/Boyb*2/Surb//C
11229.2D/4/M104) (Fig.1).



**Figure (1): Symptoms of barley stem rust caused by *Puccinia graminis* f. sp. *tritici* on barley
LBYT line showing the uredial pustules on the stem.**

Stem rust fungus isolation was carried out from a single uredial pustules (Four isolates), multiplied and the resulted spores were used to carry out the pathogenicity test on the same barley line under greenhouse conditions at Plant Pathology Research Institute, A.R.S, Giza, Egypt.

Establishment of single –pustule isolates:

Stem rust urediospores were transferred from infected samples with a sterile scalpel and transferred to upper leaf-surface of 7-days old seedlings on line which its pedigree is "Line 527/Gerbel/3/Boyb*2/Surb//C11225.2D/4/M104" [9,10]. The inoculated barley plants were directly incubated in dew chambers at 100% relative humidity for 20-24 h in the dark chambers. Inoculated seedlings returned back to the greenhouse benches. Developed individual pustules were sub cultured on leaves of healthy seedlings to propagate and generate sufficient inocula of each single-pustule [11]. Urediniospores of *P. graminis* f.sp. *tritici* of each isolate were collected and DNA was extracted from 0.50 mg urediniospores

Molecular identification:

Extraction and purification of genomic DNA:

Genomic DNA of the fungal was extracted from each four samples using a DNeasy MiniKit (Qiagen, CA, USA), according to [12].

ITS rRNA analysis:

a) PCR Reactions:

The PCR amplification performed in a total volume of 50 µl, containing 1X reaction buffer, 1.5 mM MgCl₂, 1U *Taq* DNA polymerase (promega), 2.5mM dNTPs, 30 pmol of each primer and 30 ng genomic DNA (Table 1) [13].

Table (1): Primer ITS forward and reverse, sequence and product size.

| Primer code | Sequence | Product Size |
|-------------|-----------------------------|--------------|
| (ITS-1) F | 5'- TCCGTAGGTGAACCTGCGG -3' | 600bp |

b) Thermo-cycling PCR program:

80 PCR amplification performed in a Perkin-Elmer/GeneAmp® PCR System 9700 (PE Applied Biosystems) programmed to fulfill 40 cycles after an initial denaturation cycle for 5 min at 94°C. Each cycle consisted of a denaturation step at 94°C for 30 sec., an annealing step at 53°C for 30 sec. and an elongation step at 72°C for 1 min. The primer extension segment was extended to 7 min at 72°C in the final cycle.

c) Detection of the PCR Products:

86 The intensification items settled by electrophoresis in a 1.5% agarose gel containing ethidium bromide (0.5ug/ml) in 1X TBE buffer at 95 volts. A 100bp DNA step was utilized as a genetic measure standard. PCR items were visualized on UV light and captured employing a Gene Documentation Framework (BIO-RAD 2000).

Purification of PCR Products:

91 Amplified for all PCR products were purified using EZ-10 spin column PCR products purification PCR reaction mixture was transferred to 1.5 ml microfuge tube and three volumes were added of binding buffer 1 after that the mixture solution was transferred to the EZ-10 column and let it stand at room temperature for 2 minutes after that centrifuge, 750 µl of wash solution were added to the column and centrifuged at 10.000rpm for two minutes, repeated was 10.000 rpm was spine for an additional minute to remove any residual wash solution. The column was transferred into a clean 1.5 ml microfuge tube and 50 ul of elution buffer were added, incubated at room temperature for 2 minutes and then store purified DNA at -20°C. [14].

ITS sequencing analysis:

101 The sequencing of the product PCR was conducted in an automatic sequencer ABI PRISM 3730XL analyzer by using big dye TM terminator cycle sequencing kits following supplier protocols of the manufacturer. Single-pass sequencing performed on each template using forward primer. The fluorescent-labeled fragments were purified from the

105 incorporated terminators with an ethanol precipitation protocol. The samples were
106 resuspended in distilled water and subjected to electrophoresis in an ABI 3730xl sequencer
(107 Microgen Company).

Computational analysis (BLASTn) ITS:

109 The sequences were analyzed using BLAST program
(110 <http://www.ncbi.nlm.nih.gov/BLAST>). Sequences were aligned using Align Sequences
111 Nucleotide BLAST.

112

RAPD fingerprinting:

114 **RAPD** has been successfully used for the fingerprint of four fungal isolates. This
115 method uses random sequence primers of about 10 bases in length that hybridize with
116 chromosomal DNA. The amplification reaction was carried out in 25 µl reaction volume
117 containing 1X PCR buffer, 2 mM MgCl₂, 0.2 mM dNTPs, 25pmol primer, 1 U Taq DNA
118 polymerase and 30ng templates DNA. PCR amplification was performed in a Perkin-
119 Elmer GeneAmp® PCR System 9700 (PE Applied Biosystems) programmed to fulfill 35
120 cycles after an initial denaturation cycle for 5 min at 94°C. Each cycle consisted of a
121 denaturation step at 94°C for 45s, an annealing step at 36°C for 50s, and an elongation step at
122 72°C for 1min. The primer extension segment was extended to 7 min at 72°C in the final
123 cycle. The amplification products were resolved by electrophoresis in a 1.5% agarose gel
124 containing ethidium bromide (0.5ug/ml) in 1X TBE buffer at 95 volts.

The test materials:

126 Five commercial barley varieties, *i.e.* Giza 123, Giza 124, Giza 125, Giza 126, Giza
127 and the check line (LBYT, highly susceptible) (Table 2) were tested to determine their
128 resistance to stem rust. The field trials were conducted under natural infection at Sids, Giza,
129 Nubaria and Sakha Agricultural Research Stations, during growing season 2020/2021. All
130 experiments were carried out in a randomized complete block design with three replicates,
131 each was 3m x 3.5m = 10.5 m² plot size, the grains of the tested barley varieties were sown in
132 6-row plot.


Table (2): The Egyptian barley cultivars evaluated throughout growing season 2020-2021 and their pedigree.

| No. | Barley cultivars | Pedigree |
|-----|------------------|---|
| 1 | Giza 123 | Giza 117//FAO86 |
| 2 | Giza 124 | Giza 117/Bahteem52// Giza 118/FAO 86 |
| 3 | Giza 125 | Giza117/Bahteem52//Giza118/FAO86 |
| 4 | Giza 126 | Baladi Bahteem/SD729-por12762-Bc |
| 5 | Giza 2000 | Cr366-13-1/Giza121 |
| 6 | Line (LBYT) | (Lignee527/Gerbel/3/Boyb*2/Surb//C 11225.2D/4/M104) |

Data of rust reactions were scored as response and severity of disease infection (%). Disease severity (%) was recorded weekly from the first rust appearance on any test cultivar along with the stage of the through growth season.

Final rust severity (FRS %):

Rust severity (%) of each test cultivar was recorded weekly after the initial infection was occurred, using the modified Cobb's scale (Fig. 2) [15]. Adult plant reaction scored as the percentage of rust severity (%) for each cultivar, at the time when rust was first appeared until the early dough stage [16]. Also, final rust severity (FRS%) was estimated on each cultivar under study as disease severity (%), when rust severity% reached its maximum and finally level in the control plants of the highly susceptible check line LBYT28 [17]. Also, the studied field traits were days to heading, days to maturity, plant height (cm), spike length (cm), number of grains/spike, number of spikes/m², 1000-grain weight (g), biological yield (ton/dan) and grain yield (ardab/feddan). [**feddan = 4200m² and ardab = 120kg**]



150(2): The modified Cobb's scale: A: Actual rust percentage, B: Visual rust severities
151 [15]
152

153 RESULTS


154 Rust fungi are biotrophic pathogens that attack many plant species but are particularly
155 destructive on cereal crops. The stem rust caused by *Puccinia graminis* f. sp. *tritici* has
156 historically caused severe crop losses and continue to threaten production today.

157 Stem rust race (Ug99) poses a serious threat to both wheat and barley worldwide [18].
158 Barley (*Hordeum vulgare* L.) breeders have controlled major stem rust epidemics. Several
159 barley landraces were found to possess a high level of resistance at both the adult plant stages.

160 Molecular identification:

161 ITS amplification:

162 PCR products of approximately 600bp amplified with the ITS-1 F and ITS-4R
163 primers and corresponding to the ribosomal RNA gene were obtained from the tested four
164 isolates (Fig. 3). After purification of PCR products and sequencing, the BLAST-n alignments
165 results showed that the four sequences were associated with high levels of sequence similarity
166 with ribosomal RNA gene sequences for *Puccinia graminis* f. sp. *tritici*. Sequences were
167 deposited in the GenBank database of four accession numbers, i.e. [MW931757](#), [MW931758](#),
168 [MW931759](#) and [MW931760](#) (Badawy 1, Badawy 3, Badawy 6 and Badawy 9, respectively).



170(3): Profile of ribosomal RNA-internal transcribed spacer (rRNA-ITS) amplified by
171 primers ITS1 and ITS4 from the physiologic races. Lanes: M Marker (100 bp DNA), 1, 3, 6 and
172 9.
173

174 Similarity and differences among isolates:


175 Genetic dice similarity showed differences among the races under study and the
species *Puccinia graminis* f. sp. *tritici* in the GenBank database a midst of (99.36 to 98.48 %),
according to the top genetic similarity (99.36 %) between the two races, (isolate Badawy_3
and isolate JQ688990.1, JQ688957.1, and JQ688952.1) as shown in Table (4) and Figure (5).
While the lowest value of genetic similarity (98.48 %) was observed between (isolate
Badawy_1) and (isolates KC853409.1, JQ688984.1, JQ688977.1, JQ688964.1, JQ688954.,
JQ688947.1, JQ688945.1, JQ688934.1, and JQ688902.1) as shown in Table (3) and Figure
(4). 182

Analyzing pathotypes obtained from the two main clusters:

184 The first cluster included isolate Badawy 6, while the second cluster included isolate
Badawy 9, from the individual primer analysis, it seems that isolate Badawy 6 belongs to the
cluster in tested similarly, by a percentage (99.01), (Table 5 and Figure 6), while isolate
Badawy 9 which belongs to the cluster in tested similarly by a percentage (99.19) as shown in
Table (6) and Figure (7).

189 It is therefore more likely these isolates (two members of the cluster) shared the
majority of their genetic materials and will probably meg have originated from similar source.

191 Watched the degree of difference and kinship between the races that are defined in
Figure (8) which shows Phylogenetic tree using (MEGA5) of four *Puccinia graminis tritici*
strains using *ITS rRNA*, showing names of fungus species and accession numbers.




195 **Figure (4): Phylogenetic tree using (MEGA5) of *P. graminis* f. sp. *tritici* race Badawy_1**
 196 **using *ITS rRNA*, showing names of fungus species and accession numbers.**

197
 198
 199

200 **Table (3): *ITS rRNA* of *Puccinia graminis tritici* isolate Badawy1 and related fungus**
 201 **species with the similarity percentage of more than 99%, downloaded from**
 202 **GenBank database.**

| Accession No. | E-value | Query coverage (%) | Similarity (%) |
|---------------|---------|--------------------|----------------|
| JQ688991.1 | 0.0 | 100 | 99.32 |
| KC853409.1 | 0.0 | 100 | 98.48 |
| JQ688984.1 | 0.0 | 100 | 98.48 |
| JQ688977.1 | 0.0 | 100 | 98.48 |
| JQ688964.1 | 0.0 | 100 | 98.48 |
| JQ688954.1 | 0.0 | 100 | 98.48 |
| JQ688947.1 | 0.0 | 100 | 98.48 |
| JQ688945.1 | 0.0 | 100 | 98.48 |
| JQ688934.1 | 0.0 | 100 | 98.48 |
| JQ688902.1 | 0.0 | 100 | 98.48 |


203



205 **Figure (5): Phylogenetic tree using (MEGA5) of *P. graminis* f. sp. *tritici* race Badawy_3**
 206 **using ITS rRNA, showing names of fungus species and accession numbers.**
 207

208 **Table (4): ITS rRNA of *Puccinia graminis tritici* isolate Badawy 3 and related fungus**
 209 **species with the similarity percentage downloaded from GenBank database.**

| Accession No. | E-value | Query coverage (%) | Similarity (%) |
|---------------|---------|--------------------|----------------|
| JQ688990.1 | 0.0 | 100 | 99.36 |
| JQ688943.1 | 0.0 | 100 | 98.70 |
| JQ688957.1 | 0.0 | 100 | 99.36 |
| JQ688973.1 | 0.0 | 100 | 98.70 |
| JQ688956.1 | 0.0 | 100 | 98.70 |
| JQ688969.1 | 0.0 | 100 | 98.54 |
| JQ688952.1 | 0.0 | 100 | 99.36 |
| JX047479.1 | 0.0 | 100 | 98.54 |
| JX424512.1 | 0.0 | 100 | 98.70 |
| JX047482.1 | 0.0 | 100 | 98.70 |



211 **Figure (6): Phylogenetic tree using (MEGA5) of *P. graminis* f. sp. *tritici* race Badawy_6**
 212 **using ITS rRNA, showing names of fungus species and accession numbers.**
 213

214

Table 15): ITS rRNA of *Puccinia graminis tritici* isolate Badawy 6 and related fungus species with the similarity percentage downloaded from GenBank database.

216

| Accession No. | E-value | Query coverage (%) | Similarity (%) |
|---------------|---------|--------------------|----------------|
| KY798400.1 | 0.0 | 100 | 99.01 |
| KY798389.1 | 0.0 | 99 | 99.01 |
| MT965562.1 | 0.0 | 99 | 99.01 |
| MT965559.1 | 0.0 | 99 | 99.01 |
| KC853399.1 | 0.0 | 99 | 99.01 |
| JQ688992.1 | 0.0 | 99 | 99.01 |
| JQ688988.1 | 0.0 | 99 | 99.01 |
| JQ688975.1 | 0.0 | 99 | 99.01 |
| JQ688970.1 | 0.0 | 99 | 99.01 |
| JQ688961.1 | 0.0 | 99 | 99.01 |

217




Figure 7): Phylogenetic tree using (MEGA5) of *P. graminis* f. sp. *tritici* race Badawy 9 using ITS rRNA, showing names of fungus species and accession numbers.

220

221

Table 16): ITS rRNA of *Puccinia graminis tritici* isolate Badawy 9 and related fungus species with the similarity percentage downloaded from GenBank database.

223

| Accession No. | E-value | Query coverage (%) | Similarity (%) |
|---------------|---------|--------------------|----------------|
| JQ688988.1 | 0.0 | 100 | 99.19 |
| JQ688975.1 | 0.0 | 100 | 99.19 |
| JQ688970.1 | 0.0 | 100 | 99.19 |
| JQ688961.1 | 0.0 | 100 | 99.19 |
| JQ688960.1 | 0.0 | 100 | 99.19 |
| JQ688959.1 | 0.0 | 100 | 99.19 |
| JQ688944.1 | 0.0 | 100 | 99.19 |
| JQ688940.1 | 0.0 | 100 | 99.19 |
| JQ688939.1 | 0.0 | 100 | 99.19 |
| JQ688931.1 | 0.0 | 100 | 99.19 |




Figure (8): Phylogenetic tree using (MEGA5) of four *Puccinia graminis tritici* races using *ITS rRNA*, and accession numbers.

226

227

RAPD fingerprinting:


This study at this perspective indicated differences between the *P. graminis* f.sp. *tritici* isolates, which have been collected and studied, using RAPD analysis (Figure 8). These results are similar to those reported by [19] who found in a study on a worldwide basis collections of *P. graminis* f.sp. *tritici* differences for their virulence in addition to molecular backgrounds.

Concurrently, [20] studied a sample of 115 *P. graminis tritici* isolates from the United States and identified six pathotypes and five random amplified polymorphic DNA (RAPD) groups. They found a low correlation coefficient between pathotypes and RAPD groups.

237

238

239



241 **Figure (9): RAPD fingerprinting profiles generated by arbitrarily primed PCR, show**
 242 **results for the four isolates of *Puccinia graminis* f. sp. tritici, isolate 1, isolate**
 243 **3, isolate 6 and isolate 9, respectively.**

244
 245 **Pathogenicity test:**

246 Data presented in Table (7) show that typical symptoms of rust infection, *Puccinia*
 247 *graminis* f. sp. tritici were observed on the susceptible barely line (LBYT) after 14 days from
 248 inoculation and were similar to those previously observed on the collected infected barley
 249 samples. The Egyptian barley varieties, i.e. Giza 123, Giza 124, Giza 125, Giza 126, and Giza
 250 showed high levels of resistance to stem rust by estimate final rust severity (FRS %),
 251 compared to the control line (LBYT, the highly susceptible), where all the cultivars under
 252 study the four locations recorded a (zero) Final rust severity (FRS %), Infection rate of the
 253 disease however, control (line LBYT) recorded the highest rate of infection with stem rust in
 254 Nubya, Sakha, Sids and Giza locations (76.67, 66.22, 63.31 and 60.00 %), respectively,

Table 57): Effect of infection by stem rust on barley cultivars grown under field conditions in four Egyptian locations during 2020/2021 growing season.

256

257

| No. | Barley cultivar | Location | | | |
|-----|---------------------|----------|-------|----------|-------|
| | | Sids | Giza | Nubariya | Sakha |
| 1 | Giza 123 | 0.00 | 0.00 | 0.00 | 0.00 |
| 2 | Giza 124 | 0.00 | 0.00 | 0.00 | 0.00 |
| 3 | Giza 125 | 0.00 | 0.00 | 0.00 | 0.00 |
| 4 | Giza 126 | 0.00 | 0.00 | 0.00 | 0.00 |
| 5 | Giza 2000 | 0.00 | 0.00 | 0.00 | 0.00 |
| 6 | Control (Line LBYT) | 63.31 | 60.00 | 76.67 | 66.22 |




Figure (10): Effect of stem rust on barley cultivars grown in four locations, 2020/2021 growing season.

265

266

Agronomic traits:

268 The results pointed out that the mean squares due to genotypes were significant for all base 269 on average of the four locations (Table 8).

Table (8): Analysis of variance of different agronomic traits for barley genotypes in four locations, 2020/ 2021 growing season.

271

272

| SOV | df | Days to heading | Days to maturity | Plant height | Spike length | Number of grains/spike |
|-----------|----|----------------------------------|-------------------|------------------|--------------|------------------------|
| Rep | 2 | 1.49 | 3.13 | 4.24* | 0.01 | 0.92 |
| Genotypes | 2 | 21.97** | 22.10** | 30.85** | 1.34** | 93.47** |
| Error | 4 | 1.54 | 3.29 | 0.63 | 0.01 | 0.65 |
| SOV | df | Number of spikes/ m ² | 1000-grain weight | Biological yield | Grains yield | Stem rust infection |
| Rep | 2 | 78.76** | 0.95* | 0.01* | 0.08* | 0.003 |
| Genotypes | 2 | 3955.76** | 16.05** | 1.10** | 5.37** | 13.180** |
| Error | 4 | 10.13 | 0.13 | 0.02 | 0.02 | 0.009 |

(*) and (**) significant at 0.05 and 0.01 levels of probability, respectively.

274 Overall mean values for days to heading and for days of maturity showed that the
 275 most desirable mean values towards the earliness were exhibited by the cv. Giza 125 with
 276 average values of (85.85 and 123.22 days), respectively, concerning plant height, cv. Giza
 277 125 plant were the tallest with value (115.22 cm). Meanwhile, cv. Giza 2000 had the highest
 278 mean values for spike length (8.69cm), number of grains/spike (68.68 grain), number of
 279 spikes/m² (513.1 spike), 1000-grain weight (54.34g), biological (7.45 ton) and grain yield
 280 (18.96 ardab) (Table 9).

281 **Table (9): Mean performance estimates of the studied traits for barley genotypes in four**
 282 **locations 2019/ 2020 growing season.**

| Genotypes | Days to heading | Days to maturity | Plant height (cm) | Spike length (cm) | Number of grains/spike | Number of spikes/m ² | 1000-grain weight (g) | Biological yield (ton/ feddan) | Grains yield (ardab/ feddan) |
|------------|-----------------|------------------|-------------------|-------------------|------------------------|---------------------------------|-----------------------|--------------------------------|------------------------------|
| Giza-123 | 86.86 | 126.25 | 109.08 | 8.08 | 66.66 | 440.37 | 52.52 | 6.85 | 17.96 |
| Giza-124 | 87.85 | 127.26 | 112.11 | 8.08 | 66.66 | 436.33 | 53.94 | 7.06 | 18.08 |
| Giza-125 | 85.85 | 123.22 | 115.14 | 7.58 | 63.63 | 484.82 | 50.30 | 6.58 | 17.28 |
| Giza-126 | 87.86 | 125.24 | 112.11 | 7.47 | 60.60 | 492.90 | 50.10 | 6.47 | 16.54 |
| Giza-2000 | 87.87 | 131.30 | 106.05 | 8.69 | 68.68 | 513.10 | 54.34 | 7.45 | 18.96 |
| Line-1 | 87.12 | 127.71 | 113.00 | 6.73 | 53.46 | 424.00 | 48.6 | 5.63 | 14.75 |
| Mean | 87.24 | 126.83 | 111.25 | 7.77 | 63.28 | 465.25 | 51.63 | 6.68 | 17.26 |
| L.S.D-0.05 | 1.84 | 2.70 | 1.17 | 0.15 | 1.19 | 4.73 | 0.53 | 0.08 | 0.20 |
| L.S.D-0.01 | 2.62 | 3.83 | 1.67 | 0.21 | 1.70 | 6.72 | 0.75 | 0.11 | 0.28 |

283

284 DISCUSSION

285 Stem rust of barley and wheat caused by *Puccinia graminis* f. sp. *tritici* Eriks., and E.
 286 Henrici is historically one of the most important plant diseases, stem rust epidemics often result
 287 in major grain losses [21]. All things considered, due to the pathogen's tall potential
 288 for changeability by change and sexual or agamic recombination and
 289 its capacity to duplicate quickly and spread over awesome separations, it remains
 290 a great risk that cannot be disregarded within the brief or long term. This was painfully
 291 outlined in 1999 and 2001 a new race Ug99 or TTKSK with virulence to stem rust disease
 292 tolerant cultivars was detected in Uganda and Kenya, respectively [18], but has since spread
 293 throughout East Africa and is now in the Middle East [22]. Race TTKSK is predicted to

spread to the world's most important wheat and barley growing regions in the near future [23,24] it is the most dangerous threat to cereal crop production in more than 50 yr because it is virulent to most barley and wheat cultivars grown in the major barley and wheat producing areas worldwide [23,25].

Rusts disease are biotrophic fungal pathogens (phylum: Basidiomycota) that cause disease on almost every major family of plants [26]. Puccinia is by distant the biggest class of rust parasites with more than 5000 portrayed species [27,28]. The cereal rusts, especially those attacking the major food crops such as barley and wheat caused famines throughout history and epidemics have been documented in the literature since the time of Aristotle [29].

Barley are hosts to leaf, stem, and stripe rusts disease. The stem rust pathogen *Puccinia graminis* is composed of a number of different *formae speciales* or “special forms” that principally attack one or a few hosts [30,31].

That *Puccinia g. f. sp. tritici* is one of the most important studied fungal plant pathogen systems [32]. After Eriksson sent the concept of the *formae speciales* of *P. graminis* [26], Stakman detailed an indeed better level of specialization: the capacity of *P. graminis f. sp. tritici* confines to particularly assault fair a few wheat genotypes and not others [33]. This concept of physiological specialization in rust organisms driven to the classification of races of *P. graminis f. sp. tritici* and improvement of wheat differentials to distinguish them [33].

Hence the importance of defining the races of the diseased causative agent *Puccinia graminis f. sp. tritici* in this study using molecular biology, where it is more accurate by using (RAPD) genetic fingerprinting and recording the isolates in the gene bank and determining the degree of similarity between the races that were defined under the study that were deposited in the gene bank and other races of the diseased causative *Puccinia graminis f. sp. tritici* present the registered in the gene bank from all countries of the world, where the degree of similarity between them reached (99%).

Induction of rust resistant disease cultivars considered as the primary strategy for combating the rusts [34,35]. Since the 1960s, in wheat incorporating multiple resistance genes

into cultivars has effectively controlled stem rust. On the other hand, barley stem rust disease has been kept in check since the 1940s by breeding varieties with one major durable gene, *Rpg1*; however, other factors such as a largely resistant barley and wheat crops shorter maturation period may contribute to the long-lasting disease control [35].

The Egyptian barley varieties, i.e. Giza 123, Giza 124, Giza 125, Giza 126, and Giza 2003 showed high levels of resistance to stem rust *Puccinia graminis* f. sp. *Tritici*. A similar study on barley leaf rust, confirmed, the ability of barley varieties resistance to leaf rust (*Puccinia hordei*) by [36,37,38], this is the first evaluation of these varieties against stem rust disease and this requires further research on the resistance genes present in these varieties, and their inclusion in the breeding programs for resistance, especially in the transfer of resistance genes found in barley varieties to wheat, whereas, barley, being a true diploid compared to the closely related hexaploid wheat, is the best model in which to study host-pathogen interactions with the expectation that knowledge will lead to effective control measures.

The findings of agronomic traits revealed that mean square due to genotypes of all traits were significant, such results indicated that the tested genotypes varied from each other and behaved differently from season to another [39,40,41,42]. So, the genotypes which exhibited desirable values could be used in breeding program for improving barley production.

341

REFERENCES

1. Kurt and J. Les (2005). Stem rust of small grains and grasses caused by *Puccinia graminis*. Molecular Plant Pathology, 6: 99-111.
2. Wang, A. Wan and X. Chen (2015). Barberry as alternate host is important for *Puccinia graminis* f. sp. *tritici* but not for *Puccinia striiformis* f. sp. *tritici* in the U.S. Pacific Northwest. Plant Disease, 99(11):1507-1516.
3. El-Daoudi, M. Nazim, S. Sherif, I. Shafik and M. Khalifa (1987). Genes conditioning resistance to wheat leaf and stem rust in Egypt. Proc. 5th Congress of the Egypt. Phytopathological Soc., Giza, pp. 387-404.
4. El-Daoudi, O. Mamluk, E. Bekele, Ghanem, H. Enayat, M. Solf and I. Shafik (1994). Preliminary results for leaf and stem rust of wheat, their prevalence and resistance in the Nile Valley countries and Yemen. Fifth Arab Cong. PI. Prot., Fez Morocco, Nov. 23-25 Dec. 2 (Abstr.).

5. **B. K. Mohdly, M. Sallam, F. El-Banoby and O. Boulot (2013)**. First record of stem rust (*Puccinia graminis*) on barley growing in Egypt. *Egypt. J. Phytopathol.*, 41 (1): 219-220.
6. **R. S. Leonard (2001)**. Stem rust—future enemy, in (Stem Rust of Wheat) from Ancient Enemy to Modern Enemy, Peterson, P.D., ed.). St. Paul, MN: APS Press, pp. 119-146.
7. **L. S. Bengham, D. Watters, H. Foulkes and N. Paveley (2008)**. Crop trial and the tolerance of wheat and barley to foliar disease. *Annals of App. Biology*, 54: 159-173.
8. **Watson (1981)**. Wheat and its rust parasites in Australia. In (Wheat Science – Today and Tomorrow) Evans, L.T. and Peacock, W.J., eds). London: Cambridge University Press, :309–147.
9. **R. S. Roelfs (1971)**. Races of *Puccinia graminis* f.sp. *tritici* in the USA during 1970. *Plant Disease Repr.*, 55: 986-990.
10. **D. E. Harder and M. Dunsmore (1990)**. Incidence and virulence of *Puccinia graminis* f.sp. *tritici* on wheat and barley in Canada in 1989. *Can. J. Plant. Pathology*, 12: 424-427.
11. **M. I. Abd El-Hak, N. El-Sherif, I. Shafik, A. A. Bassioni, S. E. Keddis and Y. H. El-Dokki (1982)**. Studies on wheat stem rust virulence and resistance genes in Egypt and neighboring countries, *Egypt. J. Phytopathology*, 14(1-2): 1-10.
12. **R. Nelson, K. Abarenkov, K. Arsson and U. Kõljalg (2011)**. Molecular identification of fungi: Rationale, Philosophical Concerns, and the UNITE Database. *The Open Applied Informatics Journal*; 5(Suppl 1-M9): 81-86.
13. **White, T. Bruns, S. Lee and J. Taylor (1990)**. Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. *PCR Protocols: A Guide to Methods and Applications*, 18: 315–322.
14. **K. M. Brown (1996)**. The choice of molecular marker methods for population genetic studies of plant pathogens. *New Phytol.*, 133 :181-195.
15. **R. Peterson, A. Campbell and A. Hanna (1948)**. A diagrammatic scale for estimating rust intensity on leaves and stems of cereals. *Can. J. Res.*, 26: 496-500.
16. **E. Large (1954)**. Growth stages in cereals. Illustration of the Feekes scale. *Plant Pathology*, 3:128-129.
17. **M. Das, S. Rajaram, W. Ktonstad, C. Mundt and R. Singh (1993)**. Association and genetics of three components of slow rusting in leaf rust of wheat. *Euphytica*, 68: 99-109.
18. **Z. Pretorius, R. Singh, W. Wagoire and T. Payne (2000)**. Detection of virulence to wheat stem rust resistance gene Sr 31 in *Puccinia graminis* f. sp. *tritici* in Uganda. *Plant Disease*, 84: 203.
19. **J. A. Kolmer, J. Liu and M. Sies (1995)**. Virulence and molecular polymorphism in *Puccinia recondita* f.sp. *tritici* in Canada. *Phytopathology*, 85: 276-285.
20. **X. Chen, R. Lin and H. Leung (1993)**. Relationship between virulence variation and DNA polymorphism in *Puccinia striiformis*. *Phytopathology*, 83: 1489-1497.
21. **R. Wanyera, M. G. Kinyua, Y. Jin and R. Singh (2006)**. The spread of stem rust caused by *Puccinia graminis* f. sp. *tritici*, with virulence on Sr31 in wheat in Eastern Africa, *Plant Disease*, 90: p.113.
22. **R. Nazari, M. Mafi, M. Yahyaoui, R. P. Singh and R. F. Park (2009)**. Detection of wheat stem rust (*Puccinia graminis* f. sp. *tritici*) race TTKSK (Ug99) in Iran. *Plant Dis.* 93: 317–318.
23. **R. P. Singh, D. P. Hodson, J. Huerta-Espino, Y. Jin, P. Njau, R. Wanyera, S. A. Herrera-Foessel and R. W. Ward (2008)**. Will stem rust destroy the world's wheat crop? *Agron.* 98: 271– 309.
24. **INMIMYT. (2007)**. Dangerous wheat disease jumps Red Sea: Devastating fungal pathogen spreads from eastern Africa to Yemen, following path scientists predicted. Available at <http://www.seedquest.com/News/releases/2007/january/18117.htm> (verified 8 June 2009). International Maize and Wheat Improvement Center, Mexico D.F., Mexico.
25. **J. Saloff (2005)**. Food for thought: Wheat warning—New rust could spread like wildfire. Available at <http://www.sciencenews.org/view/generic/id/6601/> (verified 4 June 2009). Science News.
26. **C. Agrios (2005)**. *Plant pathology*. 5th ed. Elsevier Academic Press, New York.

27. **Cummins and Y. Hiratsuka (2003)**. Illustrated Genera of Rust Fungi. 3rd ed. Am. Phytopathological Soc., St. Paul, MN.
28. **E.C. Swann, E.M. Frieders and D.J. McLaughlin (2001)**. Urediniomycetes. Springer, New York.
29. **W.R. Bushnell and A.P. Roelfs (1984)**. The cereal rusts. Vol. 1. Origins, specificity, structure, and physiology. Academic Press, Orlando, FL.
30. **D.F. Farr, G. F. Bills, G. P. Chamuris and A.Y. Rossman (1995)**. Fungi on plants and plant products in the United States. APS Press, St. Paul, MN.
31. **B.R. Mohdly, A. E. Khalil and K. A. Amer (2019)**. First record of an isolate (Pathotype) of *Puccinia striiformis* f. Sp. *hordei* the Causative of strip rust on barley in Egypt. Egypt. J. Phytopathol., 47 (1): 367-369.
32. **A.P. Roelfs (1985)**. Wheat and rye stem rust. p. 3– 37. In A.P. Roelfs, and W.R. Bushnell (ed.) The cereal rusts. Vol. 2. Harcourt Brace Jovanovich, New York.
33. **E.C. Stakman and F. J. Piemeisel (1917)**. Biological forms of *Puccinia graminis*. J. Agric. Res. 10: 429– 495.
34. **J.A. Kolmer (2001)**. Early research on the genetics of *Puccinia graminis* and stem rust resistance in wheat in Canada and the United States. pp. 51– 82.
35. **B.L. Steffenson (1992)**. Analysis of durable resistance to stem rust in barley. Euphytica 62: 153– 167.
36. **E.Ghobrial, L. Morsi and T. Sabet (1976)**. Sources of resistance to leaf rust of barley *Puccinia hordei* Oth. in ARE. Proceedings of the 2nd Phytopathol. Conf., 671-682.
37. **E.Chanem, E. Ghobrial, R. Rizk, A. Abdel-shafi, S. Sherif and A. Ageez (1984)**. Sources of resistance to leaf rust of barley (*Puccinia hordei*). Proceedings of EMCIP Symposium, 2: 220-226.
38. **E.Niks, U. Walther, H. Jaiser, F. Martines, D. Rubiales, O. Anderso, K. Flath, P. Gerner, F. Heinrichs, R. Jonsson, L. Kuntze, M. Rasmussen and E. Richter (2000)**. Resistance against barley leaf rust (*Puccinia hordei*) in West-European spring barley germplasm. Agronomie, 20: 769-782.
39. **H.Hartleb, U. Meyer and C. Lehmann (1990)**. Resistance behaviour of common barleys to different isolates of *Drechslera teres* (Sacc.) Shoem. Archiv für Phytopathologie und Pflanzenschutz, 26(3): 257-264.
40. **K.Zaki and A. Al-Masry (2008)**. Detection of biochemical genetic markers for net blotch disease resistance and barley grain yield. Egyptian Journal of Phytopathology, 36: 1442.
41. **Amal Mahmoud Hassan Abdel-Haleem and Mohamed Mansour Abdel-Aty (2021)**. The Relationship Between Varieties and Acrylamide Formation in Roasted Barley. Egypt. J. Chem. Vol. 64, No. 9, pp. 5357 - 5372 (2021).
42. **M.Mansour and Aziza A. Aboulila (2021)**. Molecular variability and salinity effects on growth characters and antioxidant enzymes activity in Egyptian barley genotypes. Physiological and Molecular Plant Pathology. 116, 10173.