

# Characterization, extraction of biochar and humic acid on yield parameters of maize in maize-chickpea cropping system and its residual effect on chickpea under rainfed condition

## Abstract

A field experiment was conducted during *Kharif* 2019 and 2020 in a at ICAR-Krishi Vigyana Kendra, Kalaburagi-II (Raddewadagi), Jewargi taluk, Kalaburagi district, to study the characterization, extraction of biochar and humic acid on yield parameters of maize in maize-chickpea cropping system and its residual effect on chickpea under rainfed condition .The results showed that, the biochar was highly alkaline in nature (pH of 8.91), medium in salt content with an electrical conductivity of 1.89 dS m<sup>-1</sup>. The total carbon content (78.20 g kg<sup>-1</sup>), nitrogen, phosphorus and potassium (0.64, 0.29 and 1.02 %, respectively). It also showed good amount of secondary and micronutrients. Biochar had also lower the bulk density (0.54 Mg m<sup>-3</sup>) and good water holding capacity (67.50 %). Whereas, the humic acid showed acidic in reaction (pH 4.20) and higher the nitrogen content (1.38 %). The maize was the test crop to study the direct effect and chickpea was raised to study the residual effect. The experiment were laid out in Randomized Complete Block Design with fifteen treatments and replicated thrice. The pooled analysis revealed that significantly higher significantly higher yield and yield parameters of maize and succeeding chickpea crops were recorded with application of 100 % NPK + biochar @ 5 t ha<sup>-1</sup> + HA @ 10 kg ha<sup>-1</sup> and it was on par with the treatment receiving 100 % NPK + biochar @ 2.5 t ha<sup>-1</sup> + HA @ 10 kg ha<sup>-1</sup> compare to other treatments.

**Key word:** Biochar, Humic acid, cropping system, characterization, extraction and yield

## INTRODUCTION

Globally, maize is known as queen of cereal because of the highest genetic yield potential among the cereals. It is cultivated on nearly 150 m ha in about 160 countries under diverse soil, climate, biodiversity and management practices contributing 36 per cent (782 m t) of the global grain production. The United States of America (USA) is the largest producer of maize and contributes nearly 35 per cent of the total production in the world with the

highest productivity ( $> 9.6 \text{ t ha}^{-1}$ ) which is double than the global average ( $4.92 \text{ t ha}^{-1}$ ). Whereas, in India maize is grown over an area of 9.50 million hectares with a production of 23.29 million tonnes and an average productivity of  $2.45 \text{ t ha}^{-1}$  which is much lower than most of the maize growing countries of the world. “In India, maize is grown in Karnataka, Rajasthan, Madhya Pradesh and Maharashtra. In Karnataka, it occupies an area of 1.38 million hectares with production of 3.98 million tonnes with an average productivity of  $2.88 \text{ t ha}^{-1}$ ” (Anon., 2014).

Maintaining soil organic matter is a prerequisite to ensure soil health and crop productivity in rainfed farming. India ranks first among the countries that practice rainfed agriculture. Out of the estimated 141 Mha net cultivated land in India, about 75 Mha is rainfed, spread over 177 districts of the country and produces about 40% of the total food grain in India. Efficient use of crop residue based amendment in soil is an important strategy to improve the soil fertility and productivity in rainfed areas. Annually 500 Mt crop residues are generated in India, out of which 141 Mt is surplus. Among different crops, oilseeds (29 Mt), pulses (13 Mt) and cotton (53 Mt) generate maximum residues in India, which are advertently niche crops for rainfed areas. The surplus crop residues of castor, cotton and pigeonpea stalk are estimated to be 18.0, 11.8 and 9.0 Mt, respectively. These residues are either partially utilized or unutilized due to various constraints. Surplus and unused crop residues when left unattended, often disrupt land preparation, crop establishment and early crop growth, and therefore are typically burnt on farm which causes environmental problems and substantial nutrient losses. For more effective management and disposal of the crop residues, their conversion into biochar through thermo-chemical process (slow pyrolysis) is gaining importance as a novel and economically alternative way of managing unusable and excess crop residues. Much of the stimulus for this interest has come from research on the soils of the Amazon basin, known as Terra Preta de Indio, that contain variable quantities of organic black carbon considered to be of anthropogenic origin. Conversion of crop and on-site agroforestry residues to biochar and its soil application as an amendment can turn the hitherto excess residues available in India into a useful material for enhancing soil health and crop productivity.

Varied technological methods of pyrolysis can convert feed stocks like grasses, nutshells, forestry products and animal manures to produce biochar. Pyrolysis is the heating of biomass in an oxygen limited atmosphere, causing release of volatile C structures, hydrogen (H<sub>2</sub>), methane (CH<sub>4</sub>) and carbon monoxide (CO). The volatile C structures

(alcohols, oils, tars, acids, etc.) can be re-condensed as bio-oil. The biochar that remains consists mainly of C, and contains O, H, N, and ash (calcium and potassium). Biochars with large amounts of carbon in poly-condensed aromatic structures are obtained by pyrolyzing organic feed stocks at high temperatures (400 to 700 °C), which has fewer ion exchange functional groups due to dehydration and decarboxylation, potentially limiting its usefulness in retaining soil nutrients. On the other hand, biochar produced at lower temperatures (250 to 400 °C) have higher yield recoveries and contain more C=O and CH functional groups that can serve as nutrient exchange sites after oxidation. Moreover, biochar produced at these lower pyrolysis temperatures have more diversified organic character, including aliphatic and cellulose type structures. These may be good substrates for mineralization by bacteria and fungi, which have an integral role in nutrient turnover processes and aggregate formation. Feedstock selection also has a significant influence on biochar surface properties and its elemental composition.

Biochar is a fine grained, highly porous charcoal substance that is distinguished from other charcoals in its intended use as a soil amendment. The particular heat treatment of organic biomass used to produce biochar contributes to its large surface area and its characteristic ability to persist in soils with very little biological decay (Lehmann *et al.* 2006). While raw organic materials supply nutrients to plants and soil microorganisms, biochar serves as a catalyst that enhances plant uptake of nutrients and water. Compared to other soil amendments, the high surface area and porosity of biochar enable it to adsorb or retain nutrients and retain water and also provide a habitat for beneficial microorganisms to flourish (Glaser *et al.* 2002, Lehmann *et al.* 2006 and Warnock *et al.*, 2007). However, since biochar possesses recalcitrant carbon, it places a doubt that biochar application may pose risk for soil microbes. Moreover, the pH of majority of biochar is alkaline in nature which may alter the microbially mediated nutrient transformations. The possible connections between biochar properties and the soil biota and their implications for soil processes have not yet been systematically described; hence the driving processes are still poorly identified.

“Biochar provides a suitable habitat for a large and diverse group of soil microorganisms. A higher retention of microorganisms in biochar amended soils may be responsible for greater activity and diversity due to a high surface area as well as surface hydrophobicity of both the microorganisms and biochar. A strong affinity of microbes to biochar can be expected since the adhesion of microorganisms to solids increases with higher

hydrophobicity of the surfaces. Biochar is an effective to activate living things and improve natural environment. Carbonized biomass such as rice husk charcoal or wood ash have been valuable material as soil amendment. The optimal biochar combining fertilizer and carbon storage function in soils would activate the microbial community leading to nutrient release and fertilization and would add to the decadal soil carbon pool” (Venkatesh *et al.*, 2018). “Biochar’s inherent physical quality contributes to the improvement in the soil porosity” (Lammirato *et al.*, 2011).

“Humic substances are major components of organic matter, have both direct and indirect effects on plant growth” (Sangeetha *et al.*, 2006). Humic acid (HA) improves the physical chemical and biological properties of the soil and influences plant growth. Because of its molecular structure, it provides numerous benefits to crop production.

## **MATERIAL AND METHODS**

Field investigations carried out during 2019-20 and 2020-21 to study the characterization, extraction of biochar and humic acid on yield parameters of maize in maize-chickpea cropping system and its residual effect on chickpea under rainfed condition. The soil of the experimental site was clay in texture which was alkaline in nature (pH 8.01), with low salt content ( $0.29 \text{ dSm}^{-1}$ ) and low organic carbon content ( $4.80 \text{ g ka}^{-1}$ ). The available nitrogen ( $230.50 \text{ kg ha}^{-1}$ ) was low, available phosphorus ( $31.60 \text{ kg ha}^{-1}$ ) was medium, available potassium content ( $385.63 \text{ kg ha}^{-1}$ ) was high. The experiments were laid out in Randomized Complete Block Design (RCBD) with fifteen treatments and replicated thrice. The two experiments were conducted during *kharif* 2019 and 2020 with maize main crop and the residual effect of the said treatments were studied during *rabi* seasons of 2019 and 2020 with chickpea as succeeding crop.

Recommended dose of fertilizer for maize crop is 100: 50: 25 kg N: P<sub>2</sub>O<sub>5</sub>: K<sub>2</sub>O kg ha<sup>-1</sup>. The nitrogen was given in three equal splits *viz.*, 1/3<sup>rd</sup> as basal, 1/3<sup>rd</sup> at knee high stage (30 DAS) and the remaining 1/3<sup>rd</sup> at tasseling and silking stage (60 DAS). Entire dose of phosphorous and potassium in the form of diammonium phosphate (DAP) and murate of potash (MOP) respectively were applied as basal dose to the crop.

### **Production of biochar**

Representative biochar sample was collected from Munadaragi taluk, Gadag district. The biochar was prepared at 600-700 °C and the feedstock used for preparation of biochar was *Prosopis sp.* and it is called as *Prosopis juliflora* biochar. The biochar sample was analysed for different parameter as per the methods given in Table 1.

### **Extraction of humic substance**

Twenty gram of air dried organic sample was weighed in to 250 ml conical flask 200 ml of 0.1 N NaOH was added (Schnitzer and Skinner, 1968) and shaken for 24 hours. The dark colored supernatant solution was separated by centrifugation and collected. The extraction procedure was repeated thrice using 50 ml of extractant each time for complete extraction of the humic substances.

### **Fractionation and purification of humic substances**

The precipitated humic acid fraction was separated by centrifugation. Precipitation and centrifugation was repeated to attain partial purification of humic acid fraction as described by Stevenson (1981). The fractions were further purified by treating with HCl - HF mixture (5 ml of each HCl and HF acids were dissolved in 990 ml of double distilled water) for 24 hours and this acid mixture was separated by centrifugation. The residue so obtained was thoroughly washed with distilled water and freeze dried to obtain humic acid.

### **Digestion of humic acid sample for determination of elements**

A known weight of humic acid sample was taken in a 250 ml conical flask and was pre-digested by adding 10 ml of HNO<sub>3</sub> and keeping it overnight. Diacid mixture (10 ml) in 9:4 proportion (HNO<sub>3</sub>:HClO<sub>4</sub>) was added and heated on sand bath until a snow white residue was obtained. The residue was cooled and diluted to a known volume with distilled water, filtered and made up to 100 ml using distilled water it was further used for estimation of elements. The methods adopted are same as outlined in Table .1.

### **Statistical analysis of data**

The experimental data obtained was subjected to statistical analysis using Fisher's method of analysis of variance as out lined by Gomez and Gomez (1984). The level of significance used in 'F' and 't' tests was p= 0.05. Critical difference values were calculated,

wherever 'F' test was found significant. Results have been interpreted and discussed based on the pooled data of two years (2019 and 2020).

## RESULTS AND DISCUSSION

The results obtained from the present investigation as well as relevant discussion have been summarized under following heads:

### Characterization of biochar

Representative biochar sample collected from Mundaragi taluk, Gadag district was analysed for different parameters (Table 2).

The data revealed that, the biochar recorded the maximum water holding capacity of 67.50 per cent and bulk density of  $0.42 \text{ Mg m}^{-3}$ . The chemical composition of biochar was found to be alkaline in nature with a pH of 8.91 and the electrical conductivity of  $1.89 \text{ dS m}^{-1}$ . The total carbon content of  $78.20 \text{ g kg}^{-1}$  was recorded; the total nitrogen, phosphorus and potassium were 0.64, 0.29 and 1.02 per cent, respectively. It also showed good amount of total calcium, magnesium and sulphur with consists of 0.090, 0.061 and 0.38 per cent, respectively. Biochar also recorded appreciable quantities of total iron, manganese, zinc and copper to an extent of 1.08, 0.043, 0.033 and 0.021 ppm, respectively.

The powdered biochar was characterized for different physical (BD and WHC) and chemical properties. The data showed (Table 2) that, the *Prosopis* biochar had lower bulk density of  $0.54 \text{ Mg m}^{-3}$ . Similar results were reported by Brewer *et al.* (2014) who estimated that the bulk density of biochar produced from different feedstock ranged from 0.25 to  $0.60 \text{ gcm}^{-3}$ . Biochar derived from *Prosopis juliflora* wood recorded high water holding capacity (67.5 %). This might be due to higher total carbon and surface area. The chemical analysis suggested that biochar found to be alkaline in nature with pH 8.91, medium in salt content with an electrical conductivity of  $1.89 \text{ dS m}^{-1}$ . It might be due to presence of ash residue that is dominated by carbonates of alkali and alkaline earth metals. Similar results were observed by Rumi, (2015), Wan *et al.* (2014) and Rondon *et al.* (2007) who reported that wood biochar was found to have higher saline salt content, followed by coir waste biochar. The total carbon, nitrogen, phosphorus and potassium were found to be higher with 78.20, 0.64, 0.29 and 1.02 per cent, respectively. The higher nitrogen and phosphorus content observed in the biochar which might be attributed to the optimum heat maintained during pyrolysis. It also

recorded good amount of calcium, magnesium and sulphur and the values were 0.09, 0.06 and 0.38 per cent, respectively (Lehman *et al.*, 2009). The content of iron, manganese, zinc and copper found to be in appreciable amounts in biochar. Variability in physical and chemical properties of biochar depends on the material used to produce biochar, the oxygen supply and the temperature achieved during pyrolysis. Similar results were reported by Wang *et al.* (2013) and Lua and Yang, (2004).

### **Characterization of humic acid**

The data on the characteristics of humic acid extracted from vermicompost are presented in Table 3.

The data revealed that the humic acid extracted from vermicompost was acidic in reaction (pH 4.20). The concentration of nitrogen, phosphorus, potassium, calcium, magnesium and sulphur were 1.38, 0.23, 0.56, 0.06, 0.04 and 0.81 per cent, respectively. The average concentration of humic acid content, ash content, iron, manganese, copper and zinc and were 4.98, 2.51, 3251.5, 515.31, 108.20 and 250.5 per cent respectively. It also contained some quantity of elemental composition of total carbon and hydrogen were 43.51 and 4.96 per cent. Similar findings were reported by Sathisha and Devarajan (2011) and they reported higher organic carbon in the humic acid extracted from vermicompost which may be due to higher nutrient contents of the test materials in the present study. Similar findings were also observed with respect to nutrient concentration in different organic sources (Jindo *et al.*, 2012). Atiyeh *et al.* (2002) and muter *et al.* (2015). Paola *et al.* (2006) stated that humic acid derived from municipal solid waste compost had C, H, N and ash content (52.4, 5, 4.7 and 1.21 % respectively). Similar results were presented by Kasongo *et al.* (2011) who reported that, the nutrient content of coffee pulp acts as an alternative to fertilizer with beneficial effect in improving soil properties and supply of nutrients for adequate growth and development of plants.

### **Grain and stover yield (q ha<sup>-1</sup>) of maize**

The data pertaining to grain and straw yield of maize crop as influenced by biochar and humic acid application during both the years as well as pooled data analysis are presented in Table 4.

Significantly higher grain yield ( $82.91 \text{ q ha}^{-1}$ ) was recorded with application of 100 % NPK + biochar @  $5.0 \text{ t ha}^{-1}$  + HA @  $10 \text{ kg ha}^{-1}$  ( $T_{13}$ ) as compared to other treatments. However, it was on par with  $T_{12}$  which received 100 % NPK + biochar @  $2.5 \text{ t ha}^{-1}$  + HA @  $10 \text{ kg ha}^{-1}$  ( $79.85 \text{ q ha}^{-1}$ ) followed by  $T_3$  with application of 100 % NPK + FYM @  $7.5 \text{ t ha}^{-1}$  + HA @  $10 \text{ kg ha}^{-1}$  ( $79.48 \text{ q ha}^{-1}$ ). While, significantly the lowest grain yield of  $31.44 \text{ q ha}^{-1}$  was recorded in absolute control treatment as compared to other treatments.

Similarly, significantly higher stover yield ( $105.76 \text{ q ha}^{-1}$ ) was recorded in  $T_{13}$  treatment which received 100 % NPK + biochar @  $5.0 \text{ t ha}^{-1}$  + HA @  $10 \text{ kg ha}^{-1}$  compared to other treatments. However, it was on par with  $T_{12}$  supplied with 100 % NPK + biochar @  $2.5 \text{ t ha}^{-1}$  + HA @  $10 \text{ kg ha}^{-1}$  ( $100.33 \text{ q ha}^{-1}$ ). Whereas, the lowest stover yield of  $39.09 \text{ q ha}^{-1}$  was recorded in absolute control. Similar trend was observed during individual years and as well as pooled data. Higher grain and stover yield in maize could be attributed to better uptake of essential nutrients and its translocation to economic parts as well as improvement in yield attributing characters like cob weight, cob length and number of grain per corn.

Economic yield is expressed as a function of factor that contributed to yield which is known as yield attributes. Increase in yield might be attributed to improvement in yield components due to better partitioning of carbohydrates from leaf to reproductive parts resulting in increased yield. Addition of more nutrients through biochar, humic acid and inorganic fertilizers resulted in higher grain and stover yield. Many research works have reported that biochar and humic acid enhance the yield of different crops *viz.*, sugarcane, rice and maize tomato, chickpea etc. These results supported by Asai *et al.* (2009), Chen *et al.* (2010) and Ogawa and Okimori (2010) and Madhavi (2014) reported that “application of recommended dose of NPK along with biochar at  $7.5 \text{ t ha}^{-1}$  and humic acid at  $30 \text{ kg ha}^{-1}$  were significant in increasing seed yield. Though in the absence of biochar, 75 % NPK put forth significantly lower yield than 100 % NPK. Integration with biochar at the highest level of  $7.5 \text{ t ha}^{-1}$  and humic acid at  $30 \text{ kg ha}^{-1}$ , the yields from the two levels of fertilizer (75 and also 100 % NPK) were at a par”.

### **Seed and haulm yield ( $\text{q ha}^{-1}$ ) of residual chickpea crop**

The data showed similar trend with respect to seed and haulm yield. Seed and haulm yields of chickpea were significantly influenced by the residual effect of biochar and humic acid during individual years as well as pooled data are presented in Table 5.

The perusal of pooled data indicated that, the residual effect of 100 % NPK + biochar @ 5.0 t ha<sup>-1</sup> + HA @ 10 kg ha<sup>-1</sup> (T<sub>13</sub>) treatment showed significantly higher seed and haulm yield (16.66 and 20.75 q ha<sup>-1</sup>, respectively) compared to other treatments. However, which was on par with treatment (T<sub>12</sub>) which received 100 % NPK + biochar @ 2.5 t ha<sup>-1</sup> + HA @ 10 kg ha<sup>-1</sup> (13.82 and 18.67 q ha<sup>-1</sup>, respectively). Whereas, significantly lower seed and haulm yields was obtained in absolute control (7.08 and 10.38 q ha<sup>-1</sup>, respectively). Similar trend was observed during both the years.

The yield data indicated a significant influence of residual biochar and humic acid on chickpea. This could be mainly due to attributed by nutrients present in the soil and in the biochar and humic acid, which were made available to the crop. Balanced supply of nutrients might have enabled chickpea crop to produce higher pod bearing branches and higher number of pods per plant. Yooyen *et al.* (2015) reported that “addition of biochar to soybean increased the number of seeds per plant”. Mirl *et al.* (2013) reported that “an increase in seed yield and haulm yield was observed in the biochar treatment as compared to no biochar treatment. Soil treated with biochar and humic acid increased the stover yield of crop. The better performance of chickpea crop might be due to the presence of bacterial activity which helps in increasing the phosphorus availability in soil, thereby improving plant growth and yield”. Viruel *et al.* (2014) and Bandara *et al.* (2015) they reported that “the increased the crop growth and yield by combined application of biochar and FYM. The timely availability of nitrogen with the interaction effect of biochar and humic acid mineralization might be responsible for the production of highest seed and stover yield. The superiority of *Prosopis sp.* biochar applied treatments over RDF alone was due to biochars was being protected from further decomposition thereby it could supply longer and steady source of macro and micronutrients. However, in present study residual biochar and humic acid applied in combination recorded higher yield and yield attributing parameters in the treatment T<sub>13</sub> (100 % NPK + biochar @ 5 t ha<sup>-1</sup> 8 t ha<sup>-1</sup> + HA @ 10 kg ha<sup>-1</sup>)”.

## CONCLUSION

Use of biochar in agricultural systems is one viable option that can enhance natural rates of carbon sequestration in the soil, reduce farm waste and improve the soil quality and crop productivity. Further, several studies across the world have established that biochar application increases conventional agricultural productivity and mitigate GHG emissions from agricultural soils.

Prosopis biochar was found to be a rich source of carbon and nutrients with alkaline pH, medium EC, lower density and higher maximum holding capacity. The prosopis biochar and humic acid is a good source of microorganisms and hastens the better mineralization rate and increased the efficiency of biochar in soil. Study indicates the necessity of combined application of biochar, humic acid and fertilizer for higher growth, yield and quality parameters of maize and residual chickpea crop as compared to alone application of biochar, FYM and fertilizers.

## REFERENCES

- Albuquerque JA, Salazar P, Barron V, Torrent J, Campillo MC, Gallardo, A and Villar R 2013. Enhanced wheat yield by biochar addition under different mineral fertilization levels. *Journal Agronomy for Sustainable Development* **33**: 475-484.
- Anonymous 2014. Area, production and productivity of major pulses in India. Agricultural Statistics at a Glance. *Ministry of Agriculture government of India*.
- Asai H, Samson BK, Stephan HM, Songy K, Homma K, Kiyono Y, Inoue Y, Shiraiwa T and Horie T 2009. Biochar amendment techniques for upland rice production in northern Laos. *Field Crops Research* **111**: 81-84.
- Atiyeh RM, Lee S, Edwards CA, Arancon NQ and Metzger JD 2002. The influence of humic acids derived from earthworm-processed organic wastes on plant growth. *Bio resource and Technology* **84**: 7-14.
- Bandara T, Herath I, Prasanna K, Mihiri S, Gamini S, Nishanta R, Meththika V and Yong SO 2015. Role of woody biochar and fungal bacterial co-inoculation on enzyme activity and metal immobilization in serpentine soil. *Journal of Soils and Sediments* **10**(7): 43-68.
- Chen Y, Shinogi Y and Taira M 2010. Influence of biochar use on sugarcane growth, soil parameters and ground water quality. *Australian Journal of Soil Research* **48**: 526-530.
- Glaser B, Lehmann, J and Zech W 2002. Ameliorating physical and chemical properties of highly weathered soils in the tropics with charcoal. *Biology and Fertility of Soils* **35**: 219-230.

- Gomez KA and Gomez, AA 1984. *Statistical Procedures for Agricultural Research*, 2nd Edition, A Wiley Inter Science Publications, New York, USA.
- Hesse PR 1994. *A textbook of soil chemical analysis*. Hans publication pp:111
- Jackson ML 1973. *Soil Chemical Analysis*. Prentice Hall of India Pvt. Ltd., New Delhi, pp. 498.
- Jindo K, Silvia AM, Elena CN, Pérez AF, Teresa H, Carlos G, Natália OA and Luciano P C 2012. Root growth promotion by humic acids from composted and non-composted urban organic wastes. *Plant and Soil* **353**: 209-220.
- Kasongo RK, Verdoodt A, Kanyankagote P, Baert G and Van ranst E 2011. Coffee waste as an alternative fertilizer with soil improving properties for sandy soils in humid tropical environments. *Soil Use and Management* **27**: 94-102.
- Lammirato C, Miltner A and Kaestner M 2011. Effects of wood char and activated carbon on the hydrolysis of cellobiose by b-glucosidase from *Aspergillus niger*. *Soil Biol. Biochem* **43**: 1936-1942
- Lehmann J, Czimnik C, Laird D and Sohi S 2009. Stability of biochar in the Soil. In: Biochar for Environmental Management: *Science and Technology* **12**: 176-183.
- Lehmann J, Gaunt J and Rondon M 2006. Bio-char sequestration in terrestrial ecosystems: A review. *Mitigation and Adaptation Strategies for Global Change* **11**: 403-427.
- Lua AC and Yang T 2004. Effects of vacuum pyrolysis conditions on the characteristics of activated carbons derived from pistachio-nut shells. *Journal of Colloid and Interface Science* **276**: 364-372.
- Madhavi P 2014. Effect of biochar and humic acid on fertilizer use and yield of maize (*Zea mays* L.) in alfisols of southern telangana region of andhrapradesh. *M. Sc. (Agri) Thesis, Acharya N. G. Ranga Univ. Agric. Sci., Hydrabada*.
- Mirl AH, Lal SB, Salmani M, Abid M and Khan I 2013. Growth, yield and nutrient content of blackgram (*Vigna mungo*) as influenced by levels of phosphorus, sulphur and phosphorus solubilizing bacteria. *Journals of Agriculture* **11**(1): 1-6.

- Muter O, Limane B, Strikauska S and Klavins M 2015. Effect of humic-rich peat extract on plant growth and microbial activity in contaminated soil Material. *Science and Applied Chemistry* **32**: 64-74.
- Ogawa M and Okimori Y 2010. Pioneering works in biochar research, Japan. *Soil Research* **48**: 489-500.
- Orlov DS, Rozanova ON, Matykhino RD 1992. Elemental and phase composition of soil. In: Soil Chemistry (Ed. Orlov. D.S.), Oxoman Press Pvt. Ltd., New Delhi, 29-56.
- Page AL, Miller RH and Kenay DR 1982. *Methods of Soil Analysis, part-II*, Soil Science Society of America, Inc, Publishers, Madison, Wisconsin, USA.
- Paola A, Campitelli Manuel I, Velasco Silvia B and Ceppi 2006. Chemical and physico-chemical characteristics of humic acids extracted from compost, soil and amended soil. *Talanta* **69**: 1234-1239.
- Piper CS 1966. *Soil and Plant Analysis*. Academic Press, New York, pp. 367.
- Rondon MA, Lehmann J, Ramirez J and Hurtado M 2007. Biological nitrogen fixation by common beans increases with biochar additions. *Biology and Fertility of Soils* **43**(6): 699-708.
- Rumi N, Neonjyoti Rahul SC, Bikram B, Nirmali G, Ajitabh B and Rupam K 2015. Biochar: An overview on its production, properties and potential benefits. *Biology and Biotechnology for Sustainable Development* 13-40.
- Sangeetha M, Singaram P and Devi RD 2006. Effect of lignite humic acid and fertilizers on the yield of onion and nutrient availability. *18<sup>th</sup> world cong. soil science* USA.
- Satisha GC and Devarajan L 2011. Composition and characterization of humic substances extracted from effluent-based pressmud composts. *Agropedology* 21(I): 8-17.
- Schnitzer M and Skinner SI 1968. Alkali versus acid extraction of soil organic matter, *Soil Science* **105**: 392-396.
- Stevenson FJ 1981. *Humus chemistry: Genesis, Composition and Reactions*. **110**: 496-502.

- Stevenson FJ 1994, *Humus Chemistry: Genesis, Composition and Reactions*, 2nd Edn. John Wiley and Sons, New York, pp: 378-486.
- Venkatesh G, Sammi Reddy K, Gopinath KA, Visha Kumari V and Chary RG 2018. Biochar for enhancing the soil, water and crop productivity in rainfed areas. *Rainwater Management for Climate Resilient Agriculture in Drylands* : pp 523
- Viruell E, Erazzu LE, Martinez Calsina<sup>1</sup> L, Ferrero MA, Lucca ME and Sineriz F 2014. Inoculation of maize with phosphate solubilizing bacteria effect on plant growth and yield. *Journal of Soil Science and Plant Nutrition* **14**(4): 819-831.
- Wang Y, Yuting H, Xu Z, Shenqiang W and Guangxi X 2013. Comparisons of biochar properties from wood material and crop residues at different temperatures and residence time. *Energy and Fuels* **27**(10): 5890-5899.
- Wan Q, Yuan JH, Xu KR and Li XH 2014. Pyrolysis temperature influences ameliorating effects of biochar on acidic soil. *Environmental Science and Pollution Research* **21**: 2486-2495.
- Warnock DD, Lehmann J, Kuyper TW and Rillig MC 2007. Mycorrhizal responses to biochar in soil - concepts and mechanisms. *Plant and Soils* **300**: 9-20.
- Yooyen J, Wijitkosum S and Sriburi T 2015 Increasing yield of soybean by adding biochar. *J. Environmental Research and Development* **9**(4): 1066-1074.

**Table. 1: Methods followed for analysis of biochar and humic acid**

<b>A. Biochar</b>			
<b>Sl. No.</b>	<b>Parameter</b>	<b>Method</b>	<b>Reference</b>
1	pH	Potentiometry	Jackson (1973)
2	EC(dS m <sup>-1</sup> )	Conductometry	Jackson (1973)
3	MWHC (%)	Keen's cup method	Piper (1966)
4	Bulk density	Core method	Barnyard and Henry, (1921)
5	Particle density	International pipette method	Piper (1966)
6	Porosity		
7	Total carbon (%)	Dry combustion method	Page <i>et al.</i> (1982)
8	Total nitrogen (%)	Kjeldahl digestion distillation method	Jackson (1973)
9	Total phosphorus (%)	Diacid digestion and vanadomolybdate method	Jackson (1973)
10	Total potassium (%)	Diacid digestion and flame photometer method	Jackson (1973)
11	Total calcium and magnesium (%)	Complexometric titration method	Jackson (1973)
12	Total Sulphur (ppm)	0.15% CaCl <sub>2</sub> extraction and turbidimetry	Jackson (1973)
13	DTPA extractable micronutrients like Zn, Fe, Cu & Mn (ppm)	Diacid digestion and atomic absorption spectrophotometry	Jackson (1973)
<b>B. Humic acid</b>			
1	pH	Potentiometry	Jackson (1973)
2	Total acidity (me g <sup>-1</sup> )	Ba(OH) <sub>2</sub> method	Schnitzer and Gupta (1965)
3	Elemental analysis: carbon, hydrogen, nitrogen and sulphur	dry combustion method using CHNS analyser	Orlov <i>et al.</i> (1992)
4	Total Fe, Mn, Zn and Cu (ppm)	Diacid digestion and Atomic Absorption Spectrophotometry	Jackson (1973)
5	HA Concentration (%)	Stevenson method	Stevenson (1994)
6	Ash content (%)	Hesse method	Hesse (1994)

**Table 2. Characterization of *Prosopis* biochar**

Sl. No	Parameters	Value
<b>Physical properties</b>		
1.	Bulk density (Mg m <sup>-3</sup> )	0.42
2.	MWHC (%)	67.50
<b>Chemical properties</b>		
3.	pH (1:10)	8.91
4.	EC(dS m <sup>-1</sup> )	1.89
5.	Total carbon (%)	78.20
6.	Total Nitrogen (%)	0.64
7.	Total Phosphorus (%)	0.29
8.	Total Potassium (%)	1.02
9.	Total Sulphur (%)	0.38
10.	Total Calcium (%)	0.090
11.	Total Magnesium (%)	0.061
12.	Total Zn (ppm)	0.033
13.	Total Fe (ppm)	1.08
13.	Total Cu (ppm)	0.021
14.	Total Mn (ppm)	0.043

**Table 3. Characterization of humic acid extracted from vermicompost**

<b>Sl. No</b>	<b>Parameters</b>	<b>Values</b>
1	Humic acid content (%)	4.98
2	Ash content (%)	2.51
3	Total acidity (me g <sup>-1</sup> )	5.17
4	pH (1:2.5)	4.20
5	Total carbon (%)	43.51
6	Total Hydrogen (%)	4.96
7	Total Nitrogen (%)	1.38
8	Total Phosphorus (%)	0.23
9	Total Potassium (%)	0.56
10	Total Sulphur (%)	0.81
11	Total Calcium (%)	0.06
12	Total Magnesium (%)	0.04
13	Total Zn (ppm)	250.5
14	Total Fe (ppm)	3251.5
15	Total Cu (ppm)	108.20
16	Total Mn (ppm)	515.3

**Table 4: Grain and stover of maize as influenced by application of biochar and humic acid in maize - chickpea cropping system under rainfed condition**

Treatments	Grain yield (q ha <sup>-1</sup> )			Stover yield (q ha <sup>-1</sup> )		
	2019	2020	Pooled	2019	2020	Pooled
<b>T<sub>1</sub></b> : Absolute control	32.75	30.12	31.44	40.83	37.34	39.09
<b>T<sub>2</sub></b> : 100 % NPK + FYM @ 7.5 t ha <sup>-1</sup>	55.49	58.75	57.12	60.49	67.27	63.88
<b>T<sub>3</sub></b> : T <sub>2</sub> + HA @ 10.0 kg ha <sup>-1</sup>	77.35	81.61	79.48	96.14	102.51	99.33
<b>T<sub>4</sub></b> : 75 % NPK + Biochar @ 2.5 t ha <sup>-1</sup>	49.31	52.21	50.76	57.35	62.12	59.74
<b>T<sub>5</sub></b> : 75 % NPK + Biochar @ 5.0 t ha <sup>-1</sup>	51.65	54.84	53.25	59.70	64.50	62.10
<b>T<sub>6</sub></b> : 75 % NPK + Biochar @ 2.5 t ha <sup>-1</sup> + HA @ 10 kg ha <sup>-1</sup>	67.76	71.66	69.71	75.17	81.60	78.39
<b>T<sub>7</sub></b> : 75 % NPK + Biochar @ 5.0 t ha <sup>-1</sup> + HA @ 10 kg ha <sup>-1</sup>	70.27	74.46	72.37	80.78	88.64	84.71
<b>T<sub>8</sub></b> : 75 % NPK + Biochar @ 2.5 t ha <sup>-1</sup> + FYM@ 7.5 t ha <sup>-1</sup>	62.13	66.23	64.18	67.68	72.36	70.02
<b>T<sub>9</sub></b> : 75 % NPK + Biochar @ 5.0 t ha <sup>-1</sup> + FYM@ 7.5 t ha <sup>-1</sup>	64.4	68.12	66.26	70.84	77.80	74.32
<b>T<sub>10</sub></b> : 100 % NPK + Biochar @ 2.5 t ha <sup>-1</sup>	54.17	57.45	55.81	66.19	72.20	69.20
<b>T<sub>11</sub></b> : 100 % NPK + Biochar @ 5.0 t ha <sup>-1</sup>	57.57	60.91	59.24	69.55	75.70	72.63
<b>T<sub>12</sub></b> : 100 % NPK + Biochar @ 2.5 t ha <sup>-1</sup> + HA @ 10 kg ha <sup>-1</sup>	77.82	81.87	79.85	97.21	103.45	100.33
<b>T<sub>13</sub></b> : 100 % NPK + Biochar @ 5.0 t ha <sup>-1</sup> + HA @ 10 kg ha <sup>-1</sup>	80.25	85.56	82.91	101.45	110.07	105.76
<b>T<sub>14</sub></b> : 100 % NPK + Biochar @ 2.5 t ha <sup>-1</sup> + FYM @ 7.5 t ha <sup>-1</sup>	73.78	77.49	75.64	86.04	96.02	91.03
<b>T<sub>15</sub></b> : 100 % NPK + Biochar @ 5.0 t ha <sup>-1</sup> + FYM@ 7.5 t ha <sup>-1</sup>	76.40	80.77	78.59	90.21	101.57	95.89
<b>S. Em.±</b>	<b>0.97</b>	<b>1.22</b>	<b>1.09</b>	<b>1.43</b>	<b>3.01</b>	<b>1.98</b>
<b>C.D. at 5 %</b>	<b>2.84</b>	<b>3.55</b>	<b>3.19</b>	<b>4.14</b>	<b>8.71</b>	<b>5.75</b>

**Note:**

HA : Humic acid  
FYM : Farmyard manure

**Table 5: Residual effect of biochar and humic acid on grain and haulm yield of chickpea in maize-chickpea cropping system under rainfed condition**

Treatments	Seed yield (q ha <sup>-1</sup> )			Haulm yield ((q ha <sup>-1</sup> ))		
	2019	2020	Pooled	2019	2020	Pooled
<b>T<sub>1</sub></b> : Absolute control	7.24	6.91	7.08	10.75	10.01	10.38
<b>T<sub>2</sub></b> : 100 % NPK + FYM @ 7.5 t ha <sup>-1</sup>	10.16	11.29	10.73	12.45	13.65	13.05
<b>T<sub>3</sub></b> : T <sub>2</sub> + HA @ 10.0 kg ha <sup>-1</sup>	12.58	14.57	13.58	16.45	18.17	17.31
<b>T<sub>4</sub></b> : 75 % NPK + Biochar @ 2.5 t ha <sup>-1</sup>	9.44	10.34	9.89	11.41	12.19	11.80
<b>T<sub>5</sub></b> : 75 % NPK + Biochar @ 5.0 t ha <sup>-1</sup>	9.97	10.66	10.32	11.86	12.67	12.27
<b>T<sub>6</sub></b> : 75 % NPK + Biochar @ 2.5 t ha <sup>-1</sup> + HA @ 10 kg ha <sup>-1</sup>	11.71	13.12	12.42	14.21	15.62	14.92
<b>T<sub>7</sub></b> : 75 % NPK + Biochar @ 5.0 t ha <sup>-1</sup> + HA @ 10 kg ha <sup>-1</sup>	12.10	13.34	12.72	14.58	17.18	15.88
<b>T<sub>8</sub></b> : 75 % NPK + Biochar @ 2.5 t ha <sup>-1</sup> + FYM@ 7.5 t ha <sup>-1</sup>	10.94	11.79	11.37	13.26	14.74	14.00
<b>T<sub>9</sub></b> : 75 % NPK + Biochar @ 5.0 t ha <sup>-1</sup> + FYM@ 7.5 t ha <sup>-1</sup>	11.29	12.42	11.86	13.87	15.44	14.66
<b>T<sub>10</sub></b> : 100 % NPK + Biochar @ 2.5 t ha <sup>-1</sup>	9.61	10.54	10.08	12.29	13.15	12.72
<b>T<sub>11</sub></b> : 100 % NPK + Biochar @ 5.0 t ha <sup>-1</sup>	10.05	11.14	10.60	12.74	14.22	13.48
<b>T<sub>12</sub></b> : 100 % NPK + Biochar @ 2.5 t ha <sup>-1</sup> + HA @ 10 kg ha <sup>-1</sup>	12.75	14.88	13.82	16.79	20.54	18.67
<b>T<sub>13</sub></b> : 100 % NPK + Biochar @ 5.0 t ha <sup>-1</sup> + HA @ 10 kg ha <sup>-1</sup>	15.11	18.20	16.66	19.39	22.10	20.75
<b>T<sub>14</sub></b> : 100 % NPK + Biochar @ 2.5 t ha <sup>-1</sup> + FYM @ 7.5 t ha <sup>-1</sup>	12.18	14.18	13.18	14.78	17.55	16.17
<b>T<sub>15</sub></b> : 100 % NPK + Biochar @ 5.0 t ha <sup>-1</sup> + FYM@ 7.5 t ha <sup>-1</sup>	12.55	14.45	13.50	16.22	18.01	17.12
<b>S. Em.±</b>	<b>0.77</b>	<b>0.92</b>	<b>0.85</b>	<b>0.84</b>	<b>0.98</b>	<b>0.91</b>
<b>C.D. at 5 %</b>	<b>2.22</b>	<b>2.66</b>	<b>2.41</b>	<b>2.42</b>	<b>2.82</b>	<b>2.56</b>

**Note:**

HA : Humic acid  
 FYM : Farmyard manure