

## **Original Research Article**

### **Biofertilizing and nématodes control potentials of four isolates of the native arbuscular mycorrhizal fungus on sweet pepper (*Capsicum annuum*) in Togo**

---

#### **ABSTRACT**

**Aims:** The aim of the study was to investigate the effectiveness of four isolates of the native Arbuscular Mycorrhizal Fungus (AMF) in improving pepper plant yield as well as to assess their potential for the control of parasitic nematodes.

**Study design:** The experimental set up consisted of a Randomized Block Design with 5 treatments corresponding to the four isolates of AMF– BEN10, GM142, 472, and WA330 – tested in comparison with the untreated plants as a control treatment

**Place and Duration of Study:** The study was conducted in 2020 from June to September at « at the Research Station of the Faculty of Agronomy, University of Lomé, Togo».

**Methodology:** The four isolates of AMF were inoculated to the seeds in the nursery. Six weeks after, the seedlings were transplanted on 3mx6m beds. Several growth parameters such as the number of leaves and branches, plant height, number of flower buds and fruit weight were recorded each three weeks after transplantation. Nematode density was assessed four times (before transplanting, at the flowering, during fruiting, and after the last harvest of pepper fruits).

**Results:** The AMF increased the marketable pepper fruit weight by 39%, while reducing significantly the root nematode density by 20-34%. A positive correlation ( $P < 0.0001$ ;  $r = 0.816$ ) was observed between the mycorrhization frequency and mycorrhization intensity, the former varying from 70% to 91% and the latter from 60% to 85%. The plant height, the number of leaves, the branches and the number of flower buds were not affected by AMF inoculation. **Conclusion:** The present study shows the potential of AMF to be considered candidate alternative to chemical fertilizers and pesticides for sustainable production of sweet pepper in Togo.

**Keywords:** arbuscular mycorrhizal fungi, mycorrhization, nematodes, reduction, yield.

## 1. INTRODUCTION

The sweet pepper, *Capsicum annuum*, is a fresh dietary vegetable fruit rich in vitamin C and provitamin A (carotene) [1]. The sweet pepper is an excellent source of many nutrients and important secondary metabolites for human health (potassium, flavonoids, antioxidants), and which are known to reduce the risk of many diseases such as cancer and cardiovascular diseases [2]. This crop is very popular crop that contributes to the income of the smallholder farmers, with high international export potential. [2].

In Togo, the sweet pepper is grown mainly in the urban and peri-urban areas. Although these production areas increase each year to respond to the increase in the market demand at the national and international levels the yield remains very low [3]. This low productivity of pepper plants is attributed to several factors among which the soil fertility, coupled with the impact of some biotic components including insect pests, diseases and roots feeders. Among roots feeders, nematodes are the most important pests [3,4,5], the key damaging one being *Meloidogyne* spp. [6, 7]. Currently, the management of nematode pests on vegetable crops including sweet pepper is done mainly using the synthetic chemical nematicides [8, 1]. However, their highly hazardous nature has led to many of these products being removed from the market and their use discontinued [9]. Other nematode management practices such as botanical [10, 11], organic fertilizers [12, 13] or cultural control [14] have been explored for vegetables crops, with some success. There is therefore a need to find alternatives to chemicals for the sustainable production

Recently, many studies established that Arbuscular mycorrhizal (AM) fungi favor plant growth by improving nutrient acquisition [15], but also by increasing their resistance against abiotic and biotic stress, including nematodes [16, 17, 18]. Thus, control of the nematodes using mutualistic micro-organisms such as Arbuscular mycorrhizal fungi (AMF) has been suggested as a potential alternative to chemical control [16]. AMF have mutualistic relationships with more than 80% of terrestrial plant species [19]. This symbiotic relationship is ancient and would have had important roles in establishment of plants on land [19]. In this symbiosis, the fungus provides the plant host with mineral nutrients, especially phosphate, receiving in turn carbohydrates [19, 20]. In this way, the association with AMF can improve the provision of poorly mobile nutrients, especially phosphorus (P), but also ammonium, copper, zinc and other micronutrients [21]. Fungi seems to generate a smaller carbon cost per absorption area unit than roots, and they also allow a higher exploration of

soil not accessed by roots [19]. Moreover, other potential benefits of mycorrhizal symbiosis have been mentioned, such as improved plant water relations and reduction of pathogenic infections [16], promotion of soil aggregation [21] as well as synergistic effects with other microorganisms [22, 23]. We hypothesize that AMF can constitute a viable alternative to the use of chemical fertilizers and pesticides for sustainable sweet pepper production without adverse effects on the environment and human health in Togo. Therefore, the present study was conducted to evaluate the biofertilizing and the pests control potentials of AMF on sweet pepper under field conditions.

## **2. MATERIAL AND METHODS**

### **2.1. Expérimental site**

The field experiment was conducted at the Research Station of the Faculty of Agronomy, University of Lomé, Togo (6°10.563N and 1°12.782E). The site is characterized by Guinean climate with two rainy seasons, April to July and September to November with two dry seasons in between. The soil of the experimental site is classified as a ferralsol soil [24] with the following characteristics: organic matter (OM) 1.87%; total N 0.15%; pH 6.50; available Phosphorus (P<sub>2</sub>O<sub>5</sub>) 0.5 mg/kg; Potassium (K<sub>2</sub>O) 0.46mg/kg and Magnesium (MgO) 0.01mg/kg [24]

### **2.2. Mycorrhiza inoculum used**

Four pure isolates (Ben 10, 472, WA330, GM142 at a dose of 6000 spores) were used. These isolates were obtained from the "Laboratoire des Sciences Agronomiques et de Biologie Appliquée", (La.S.A.B.A.-University of Kara).

### **2.3. Nursery and inoculation of seedling**

The nursery is done in greenhouse. The substrate used for the nursery consisted the mixture (w/w, 1:2) of the arable soil collected at the agronomy station and the beach sand. The soil

was collected from a depth of 0–25 cm and passed through a 1 mm aperture sieve to remove roots and debris. The beach sand was thoroughly washed with tap water to remove salt. The substrate mixture was sterilized at 80 °C for 72 h. Substrate pH (H<sub>2</sub>O) was 7.7, the organic carbon 20 g C kg<sup>-1</sup>, and the total N and available P (P-Brayl) were 3.4 g N kg<sup>-1</sup> and 19 mg P kg<sup>-1</sup>, respectively, analyzed at the ITRA (Institut Togolais de la Recherche Agronomique).

Four AMF inocula (Ben 10, 472, WA330, GM142 at a dose of 6000 spores) were used to inoculate seedlings. Seeds inoculation was done during nursery period in plastic tanks? (50×40×20cm) in the greenhouse. The tank filled with sterilized substrate was watered and three stripes were made in the length direction of the plastic tank about 1cm deep as a seedbed. Thereafter, 50g of corresponding isolates inoculum was spread in each stripe before putting the seeds and closed it with the sterilized marine sand. The control plastic tank had not received any AMF, but sterilized substrate used for inocula production. One plastic tank was used for each inoculum making in total, five plastic tanks.

#### **2.4. Experimental design**

The treatments were arranged in a completely randomized block design with five treatments with four repetitions. Each block consisted of 3×6m plots, separated by one-meter space, while blocks were separated by 2 meters. Six-weeks old plants from the nursery were transplanted at 25×25cm in each plot, making a total number of plants per plot was 120. The plots were regularly watered and weeded until harvest.

#### **2.5. Assessment of AMF root colonization**

Root colonization by AMF was assessed two months after transplanting in the field. Roots were extracted by wet sieving [25]. AMF root colonization was determined according to [26], using trypan blue to stain mycorrhizal structures. A 1.0g subsample of the roots excised from the five plants, to assess the percentage of AMF colonisation. At 90~ on a hot plate, the root samples were cleared in KOH (100g/l) for 1 h and stained with trypan blue (0.5g/l) in lactoglycerol (26) at 90~ for 30 min. Percentage colonization of host plant roots was estimated by visual observations of stained root segments mounted in lactoglycerol by the grid-line intercept method (27) by subtracting the initial weight of the filter paper from the weight of mycelia and filter paper

## **2.6. Effect of AMF on the agronomic parameters of sweet pepper**

Starting from the third week after transplanting and at a frequency of three weeks, twenty plants per treatment (five per plot unit) were examined and parameters such as plant height (from crown to apex), number of leaves, number of branches were assessed.

## **2.7. Yield estimation (kg/18 m<sup>2</sup>)**

A total of five fruits harvests have been done, the first harvest was done two months after the transplantation of plants, and subsequently at 15-days intervals until the end of harvest, leading to a total of five harvests for a total harvest period that lasted two months. The pepper fruits collected at each harvest time on each elementary plot were weighted to determine the pepper fresh fruit weight (Kg) per plot per harvest. The fruit weights of the five harvests were then summed up to determine the total weight of harvested pepper fruits per elementary plot. The mean mean pepper fresh fruit weight (kg/18 m<sup>2</sup>) for each treatment was calculated by adding the total weight from the five replication and divided the result by five.

## **2.8. Sampling, extraction and evaluation of the density of nematodes**

The soil was collected from up to a depth of 15- 20 cm from different treatments plots and 1cm from the selected plant to be sampled. In each plot, three soil samples were collected by boring at different locations randomly selected, and mixed according to the method described by [25] using a plastic bag to form a representative sample of about 600g per plot. In the laboratory, pepper plant roots contained in each soil sample were removed and used to determine the density of the root nematodes, after crushing the roots with moulinex. Subsequently, nematode density in the root-free soil was also determined. Extraction of the nematodes from soil/root sample was done using a modified Baermann plate method [28].. For both purpose, 100 g of soil or 5 g of roots were weighed in the laboratory. Roots were previously cleaned and crushed using a moulinex. Each sample is weighed into a sieve lined inside by the toilet paper for filter and the whole is placed in a plastic basin. Each sample is scattered in the screen using tweezers. Then water was added until the sample was lightly covered, thus promoting the migration of nematodes to the water which is the extraction medium.

After 24 hours incubation for the soil samples and 48 hours for the roots, the sieve containing filter paper on which is deposited the sample is gently removed from the basin. The water from the bowl containing the nematodes is collected in graduated tubes and allowed to settle for 30 minutes. Then, the volume of the extract medium was then reduced

to 100 ml which was used as the final extract medium for the observation of nematodes. 10ml is removed from the 100 ml extract medium using a pipette in a petri dish and nematodes were counted with a Leica Wild M3C microscope. This operation is repeated three times for each sample.

## **2.9. Statistical analysis**

Data on mycorrhizal parameters, agronomic performance, and pest pressure were used to compare the different treatments by the Generalized Linear Model (GLM) procedure using SPSS 25 (Statistical Package for the Social Sciences) version 2018. Data on density and percentages were log- or arcsin-transformed, respectively, before being subjected to statistical analysis [29]. In the case of significant differences, means were discriminated using the Student-Newmann-Keuls) multiple range test.

## **3. RESULTS AND DISCUSSION**

### **3.1. Results**

#### **3.1.1. Sweet Pepper root rate colonization by mycorrhizal fungus**

The presence of mycorrhizal structures was detected in all the plants at the three stages of assessment. The Roots mycorrhization was observed in all the plants included the control plant. (figure1) and the intensity of root mycorrhization was found to increase with the age of the plants. Overall, the AMF colonization frequency was significantly higher ( $P < .0001$ ) for the AMF-treated plants compared to the untreated plants (figure 2), with the isolates GM 142 and WA472, recording the highest frequencies. A positive correlation was found between mycorrhization frequency and mycorrhization intensity ( $P < .0001$ ;  $r = 0.816$ ) (Figure 3).




Figure 1. Mycorrhization intensity in the roots of bell pepper plants inoculated or not with four AMF isolates (Ben10, GM142, 472 and WA330).  
 Test: significant from normal control \* $P < 0.05\%$ . , Average mean  $\pm$  SE = Standard error.




Figure 2: Mycorrhization frequency in the roots of bell pepper plants inoculated or not with AMF isolates (Ben10, GM142, 472 and WA330).  
 Test: significant from normal control \* $P < 0.05\%$ . , Average mean  $\pm$  SE = Standard error.





Figure 1: Relationship between mycorrhization frequency and mycorrhization intensity from the root of sweet pepper inoculated from four AMF isolates (Ben10, GM142, 472 and WA330).

### 3.1.2. Effect of AMF on height, number of branches, number of leaves and number of flower buds

Data on plant height, number of branches, number of leaves and number of flower buds are shown in Figure 4, Table 1, Table 2 and Table 3, respectively. These results show that the supply of AMF did have any effect on these parameters, although the mycorrhizal plants showed a slight increase in vivacity compared to the control in the range of 1.51% to 4.22% in week 9.. Only the mean number of flower buds was significantly different in week 11. Plants inoculated with GM142 and 472 had 17.44% and 22.14% respectively more flower buds compared to the control.



**Figure 4:** Effects of four AMF isolates (Ben10, GM142, 472 and WA330) on the height of sweet pepper plant at 4 growing stages.

**Table 1:** Effects of four AMF isolates (Ben10, GM142, 472 and WA330) on the number of branches of sweet pepper plant at four interval times of growing stage.

| AMF isolates      | Number of branches per plant (mean ± SE) |             |             |              |
|-------------------|--|-------------|-------------|--------------|
|                   | Week 2                                   | Week 5      | Week 7      | Week 9       |
| <b>BEN10</b>      | 0.00±0.00a                               | 4.35±0.48a  | 10.70±0.58a | 13.40±0.37a  |
| <b>GM142</b>      | 0.00±0.00a                               | 6.00±3.92a  | 12.45±0.98a | 14.65±0.76a  |
| <b>Strain 472</b> | 0.00±0.00a                               | 6.25±1.20a  | 12.80±1.23a | 16.75±0.96a  |
| <b>WA330</b>      | 0.00±0.00a                               | 4.95±0.70a  | 11.45±0.76a | 13.75±0.49a  |
| <b>Witness</b>    | 0.00±0.00a                               | 5.90±0.63a  | 12.45±0.65a | 13.60±0.53a  |
| <b>cv%</b>        |  | <b>9.6</b>  | <b>18.6</b> | <b>15.67</b> |
| <b>p</b>          |  | <b>0.42</b> | <b>0.41</b> | <b>0.52</b>  |

NB: Comparisons are made by column. Means in the same column followed by the same lower case letter are not statistically different (Student-Newman-Keuls,  $P < 0.05$ ). SE = Standard

**Table 2 :** Effect of AMF isolates (Ben10, GM142, 472 and WA330) on the average number of leaves on sweet pepper plant at different times of growing stage.

| AMF isolates      | Average number of leaves per plant (mean $\pm$ SE) |                   |                   |                   |
|-------------------|--|-------------------|-------------------|-------------------|
|                   | Week 2   | Week 5            | Week 7            | Week 9            |
| <b>BEN10</b>      | 7.20 $\pm$ 0.46a                                   | 34.15 $\pm$ 3.24a | 59.09 $\pm$ 2.63a | 68.45 $\pm$ 2.09a |
| <b>GM142</b>      | 7.35 $\pm$ 0.37a                                   | 41.40 $\pm$ 5.49a | 62.10 $\pm$ 4.04a | 70.30 $\pm$ 3.17a |
| <b>Strain 472</b> | 7.15 $\pm$ 0.35a                                   | 40.00 $\pm$ 3.25a | 57.15 $\pm$ 2.24a | 67.60 $\pm$ 2.64a |
| <b>WA330</b>      | 7.20 $\pm$ 0.31a                                   | 43.25 $\pm$ 5.11a | 59.50 $\pm$ 4.07a | 68.05 $\pm$ 3.46a |
| <b>Witness</b>    | 7.40 $\pm$ 0.32a                                   | 44.65 $\pm$ 4.46a | 59.15 $\pm$ 3.78a | 66.35 $\pm$ 2.55a |
| <b>cv%</b>        | <b>15.4</b>  | <b>11.4</b>       | <b>14.9</b>       | <b>15.7</b>       |
| <b>p</b>          | <b>0.97</b>  | <b>0.49</b>       | <b>0.88</b>       | <b>0.90</b>       |

NB: Comparisons are made by column. Means in the same column followed by the same lowercase letter are not statistically different (Student-Newman-Keuls,  $P < 0.05$ ). SE = Standard error.

**Table 3:** Effect of AMF isolates (Ben10, GM142, 472 and WA330) on the average number of flower buds sweet pepper at four different times of growing stage.

| AMF strains    | Number of flower buds of bell pepper (mean $\pm$ SE) |                   |                   |                    |
|----------------|--|-------------------|-------------------|--------------------|
|                | Week 5   | Week 7            | Week 9            | Week 11            |
| <b>BEN10</b>   | 0.95 $\pm$ 0.23a                                     | 9.05 $\pm$ 1.46a  | 17.70 $\pm$ 0.77a | 16.90 $\pm$ 0.78ab |
| <b>GM142</b>   | 0.50 $\pm$ 0.50b                                     | 8.40 $\pm$ 1.33a  | 17.65 $\pm$ 1.02a | 17.50 $\pm$ 0.52a  |
| <b>472</b>     | 0.35 $\pm$ 0.22b                                     | 11.85 $\pm$ 1.96a | 19.10 $\pm$ 0.80a | 18.20 $\pm$ 0.80a  |
| <b>WA330</b>   | 0.25 $\pm$ 0.12b                                     | 8.85 $\pm$ 1.51a  | 16.35 $\pm$ 0.66a | 15.40 $\pm$ 0.49b  |
| <b>control</b> | 0.20 $\pm$ 0.09b                                     | 10.60 $\pm$ 1.88a | 16.30 $\pm$ 0.90a | 14.90 $\pm$ 0.75b  |
| <b>cv%</b>     | 14.9   | 9.75              | 11.6              | 14.2               |
| <b>p</b>       | 0.02   | 0.55              | 0.12              | 0.04               |

NB: Comparisons are made by column. Means in the same column followed by the same lower-case letter are not statistically different (Student-Newman-Keuls,  $P < 0.05$ ). SE = Standard error.

### 3.1.3. Effects of AMF isolates inoculation on sweet pepper fruit weight

The fresh fruit weight of sweet pepper is shown in Table 4. The results showed a significant effect of AMF isolates on pepper fresh fruit weight. The highest weight was obtained with the inoculum 472 with an average of 8.73 kg/18 m<sup>2</sup> about 39-fold higher. It increases the fruit fresh fruit pepper by 39.45% compared that obtained for untreated plants.

Table 4: Effects of four AMF isolates (Ben10, GM142, 472 and WA330) on the sweet pepper fresh fruit weight and average fresh fruit weight gain.

| Mycorrhizae strains | Fruit yield over the whole period in kg/18 m <sup>2</sup> (average ± SE) |                |
|---------------------|--|----------------|
|                     | Yield  | Gain in weight |
| <b>BEN10</b>        | 6.99±1.48b   | <b>11,66%b</b> |
| <b>GM142</b>        | 6.89±1.51b   | <b>10,06%b</b> |
| <b>472</b>          | 8.73±1.01a   | <b>39,45%a</b> |
| <b>WA330</b>        | 7.12±1.35ab  | <b>13,73%b</b> |
| <b>Control</b>      | 6.26±1.49b   | -              |
| <b>cv%</b>          | <b>13.2</b>  | <b>13.6</b>    |
| <b>p</b>            | <b>0.02</b>  | <b>0.03</b>    |

NB: Comparisons are made by column. Means in the same column followed by the same lower-case letter are not statistically different (Student-Newman-Keuls,  $P < 0.05$ ). SE = Standard Error

### 3.1.4. Effects of AMF isolates on nematode density

The results regarding nematode density from the roots and the soil are shown in the Table 5. Before transplanting, the result shows that the soil was heavily infected with nematodes. No significant differences between plots are observed ( $P = 0,09$ ). The same trend was observed between treatments regarding nematode's density at the flowering, the fruiting and at the harvest ( $P = 0,08$ ,  $P = 0,78$  and  $0,09$  respectively).

Concerning the nematode density in the roots, a progressive evolution of the number of nematodes in the roots of sweetpepper has been noted from the transplanting to the harvest for all treatments (Table5). The analysis of the variance shows that the reduction of nematodes density by each of the four AMF strains used is statistically significant compared

to control without AMF at the flowering and at the harvest ( $P \leq 0,04$ ). However, comparison within the different strains of AMF used reveals similar actions.

The correlation analysis (figure 5) shows a negative linear relationship between mycorrhization intensity and nematode population density ( $P < 0.05$ ;  $r = - 0.48$ ). Indeed, it was found that nematode density decreased with plant age while mycorrhization intensity did not increase.

\*




Figure 5: Relationship between mycorrhization intensity from four AMF isolates (Ben10, GM142, 472 and WA330) and nematode density in the root of sweet pepper.

**Table 5:** Effect of four AMF isolates (Ben10, GM142, 472 and WA330) on average number of nematodes (Mean± SE) extracted from soil and roots of sweet pepper, before transplanting, at flowering, during fruiting and at harvest.

| Mycorrhizae strains | Density of nematodes in soil and roots (mean ± SE) |                        |              |                 |              |                |              |
|---------------------|--|------------------------|--------------|-----------------|--------------|----------------|--------------|
|                     | Before transplanting                               | Beginning of flowering |              | During fruiting |              | After harvest  |              |
|                     | Soil   | Soil                   | Root         | Soil            | Root         | Soil           | Root         |
| <b>BEN10</b>        | 129.00 ±17.43a                                     | 161.6 ±15.91a          | 19.51 ±3.22a | 182.53 ±12.11a  | 20.5 ±3.21a  | 128.31 ±13.15a | 19.13 ±6.83b |
| <b>GM142</b>        | 129.10 ±17.11a                                     | 183.3 ±20.96a          | 20.81 ±3.21a | 168.34 ±10.51a  | 18.61 ±3.12a | 139.15 ±7.14a  | 21.65 ±6.62b |
| <b>472</b>          | 129.12 ±18.64a                                     | 180.8 ±54.60a          | 20.83 ±3.93a | 170.00 ±42.41a  | 20.82 ±3.91a | 108.37 ±12.92a | 21.14 ±5.91b |
| <b>WA330</b>        | 129.10 ±18.31a                                     | 283.3 ±43.66a          | 15.84 ±2.21b | 198.36 ±18.83a  | 16.3 ±2.21a  | 165.00 ±2.26a  | 17.10 ±7.00b |
| <b>Control</b>      | 129.1 ±11.10a                                      | 272.51 ±42.10a         | 22.10 ±3.32a | 248.31 ±22.72a  | 20.12 ±3.33a | 170.84 ±14.32a | 27.11 ±9.11a |
| <b>cv%</b>          | 14.9   | 22.7                   | 17.1         | 11.7            | 16.2         | 21.8           | 11.2         |
| <b>P</b>            | 0.09   | 0.08                   | 0.04         | 0.78            | 0.89         | 0.09           | 0.04         |

NB: Comparisons are made by column. Means in the same column followed by the same lower-case letter are not statistically different (Student-Newman-Keuls,  $P < 0.05$ ). SE = Standard Error

### 3.2. Discussion

This study on the effect of four local AMF isolates on growth parameters, fruit yield and nematodes pressure of sweet pepper is one of the first carried out in the vegetable crops in Togo. The use of spores as inoculum to inoculate sweet pepper plants must meet two main criteria; on the one hand, the fungus must be able to intensively colonize the sweet pepper roots and on the other hand, this colonization must promote yield increase [18, 20] and the reinforcement of the plants' immune system for protection against nematode pests [30, 18, 31].

From the experiment, it was found that most of the sweet pepper plants inoculated with the different isolates of AMF showed mycorrhizal structures in their root cortex, even the control plants. Our result suggests sweet pepper to be a mycorrhizal plant that is compatible for a symbiosis relationship with AMF [32]. The mycorrhization of the control plants a few weeks after transplanting could be due to the presence of native mycorrhizal strains in the garden soil [19]. The frequency and intensity of mycorrhization of inoculated plants varied slightly according to the isolates used [33]. In most cases, the beneficial effect of mycorrhizae is due to an improvement of the mineral nutrition of the host plant, especially with regard to the elements that are not very mobile, such as P, Zn, Cu, and does not exclude nitrogenous nutrition [34] and also the photosynthesis activities from the plant [34, 35]. The more the mycorrhizal growth is increased, the more the number and the leaf surface are accentuated [36, 37]. In the present study, yet no significant difference was observed during the trial on the growth parameters (height, number of leaves, branching) of the plants, regardless of the treatment. Indeed, [38] reported that the growth response of plants due to AMF infection could be positive, negative or neutral depending on many factors (edaphic, environmental, mycorrhizal and/or plant). [39] showed in a similar study that infection of wheat with *Glomus falciculatus* increased its drought resistance and helped the plant to grow, whereas *Glomus mosseae* had no effect. Therefore, it would be appropriate to look for the most suitable isolates in terms of their effectiveness on the growth parameters of bell pepper. The results of the present experiment showed that the contribution of AMF seems rather beneficial in improving the fruit yield of bell pepper. Isolates 472 optimized yield (by more than 39% compared to the control) [40]. However, the yields obtained in our study (8.73 t/ha obtained by isolates 472) are below those obtained in Egypt 14.9 t/ha [41] and in the sub-region. This suggests other yield parameters including intrinsic and extrinsic characteristics (climate, soil, pests, diseases, etc.) of the crop [41, 42]. It is also possible to think of combination fertilizer with AMF to increase yields, because, according to [42] the humus factor is of capital importance for demanding crops such as Solanaceae.

Concerning the nematode density, the results showed that there was no significant difference between treatments for soil nematodes before transplanting at which 129 nematodes per 100g soil on average has been found. This presence of nematodes can be explained by the humectation of superior layer of soil by watering dragging their migration toward the superior layer and due to the presence of the grasses which are the natural host of nematodes. [43, 44].

In relation root nematodes/ml, the significant difference was observed between inoculated plants and control plants at flowering and after harvest. These results are in line with many other studies which have reviewed the effects of AMF on plant growth and their interactions [41, 42]. A general conclusion from these reviews suggests that AMF increase resistance to nematode infestation by slowing down nematode development. [43] established that the efficiency of the arbuscular mycorrhizal fungi native to Benin were as well very efficient in greenhouse and in the field to reduce the rate of

*Meloidogyne* spp density in the soil of culture and in the roots of the tomato. In the other hand, the present results showed that the effect of AMF did not block the multiplication of nematodes but reduced their multiplication rate and the action would not be direct but rather indirect [44]. However, the reduction level of nematodes density in the roots of *S. annuum* is not linear, which can be explained by the fact that the effect of AMF inoculation was not constant during the experiment [45, 46]. The lack of effectiveness consistency may be attributed to several factors, including slight variation in experimental set up, but more possibly different feeding styles of nematodes assessed [47].

#### 4. Conclusion

This study showed AMF as alternatives to chemical fertilizers and synthetic pesticides for sustainable production of sweet pepper in Togo. Each isolate of AMF tested was able to reduce population density of nematodes on roots and promote increase in pepper fruits yield. In terms of perspectives, we believe that the identification of indigenous strains for nematodes control would be an important step towards the quantitative and qualitative improvement of sweet pepper yield in Togo. Similarly, the strains should be tested on other crops, especially those known to be susceptible to nematodes.

#### CONSENT (WHERE EVER APPLICABLE)

Not applicable

#### ETHICAL APPROVAL (WHERE EVER APPLICABLE)

Not applicable

#### REFERENCES

- [1] Fratianni F., d'Acerno A., Cozzolino A., Spigno P., Riccardi R., Raimo F et al. Biochemical Characterization of Traditional Varieties of Sweet Pepper (*Capsicum annuum* L.) of the Campania Region, Southern Italy. *Antioxidants* 2020, 9, 556; doi:10.3390/antiox9060556
- [2] FAO (Food and Agriculture Organization of the United Nations) Crop prospects and food situation—quarterly global report No. 1. *Rome*. 2020
- [3] Kanda, M., Wala, K., Batawila, K., Djaneye-Boundjou, G., Ahanchede, A., & Akpagana, K.. Le maraîchage périurbain à Lomé : Pratiques culturelles, risques sanitaires et dynamiques spatiales. *Cahiers Agricultures*. 2009; 18(4), 356-363. French.
- [4] Grassini, P., Eskridge, K. & Cassman, K. Distinguishing between yield advances and yield plateaus in historical crop production trends. *Nat Commun*. 2013; 4, 2918. <https://doi.org/10.1038/ncomms3918>
- [5] Zhao, C., B. Liu, S. Piao, X. Wang, D.B. Lobell, Y. Huang, M. Huang, et al: Temperature increase reduces global yields of major crops in four independent estimates. *Proc. Natl. Acad. Sci*. 2017; no. 35, 9326-9331, doi:10.1073/pnas.1701762114.
- [6] Sikora RA, Bridge J, Starr JL. Management practices: an overview of integrated nematode management technologies. In, Luc M, Sikora RA, Bridge J, eds. *Plant parasitic nematodes in Subtropical and Tropical Agriculture*. 2nd Edition, CABI Wallingford, UK. 2005; p 793–825.
- [7] Collange B, Navarrete M, Peyre G, Mateille T, Tchamitchian M. Root-knot nematode (*Meloidogyne*spp. ) management in vegetable crop production: The challenge of an agronomic system analysis. *Crop Prot*.2011; 30(10): 1251–1262.
- [8] Abbas H, Javed N, Khan SA, Ahmad S. Exploitation of the Nematicidal Potential of Bio- and Synthetic Chemicals Against *Meloidogyne incognita* and Their Impact on Phytotoxicity and Nematode Reproduction. *Pak J Zool* 2015; 47(6): 1587–600.

- [9] Aktar MW, Sengupta D, Chowdhury A. Impact of pesticides use in agriculture: their benefits and hazards. *Interdisc Toxicol* 2009; 2(1): 1-12.
- [10] D'Addabbo T, Laquale S, Lovelli S, Candido V, Avato P. Biocide plants as a sustainable tool for the control of pests and pathogens in vegetable cropping systems. *Ital J Agron*. 2014; 9(4): 137–45.
- [11] Rocha TL, Soll CB, Boughton BA, Silva TS, Oldach K, Firmino AAP, et al. Prospection and identification of nematotoxic compounds from *Canavalia ensiformis* seeds effective in the control of the root knot nematode *Meloidogyne incognita*. *Biotechnol Res Innov*. 2017; 1(1): 87–100.
- [12] McSorley, R. Nematodes. In Nanipieri P., Paul E., Li Y. and Sumner M. (eds) *Handbook of soil science*; 2<sup>nd</sup> Ed. Taylor and Francis. 2011; Pp 25-5 to 25-12.
- [13] Bissadou, K. D., Tchabi, A., Tounou, A. K., Ayessom, A. & Gumedzoe, M. Impact de la fumure organique appliquée seule et en combinaison avec une souche indigène de champignon mycorrhizien arbusculaire *Glomus mosseae* sur *Meloidogyne* spp, principal nématode parasitaire de la tomate au Togo. *Journal of Applied Biosciences*. 2012; 55: 3973- 3986. French
- [14] Claudius-Cole AO, Fawole B, Asiedu R, Coyne DL (2014) Management of *Meloidogyne incognita* in yam-based cropping systems with cover crops. *Crop Prot* 63:97–102
- [15] Clark, R.B., and S.K. Zeto. Mineral acquisition by arbuscular mycorrhizal plants. *Journal of Plant Nutrition*. 2000; 23:867-902.
- [16] Hol W H G et Cook R. An overview of arbuscular mycorrhizal fungi–nematode interactions. *Basic an Applied Ecology*. 2005; 6: 489-503.
- [17] Tchabi A, Hountondji F C C, Ogunsola B, Lawouin L, Coyne D, Wiemken A. Effect of two species of arbuscular mycorrhizal fungi inoculation on development of micro-propagated yam plantlets and suppression of *Scutellonemabradys* (Tylenchidae). *Journal of Entomology and Nematology*. 2016; 8, 1-10.
- [18] Gnamkoulamba, A., Tounou, AK., Agboka, K., Tchao, M., Tchabi, A., Adjevi, K. M. et al. Evaluation of the bioprotective potential of arbuscular mycorrhizal fungi against plant-parasitic nematodes and insect pests of rice in Togo. *Journal of Scientific Research of the University of Lomé*. 2018; 20 (3), 143-166. French
- [19] Smith S E, Read D J. Mineral nutrition, toxic element accumulation and water relations of arbuscular mycorrhizal plants. In: Smith, SE and Read. DJ. *Mycorrhizal Symbiosis*. Mycorrhizal symbiosis. 3<sup>rd</sup> ed. Academic Press, San Diego, California, USA. 2008.
- [20] Pacovsky, R.S., Micronutrient uptake and distribution in mycorrhizal or phosphorus-fertilized soybeans, *Plant and Soil*. 1986; 10.1007/BF02374618, 95, 3, (379-388).
- [21] Koide R., Nutrient supply, nutrient demand and plant response to mycorrhizal infection. *New Phytologist*. 1991; 117: 365-386.
- [22] Srinath, J., Bagyaraj, DJ., Satyanarayana, BN. Enhanced growth and nutrition of micropropagated *Ficus benjamina* to *Glomus mosseae* co-inoculated with *Trichoderma harzianum* and *Bacillus coagulans*. *World Journal of Microbiology and Biotechnology*. 2003; 19(1), 69-72.
- [23] Osorio, NW; Habte, M.. Synergistic effect of a phosphate solubilizing fungus and an arbuscular mycorrhizal fungus on leucaena seedlings in an oxisol fertilized with rock phosphate. *Botany*. 2013; 91: 274-281. DOI: 10.1139/cjb-2012-0226.
- [24] Toundou O. Évaluation des caractéristiques chimiques et agronomiques de cinq composts de déchets et étude de leurs effets sur les propriétés chimiques du sol, la physiologie et le rendement du maïs (*Zea mays* L. Var. *Ikenne*) et de la tomate (*Lycopersicum esculentum* L. Var. *Tropimech*) sous deux régimes hydriques au Togo. Thèse de doctorat de l'Université de Lomé en cotutelle avec l'Université de Limoges. 2016; 213p. French.
- [25] Oehl F, Sieverding E, Ineichen K, Mäder P, Boller T, Wiemken A., Impact of land use intensity on the species diversity of arbuscular mycorrhizal fungi in agroecosystems of Central Europe. *Applied Environmental Microbiology*, 2003; 69: 2816-2824.
- [26] Brundrett, M., Bougher, N. Dell, B. Grove, T., Malajczuk. N. Working with Mycorrhizas in Forestry and Agriculture. ACIAR Monograph Australian Centre for International Agricultural Research, Canberra. 1996; 32, 374 pp.

- [27] Giovannetti M, Mosse B. An evaluation of techniques for measuring vesicular arbuscular mycorrhizal infection in roots. *New Phytologist*, 1980; 84: 489-500.
- [28] Coyne DL, Nicol JM, Claudius-Cole B. Practical plant nematology: A field and laboratory guide. 2007
- [29] Gomez, K A., Gomez, AA. Statistical Procedures for Agricultural Research. 2 édition, John Wiley and Sons Inc., New York, USA. 1984; 680pp
- [30] Smith, FA., & Smith, S. Tansley review no. 96 structural diversity in (vesicular)-arbuscular mycorrhizal symbioses. *The New Phytologist*. 1997; 137 (3), 373-388.
- [31] Yang H., Xu J., Guo Y., Koide R. T., Dai Y., Xu M., et al. Predicting plant response to arbuscular mycorrhizas: the role of host functional traits. *Fungal Ecol.* 2016; 20, P.79–83.
- [32] McGonigle TP, Fitter AH. Ecological specificity of vesicular-arbuscular mycorrhizal associations. *Mycological Research*. 1990; 94, 120–122
- [33] Srinath, J., Bagyaraj, DJ., Satyanarayana, BN. Enhanced growth and nutrition of micropropagated *Ficus benjamina* to *Glomus mosseae* co-inoculated with *Trichoderma harzianum* and *Bacillus coagulans*. *World Journal of Microbiology and Biotechnology*. 2003; 19(1), 69-72.
- [34] Li S., Yang W., Guo J., Li X., Lin J., Zhu X. Changes in photosynthesis and respiratory metabolism of maize seedlings growing under low temperature stress may be regulated by arbuscular mycorrhizal fungi. *Plant Physiology and Biochemistry*. 2020; 154:1-10.
- [35] Liang BB., Wang WJ., Fan XX., Kurakov AV., Liu YF., Song FQ., et al.. Arbuscular mycorrhizal fungi can ameliorate salt stress in *Elaeagnus angustifolia* by improving leaf photosynthetic function and ultrastructure. *Plant Biology*. 2021; 23 232-241.
- [36] Nouaïm R., Chaussod R., - Rôle des mycorhizes dans l'alimentation hydrique et minérale des plantes, notamment des ligneux de zones arides. *Options Méditerranéennes*. 1996 ; 20 :9-26. French.
- [37] Nzanza, B., Marais, D. and Soundy, P. Tomato (*Solanum lycopersicum* L.) seedling growth and development as influenced by *Trichoderma harzianum* and arbuscular mycorrhizal fungi. *African Journal of Microbiology Research*. 2011; 5: 425–431
- [38] Johnson NC, Graham JH, Smith FA. Functioning of mycorrhizal associations along the mutualism–parasitism continuum. *New Phytol.* 1997; 135:575–585
- [38] Mitra D., Djebaili R., Pellegrini M., Bhaswatimayee M., Sarker A., Chaudhary P. et al. Arbuscular mycorrhizal symbiosis: plant growth improvement and induction of resistance under stressful conditions. *Journal of Plant Nutrition*. 2021; 44:13, pages 1993-2028.
- [39] Yang H., Zhang Q., Dai Y., Liu Q., Tang J., Bian X., et al. Effects of arbuscular mycorrhizal fungi on plant growth depend on root system: a meta-analysis. *Plant Soil*. 2015; 389, P. 361–374.
- [40] Laumonier R. Les cultures légumières et maraîchères, tome III. 3e édition. Collection « Encyclopédie Agricole » Editions J-B. Baillière, Paris, France. 1979; 276p. French.
- [41] Smith S, Sally E., Read, DJ. *Mycorrhizal symbiosis*. Academic press. 2010.
- [42] Keikantsemang NN., , Greenhouse and field evaluations of commonly occurring weed species for their host suitability to *Meloidogyne* species. *International Journal of Pest Management*. 2015; 62(1):1-9
- [43] Hol WHG, Cook R. An overview of arbuscular mycorrhizal fungi–nematode interactions. *Basic and Applied Ecology*. 2005; 6: 489-503.
- [44] Gough EC., Owen KJ., Zwart RS., Thompson JP. A Systematic Review of the Effects of Arbuscular Mycorrhizal Fungi on Root-Lesion Nematodes, *Pratylenchus* spp. *Front in Plant Science* (11). 2020; P.11: 923
- [45] Affokpon, A., Coyne, DL., Lawouin, L., Tossou, C., Agbèdè, RD., Coosemans, J. Effectiveness of native West African arbuscular mycorrhizal fungi in protecting vegetable crops against root-knot nematodes. *Biology and fertility of soils*. 2011; 47(2), 207-217.
- [46] Sangwan S., Prasanna R. Mycorrhizae Helper Bacteria: Unlocking Their Potential as Bioenhancers of Plant-Arbuscular Mycorrhizal Fungal Associations. *Microbiology and Ecology*. 2022; 84(1):1-10.

[47] Topalović O, Hussain M, Heuer H. Plants and Associated Soil Microbiota Cooperatively Suppress Plant-Parasitic Nematodes. *Frontiers Microbiology*. 2020; doi:10.3389/fmicb.2020.00313.

**DEFINITIONS, ACRONYMS, ABBREVIATIONS**

Not applicable

UNDER PEER REVIEW