

## Original Research Article

### **Effect of seed soaking and growing media on germination and survivability of acid lime (*Citrus aurantifolia* Swingle)**

#### **Abstract**

The experiment was laid out in a Completely Randomized Design with Factorial concept by three repetitions. The seeds of acid lime were treated with different seed soaking *viz.*, water, GA<sub>3</sub> @ 50 ppm, GA<sub>3</sub> @ 100 ppm, cow urine @ 25 %, cow urine @ 50 % and cow dung slurry (1:1 w/w) and sowing in different growing Media *viz.*, sand, vermicompost, FYM and coco peat. The result indicated that GA<sub>3</sub> @ 100 ppm is found to be the most effective for better germination parameters, growth parameters and survival percentage of acid lime seedlings. Among different growing media Sand + Vermicompost + Cocopeat (1:1:1 v/v/v) recorded higher values for germination parameters, growth parameters and survival percentage of acid lime seedlings. The treatment combination of GA<sub>3</sub> @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v) showed better germination parameters, growth parameters and survival percentage. Therefore, the combination of GA<sub>3</sub> @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v) was found most suitable for better growth of acid lime seedling.

**Keywords:** Acid lime, seed soaking, growing media, germination, survivability

#### **Introduction**

Citrus is one of the most important fruit crops of tropical and subtropical regions. It occupies 3<sup>rd</sup> rank after mango and banana in India. It is grown in over 100 countries and often regarded as golden fruit. The total area under cultivation of acid lime & lemon in India is 316 thousand hectare with production of around 3628 thousand MT (Anonymous, 2020-21). In Gujarat, area under citrus cultivation is 48503 hectare with 625833 MT production (Anonymous, 2020-21). It is mainly grown in Ahmedabad, Kheda, Mehsana, Bhavnagar, Gandhinagar and Junagadh districts of Gujarat.

Propagation of acid lime is generally done by grafting, budding, air layering and by seed. The commercial practice of propagation is by seed (Khatana and others 2015). It produces true to type, because of high degree (39-60 %) of nucellar embryony. The resultant seedlings are known to be free from tristeza virus and thus perform better. In acid lime

germination percentage is low and it takes about 3 weeks to germinate (Cheema and others 1954). Some of the problems faced by acid lime growers are slow, erratic and incomplete germination with high initial seedling mortality. The possible reason of slow germination is presence of the growth inhibitors and physical resistance of seed coat to radical protrusion (Khan and others 2002). The growth of acid lime seedling is very slow in nursery as well as in the field. The increasing germination percentage and producing healthier seedling are major challenges for farmers.

So, the seeds of acid lime cv. Kagzi Lime are soaked with plant growth regulator such as gibberellic acid ( $GA_3$ ) as well as with organic substance such as cow urine, cow dung slurry and water for better seed germination.  $GA_3$  controls mobilization of starch which acts as a respiratory substrate leading to immediate enhancement in cell elongation. Gibberellins also help in enhancing the availability of reserved mineral elements which promote the germination process. Cow urine contains iron, urea, uric acid, estrogens and progesterone which affect the inhibitory responses to seed germination, shoot growth and seedling vigour. Cow dung slurry also contains some growth promoting substances (auxin), N, P, K, micronutrients and biodegradable enzymes which have been cause for softening of seed coat and thereby enhances seed germination and growth of seedlings (Raj and others 2014). Water soaking of seeds is done to modify hard seed coats, remove inhibitors, soften seeds and reduce the time of germination.

Media not only act as a growing place but also as a source of nutrient for plant growth. It is a substrate that provides the required elements and physical support to the growing plants. Vermicompost provides sufficient levels of oxygen to roots, adequate storage of water and nutrient for the plants. FYM is having good water holding capacity as well as sufficient porosity. Cocopeat provides excellent pore space (25-30 %) and fine structure required for proper growth. It is a rich source of nutrients and can easily mixed with other growing media used in seed germination and growth of seedling. Sand is used as a rooting media or adding the coarser texture needed to induce proper drainage and aeration. Keeping in view above, the aims of the study was to identify suitable treatment combinations for better seed germination and seedling growth of acid lime.

### **Materials and Methods**

The experiment was carried out under shade net house at College Farm, College of Horticulture, Sardarkrushinagar Dantiwada Agricultural University, Jagudan, Dist. Mehsana, Gujarat during July, 2021 to December, 2021. The experiment was laid out by Completely Randomized Design with Factorial concept (FCRD), which included 24 treatments with 3

replications, there were two factors which comprised six levels of seed soaking viz., S<sub>1</sub> (Water), S<sub>2</sub> (GA<sub>3</sub> @ 50 ppm), S<sub>3</sub> (GA<sub>3</sub> @ 100 ppm), S<sub>4</sub> (Cow urine @ 25 %), S<sub>5</sub> (Cow urine @ 50 %) and S<sub>6</sub> (Cow dung slurry; 1:1 w/w) and four levels of different growing media viz., G<sub>1</sub> (Sand + Vermicompost), G<sub>2</sub> (Sand + FYM), G<sub>3</sub> (Sand + Vermicompost + Cocopeat) and G<sub>4</sub> (Sand + FYM + Cocopeat) which were used in 1:1 proportion on volume basis. The seeds of acid lime were soaked before sowing in different seed soaking treatments for 12 hrs in beaker. The seeds were dried for 10 minutes in shade after soaking. The dried seeds were immediately sown in the polythene bags at 1.2 cm depth. The polythene bags were watered by water cane.

The days taken for germination of 50 % seeds from the date of sowing were observed daily. The number of days was counted from the date of initiation of germination up to the date of germination of 50 % seeds. Germination percentage was calculated by dividing the total number of seeds germinated by the total number of seeds sown and multiplying it by 100. Observations were recorded with respect to seedling height, stem diameter and number of leaves at 90, 120 and 150 days after sowing. Seedling height was measured from base of seedling to highest tip of plant. Stem diameter was measured 1 cm above from the base of the stem using vernier caliper. Fresh and dry weight of shoot as well as fresh and dry weight of root were measured by destructive method of uprooting the plant and taking measurement by standard method.

Germination percentage was recorded by using following formula:

$$\text{Germination (\%)} = \frac{\text{Number of germinated seeds}}{\text{Total number of seedlings}} \times 100$$

Survival per cent was recorded by using following formula:

$$\text{Survival (\%)} = \frac{\text{Total number of survived seedlings}}{\text{Total number of seedlings}} \times 100$$

The recorded data were analyzed statistically using various techniques as described by Panse and Sukhatme (1985). The treatment means were compared with C.D. at 5 per cent level.

## Results and Discussion

As per the results, the significant different was observed between the treatment on seed germination and seedling growth attributes.

### Germination parameters

The germination parameters were significantly influenced by application of different seed soaking treatments and growing media combinations. The minimum number of days taken to 50 % germination (16.09) as well as maximum germination percentage at 30 DAS (87.50 %) were recorded with treatment S<sub>3</sub> (GA<sub>3</sub> @ 100 ppm). The minimum number of days taken to 50 % germination (16.49) as well as maximum germination percentage at 30 DAS (85.56 %) were observed when seeds of acid lime were sown in growing media G<sub>3</sub> [Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)]. Interaction of S<sub>3</sub>G<sub>3</sub> [GA<sub>3</sub> @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded minimum number of days taken to 50 % germination (15.00) as well as maximum germination percentage (93.33 %) at 30 DAS.

This might be due to the promising effect of GA<sub>3</sub> on seed germination owing to its participation in the activity of alpha-amylase which catalyzed the starch conversion into simple carbohydrates and chemical energy was liberated used in the activation of embryo and as growing media have appropriate cation exchange capacity for retention of nutrients and have the properties of good water holding capacity as well as being sufficient porous for permitting adequate moisture and exchange of gases between media and embryo as reported by Ramteke and others(2015) in papaya. Similar results were also found by Parasana and others (2013) in mango and Khatana and others (2015) in Kagzi Lime.




Fig. 1: Effect of seed soaking and growing media on number of days taken to 50 % germination of acid lime

### Growth parameters

The growth parameters were significantly influenced by application of different seed soaking treatments and growing media combinations. The various growth parameters such as seedling height, stem diameter, number of leaves, fresh weight of shoot, dry weight of shoot, fresh weight of root, dry weight of root and survival percentage of seedling were found significant among different treatments.

### Height of seedling (cm) at 90, 120 and 150 DAS

Among seed soaking treatments, maximum height of seedling (16.68 cm, 23.40 cm and 27.81 cm) at 90, 120 and 150 DAS, respectively was observed in S<sub>3</sub> (GA<sub>3</sub>@ 100 ppm). Among growing media, maximum height of seedling (15.79 cm, 22.19 cm and 26.76 cm) at 90, 120 and 150 DAS, respectively was observed in G<sub>3</sub> [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S<sub>3</sub>G<sub>3</sub> [GA<sub>3</sub> @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum seedling height (18.30 cm, 26.13 cm and 30.50 cm) at 90, 120 and 150 DAS, respectively.

**Table 1:** Effect of seed soaking and growing media on number of days taken to 50 % germination, germination percentage at 30 DAS and seedling height (cm) of acid lime seedling

Treatments	Number of days taken to 50 % germination	Germination percentage at 30 DAS	Seedling height (cm)		
			90 DAS	120 DAS	150 DAS
<b>Factor A: Seed soaking (s)</b>					
S <sub>1</sub> : Water	20.47	65.83	9.46	14.69	19.20
S <sub>2</sub> : GA <sub>3</sub> @ 50 ppm	16.49	84.58	14.35	20.35	25.11
S <sub>3</sub> : GA <sub>3</sub> @ 100 ppm	16.09	87.50	16.68	23.40	27.81
S <sub>4</sub> : Cow urine @ 25 %	19.16	71.67	10.98	16.49	21.31
S <sub>5</sub> : Cow urine @ 50 %	18.86	73.33	11.35	17.12	21.72
S <sub>6</sub> : Cow dung slurry (1:1 w/w)	17.83	75.00	12.16	17.94	22.67
S.Em±	0.25	1.09	0.16	0.23	0.31
C.D. at 5 %	0.70	3.10	0.46	0.65	0.89
<b>Factor-B: Growing media (g)</b>					
G <sub>1</sub> : Sand + Vermicompost (1:1 v/v)	17.94	74.72	11.94	17.52	22.30
G <sub>2</sub> : Sand + FYM (1:1 v/v)	20.41	67.78	10.04	15.51	20.12
G <sub>3</sub> : Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)	16.49	85.56	15.79	22.19	26.76
G <sub>4</sub> : Sand + FYM + Cocopeat (1:1:1 v/v/v)	17.76	77.22	12.22	18.11	22.69
S.Em±	0.20	0.89	0.13	0.19	0.25
C.D. at 5 %	0.57	2.53	0.38	0.53	0.73
<b>Interaction (S x G)</b>					
S.Em±	0.49	2.18	0.33	0.45	0.62
C.D. at 5 %	1.41	6.20	0.93	1.29	1.78
C.V. %	4.72	4.94	4.51	4.30	4.71

This might due to the fact that the effect of gibberellic acid in increasing the osmotic uptake of nutrients, and thereby causing cell multiplication and cell elongation reflects in greater intermodal length, ultimately resulting in increase in plant height as reported by Deb and others (2014) in papaya and conducive effect of the media composition have low bulk density and higher water holding capacity resulting in better drainage of water and aeration which might have sustained good root and shoot growth leading to increase in seedling height (Abirami and others 2011 in nutmeg). Synergistic effect of both factors which help in increasing seedling height.

### **Stem diameter (mm) at 90, 120 and 150 DAS**

Among seed soaking treatments, maximum stem diameter (16.68 cm, 23.40 cm and 27.81 cm) at 90, 120 and 150 DAS, respectively was observed in S<sub>3</sub> (GA<sub>3</sub> @ 100 ppm). Among growing media, maximum stem diameter (15.79 cm, 22.19 cm and 26.76 cm) at 90, 120 and 150 DAS, respectively was observed in G<sub>3</sub> [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S<sub>3</sub>G<sub>3</sub> [GA<sub>3</sub> @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum stem diameter (2.09 mm, 2.90 mm and 3.49 mm) at 90, 120 and 150 DAS, respectively.

This might be due to GA<sub>3</sub> increases somatic uptake of nutrients causing cell multiplication, cell expansion and elongation in the cambium tissue thus it increases stem diameter of seedling (Feucht and Watson, 1958 in pea) and synergistic combinations of these growing media attributed proper aeration, high water holding capacity and better nutrient availability that enhance the physical condition of growing media leading to higher production of photosynthetically functional leaves in this treatment resulting in better diameter of seedling (Borah and others 1994 in silk cotton). These findings are in close accordance with the result of Ramteke and others (2015) in papaya, Rahangdale (2019) in custard apple.

### **Number of leaves at 90, 120 and 150 DAS**

Among seed soaking treatments, maximum number of leaves (17.82, 24.10 and 28.78) at 90, 120 and 150 DAS, respectively was observed in S<sub>3</sub> (GA<sub>3</sub> @ 100 ppm). Among growing media, maximum number of leaves (16.94, 23.49 and 27.66) at 90, 120 and 150 DAS, respectively was observed in G<sub>3</sub> [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S<sub>3</sub>G<sub>3</sub> [GA<sub>3</sub> @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum number of leaves (19.87, 27.07 and 32.27) at 90, 120 and 150 DAS, respectively.

This might be due to the activity of GA<sub>3</sub> at apical meristem resulting in more synthesis of nucleoprotein which is responsible for increasing leaf number and expansion and a appropriate media mixture provides better root environment to the plant leading to better nutrient availability to the photo synthetically functional leaves that ultimately utilized for more number of leaves (Ramteke and others 2015). Similar findings were also reported by Meena and others (2012) and Anjanawe and others (2013) in Papaya and Parasana and others (2014) in mango.

### **Fresh and dry weight of shoot (gm) at 150 DAS**

Among seed soaking treatments, maximum fresh and dry weight of shoot (6.90 g and 2.60 g, respectively) at 150 DAS was observed in S<sub>3</sub> (GA<sub>3</sub> @ 100 ppm). Among growing

media, maximum fresh and dry weight of shoot (6.60 g and 2.57 g, respectively) at 150 DAS was observed in G<sub>3</sub> [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S<sub>3</sub>G<sub>3</sub> [GA<sub>3</sub> @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum fresh and dry weight of shoot (7.57 g and 3.05 g, respectively) at 150 DAS.

This might be due to the combination containing many macro and micro nutrients, humic acid, which maintaining proper aeration and porosity and GA<sub>3</sub>, which increases cell division and uptake of nutrients, thus increasing the growth of seedling. The increase in fresh weight and dry weight might be because the combination of GA<sub>3</sub> and media increases the water and nutrient transportation to aerial parts, which leads to production of photosynthetic product and translocation of various plant parts, resulting in a higher fresh and dry weight of shoot. Similar findings were obtained by Kumawat and others (2014) and Dayeshwari and others (2017) in papaya.

**Table 2:** Effect of seed soaking and growing media on seedling diameter (cm) and number of leaves of acid lime seedling

Treatments	Seedling diameter (cm)			Number of leaves		
	90 DAS	120 DAS	150 DAS	90 DAS	120 DAS	150 DAS
<b>Factor A: Seed soaking (s)</b>						
S <sub>1</sub> : Water	1.44	1.87	2.27	10.15	15.52	19.42
S <sub>2</sub> : GA <sub>3</sub> @ 50 ppm	1.80	2.45	2.79	15.63	21.58	25.83
S <sub>3</sub> : GA <sub>3</sub> @ 100 ppm	1.94	2.66	3.11	17.82	24.10	28.78
S <sub>4</sub> : Cow urine @ 25 %	1.53	2.03	2.40	12.18	17.55	21.87
S <sub>5</sub> : Cow urine @ 50 %	1.60	2.10	2.44	12.63	18.07	22.45
S <sub>6</sub> : Cow dung slurry (1:1 w/w)	1.66	2.18	2.56	13.43	19.13	23.33
S.Em±	0.02	0.03	0.03	0.19	0.20	0.24
C.D. at 5 %	0.06	0.08	0.09	0.54	0.56	0.67
<b>Factor-B: Growing media (g)</b>						
G <sub>1</sub> : Sand + Vermicompost (1:1 v/v)	1.63	2.14	2.48	13.08	18.52	22.94
G <sub>2</sub> : Sand + FYM (1:1 v/v)	1.47	1.95	2.33	11.20	16.24	20.41
G <sub>3</sub> : Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)	1.90	2.57	3.02	16.94	23.49	27.66
G <sub>4</sub> : Sand + FYM + Cocopeat (1:1:1 v/v/v)	1.64	2.20	2.55	13.34	19.04	23.44
S.Em±	0.02	0.02	0.02	0.16	0.16	0.19
C.D. at 5 %	0.05	0.07	0.07	0.44	0.46	0.55
<b>Interaction (S x G)</b>						
S.Em±	0.04	0.06	0.06	0.38	0.40	0.47
C.D. at 5 %	0.12	0.16	0.18	1.08	1.13	1.34
C.V. %	4.35	4.50	4.14	4.84	3.55	3.45




Fig. 2: Effect of seed soaking and growing media on stem diameter (mm) at 90 DAS, 120 DAS and 150 DAS of acid lime

#### Fresh and dry weight of root (gm) at 150 DAS

Among seed soaking treatments, maximum fresh and dry weight of root (1.90 g and 0.62 g, respectively) at 150 DAS was observed in S<sub>3</sub> (GA<sub>3</sub> @ 100 ppm). Among growing media, maximum fresh and dry weight of root (1.83 g and 0.59 g, respectively) at 150 DAS was observed in G<sub>3</sub> [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S<sub>3</sub>G<sub>3</sub> [GA<sub>3</sub> @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum fresh and dry weight of root (2.59 g and 0.81 g, respectively) at 150 DAS.

The increased root growth is attributed to the fact that the combination of GA<sub>3</sub> and media provides a better root environment for the plant, resulting in better nutrient availability to the photosynthetically functional leaves and it also increases the translocation of food into the root zone, which leads to increase in fresh weight of root thereby increased dry weight of root. A similar result was recorded by Ramteke and others (2015) in papaya, Kaur (2017) in mango, Parab and others (2017) in papaya.

**Table 3:** Effect of seed soaking and growing media on fresh weight of shoot, dry weight of shoot, fresh weight of root, dry weight of root and survival percentage of acid lime seedling

Treatments	Fresh weight of shoot (g)	Dry weight of shoot (g)	Fresh weight of root (g)	Dry weight of root (g)	Survival percentage
<b>Factor A: Seed soaking (s)</b>					
S <sub>1</sub> : Water	4.33	1.54	0.84	0.21	62.50
S <sub>2</sub> : GA <sub>3</sub> @ 50 ppm	6.02	2.33	1.61	0.52	80.83
S <sub>3</sub> : GA <sub>3</sub> @ 100 ppm	6.90	2.60	1.90	0.62	88.75
S <sub>4</sub> : Cow urine @ 25 %	4.92	1.83	1.00	0.29	69.58
S <sub>5</sub> : Cow urine @ 50 %	5.14	1.99	1.06	0.31	71.67
S <sub>6</sub> : Cow dung slurry (1:1 w/w)	5.22	2.02	1.18	0.36	73.75
S.Em±	0.06	0.02	0.01	0.01	0.95
C.D. at 5 %	0.18	0.06	0.04	0.01	2.70
<b>Factor-B: Growing media (g)</b>					
G <sub>1</sub> : Sand + Vermicompost (1:1 v/v)	5.24	2.01	1.14	0.34	73.33
G <sub>2</sub> : Sand + FYM (1:1 v/v)	4.59	1.59	0.91	0.25	65.00
G <sub>3</sub> : Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)	6.60	2.57	1.83	0.59	86.11
G <sub>4</sub> : Sand + FYM + Cocopeat (1:1:1 v/v/v)	5.26	2.04	1.19	0.37	73.61
S.Em±	0.05	0.02	0.01	0.00	0.77
C.D. at 5 %	0.15	0.05	0.03	0.01	2.20
<b>Interaction (S x G)</b>					
S.Em±	0.13	0.04	0.03	0.01	1.89
C.D. at 5 %	0.36	0.12	0.08	0.03	5.39
C.V. %	4.10	3.48	3.96	4.35	4.40

### Survival percentage of seedling at 150 DAS

Among seed soaking treatments, maximum survival percentage of seedling (88.75 %) at 150 DAS was observed in S<sub>3</sub> (GA<sub>3</sub> @ 100 ppm). Among growing media, maximum survival percentage (86.11 %) at 150 DAS, respectively was observed in G<sub>3</sub> [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S<sub>3</sub>G<sub>3</sub> [GA<sub>3</sub> @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum survival percentage of seedling (96.67 %) at 150 DAS.

This may be due to GA<sub>3</sub> helps in cell expansion and its elongation resulting better root and shoot growth, which supports and encourage better survival of the seedlings (Rahangdale 2019) and media containing vermicompost and cocopeat as most of the constituents provided a start for establishment of seedlings which further got supplemented by PGPR's. Good physical and biological conditions in media combination had positive effect on root and shoot growth which also helps in better survival. Similar results were obtained by Ramteke and

others (2015) for papaya. These findings are in agreement with the results obtained by Gupta (1989) and Khatana and others (2015) in Kagzi Lime.

### Conclusion

From the present investigation, it can be concluded that seed soaking with GA<sub>3</sub> 100 ppm and growing media Sand + Vermicompost + Cocopeat (1:1:1) significantly influenced the seed germination, seedling growth and survivability of acid lime. The treatment combination S<sub>3</sub>G<sub>3</sub> [GA<sub>3</sub> @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] was found superior and most effective for better physiological development of acid lime seedlings over the rest of the treatment combinations.

### References

- Abirami K, Rema JR, Mathew PA and Sril V (2011) Response of nutmeg seeds to different nursery media. *Indian J Hort* 67(4):584-586
- Anjanawe SR, Kanpure RN, Kachouli BK and Mandloi DS (2013) Effect of plant growth regulators and growth media on seed germination and growth vigour of papaya. *Ann Plant Soil Res* 15(1):31-34
- Anonymous, (2020-21) 2<sup>nd</sup> Advance Estimates, National Horticulture Database. National Horticulture Board, Ministry of Agriculture and Farmers Welfare. Horticulture Statistics Division, Department of Agriculture, Cooperation and Farmers Welfare, Government of India, New Delhi
- Borah AS, Ray AN, Bhat AK, Maheswarappa R, Subramanian HP, Dileep PM, Sudhakara K, Santhoshkumar AV, Nazeema KK and Ashokan PK (1994) Effect of seed size, rooting medium and fertilizers on the growth of seedlings of silk cotton (*Ceiba pentandra* Linn). *Indian J For* 17(4):293-300
- Cheema GS, Bhat SS and Naik KC (1954) Commercial fruits of India with special reference to Western India. Macmillan and Co, London, pp 153-156
- Dayeshwari D, Rayaprolu S and Jone A (2017) Effect of potting media on seed germination, seedling growth and vigour in papaya cv. CO-8. *Int J Pure Appl Biosci* 3(5):505-512
- Deb P, Das A, Ghosh SK and Suresh CP (2014) Improvement of seed germination and seedling growth of papaya (*Carica papaya* L.) through different pre-sowing seed treatments. *Acta Hort* 8(51):313-316
- Feucht JR and Watson DP (1958) The effect of gibberellins on internal tissues of pea (*Phaseolus vulgaris* L.). *Am J Bot* 45:520-522
- Gupta OP (1989) Effect of gibberellic acid on seed germination in acid lime (*Citrus aurantifolia* Swingle) cv. Kagzi Lime. *Progress Hort* 21(3-4):246-248.
- Kaur S (2017) Effect of growing mixtures on seed germination and seedling growth of different mango (*Mangifera indica* L.) cultivars under submountainous conditions of Punjab. *Chem Sci Rev Lett* 6(23):1599-1603
- Khan MM, Usman M, Waseem R and Ali MA (2002) Role of gibberellic acid (GA<sub>3</sub>) on citrus seed germination and study of some morphological characteristics. *Pak J Agric Sci* 39(2):113-118
- Khatana KJ, Jadav RG and Nehete DS (2015) Influence of GA<sub>3</sub> on germination and growth of acid lime cv. Kagzi Lime seed (*Citrus aurantifolia* Swingle) under field as well as net house conditions. *J Asian Hort* 10(1):11-16

- Kumawat R, Maji S, Govind and Meena DC (2014) Studies on seed germination and seedling growth of papaya (*Carica papaya* L.) cv. Coorg Honey Dew as influenced by media and chemicals. *J Crop Weed* 10(2):281-286
- Meena RR, Jain MC and Mukerjee S (2012) Effect of pre-sowing dip treatment with gibberellic acid on germination and survivability of papaya. *Ann Plant Soil Res* 5(1):120-121
- Panse VG and Sukhatme PV (1985) Statistical methods for agricultural workers. 4<sup>th</sup> Edition., ICAR, New Delhi, pp 87-89
- Parab AM, Mathad TC and Malshe KV (2017) Effect of pre-soaking chemicals on germination and subsequent seedling growth of papaya (*Carica papaya* L.) cv. Solo. *Int J Chem Stud* 5(4):1812-1816
- Parasana JS, Leua HN and Ray NR (2013) Effect of different growing medias mixture on germination and seedling growth of mango (*Mangifera indica* L.) cultivars under net house condition. *Bioscan* 8(3):897-900
- Rahangdale P (2019) Effect of GA<sub>3</sub> and date of sowing on seed germination, growth and survival of custard apple (*Annona squamosa* L.) seedlings. Thesis M.Sc. (Horti.), submitted to JNKV, Jabalpur
- Raj A, Jhariya M and Toppo P (2014) Cow dung for eco-friendly and sustainable productive farming. *Int J Sci Res* 3(8):201-202
- Ramteke V, Paithankar DH, Ningot EP and Kurrey VK (2015) Effect of GA<sub>3</sub> and propagation media on germination, growth and vigour of papaya cv. Coorg Honey Dew. *The Bioscan* 10(3):1011-1016