

Original Research Article

Effects of seed soaking and growing media on germination and survivability of acid lime (*Citrus aurantifolia* Swingle)

Abstract

The experiments was laid out in a Completely Randomized Design with Factorial concept by three repetitions. The seeds of acid lime were treated with different seed soaking viz., water, GA₃ @ 50 ppm, GA₃ @ 100 ppm, cow urine @ 25 %, cow urine @ 50 % and cow dung slurry (1:1 w/w) and sowing in different growing Media viz., sand, vermicompost, FYM and coco peat. The result indicated that GA₃ @ 100 ppm is found to be the most effective for better germination parameters, growth parameters and survival percentage of acid lime seedlings. Among the different growing media, Sand + Vermicompost + Cocopeat (1:1:1 v/v/v) recorded higher values for germination parameters, growth parameters and survival percentage of acid lime seedlings. The treatment combination of GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v) showed better germination parameters, growth parameters and survival percentage. Therefore, the combination of GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v) was found most suitable for better growth of acid lime seedling.

Keywords: Acid lime, seed soaking, growing media, germination, survivability

Introduction

Citrus is one of the most important fruit crops of tropical and subtropical regions. It occupies 3rd rank after mango and banana in India. It is grown in over 100 countries and often regarded as golden fruit. The total area under cultivation of acid lime & lemon in India is 316 thousand hectares with production of around 3628 thousand MT (Anonymous, 2020-21). In Gujarat, area under citrus cultivation is 48503 hectares with 625833 MT production (Anonymous, 2020-21). It is mainly grown in Ahmedabad, Kheda, Mehsana, Bhavnagar, Gandhinagar and Junagadh districts of Gujarat.

Propagation of acid lime is generally done by grafting, budding, air layering and by seed. The commercial practice of propagation is by seed (Khatana and others 2015). It produces true to type, because of high degree (39-60 %) of nucellar embryony. The resultant seedlings are known to be free from tristeza virus and thus perform better. In acid lime,

Comment [N1]: Begin the abstract with a brief introduction which describes the study's rationale. Also mention the aim of the study, so that your reader is well guided. Also, use single line space for the abstract. Use capital letters for writing the abstract.

Comment [N2]: Close up the space.

Comment [N3]: as

Comment [N4]: with three replicates

Comment [N5]: samples:

Comment [N6]: Write in full the first time.

Comment [N7]: sowed

Comment [N8]: :

Comment [N9]: Write in full the first time.

Comment [N10]: was

Comment [N11]: Mention the treatment combinations before the results

Comment [N12]: In conclusion

Comment [N13]: Use capital letters for writing the keywords.

Comment [N14]: Use capital letters for writing the introduction.

Comment [N15]: Close up the space.

Comment [N16]: Include the source.

Comment [N17]: Include the source.

Comment [N18]: Include the source.

Comment [N19]: and

Comment [N20]: Close up the space.

Comment [N21]: Close up the space.

Comment [N22]: Include the source.

Comment [N23]: Close up the space.

Comment [N24]: Include the source.

Comment [N25]: et al.,

Comment [N26]: Include the source.

Comment [N27]: Include the source.

germination percentage is low, and it takes about 3 weeks to germinate (Cheema and others 1954). Some of the problems faced by acid lime growers are slow, erratic and incomplete germination with high initial seedling mortality. The possible reason of slow germination is presence of the growth inhibitors and physical resistance of seed coat to radical protrusion (Khan and others 2002). The growth of acid lime seedling is very slow in nursery as well as in the field. The increasing germination percentage and producing healthier seedling are major challenges for farmers.

So, the seeds of acid lime cv. Kagzi Lime are soaked with plant growth regulator such as gibberellic acid (GA₃) as well as with organic substance such as cow urine, cow dung slurry and water for better seed germination. GA₃ controls mobilization of starch which acts as a respiratory substrate leading to immediate enhancement in cell elongation. Gibberellins also help in enhancing the availability of reserved mineral elements which promote the germination process. Cow urine contains iron, urea, uric acid, estrogens and progesterone which affect the inhibitory responses to seed germination, shoot growth and seedling vigour. Cow dung slurry also contains some growth promoting substances (auxin), N, P, K, micronutrients and biodigestible enzymes which have been cause for softening of seed coat and thereby enhances seed germination and growth of seedlings (Raj and others 2014). Water soaking of seeds is done to modify hard seed coats, remove inhibitors, soften seeds and reduce the time of germination.

Media not only act as a growing place but also as a source of nutrient for plant growth. It is a substrate that provides the required elements and physical support to the growing plants. Vermicompost provides sufficient levels of oxygen to roots, adequate storage of water and nutrient for the plants. FYM is having good water holding capacity as well as sufficient porosity. Cocopeat provides excellent pore space (25-30 %) and fine structure required for proper growth. It is a rich source of nutrients and can easily mixed with other growing media used in seed germination and growth of seedling. Sand is used as a rooting media or adding the coarser texture needed to induce proper drainage and aeration. Keeping in view above, the aims of the study was to identify suitable treatment combinations for better seed germination and seedling growth of acid lime.

Materials and Methods

The experiment was carried out under a shade net house at College Farm, College of Horticulture, Sardarkrushinagar Dantiwada Agricultural University, Jagudan, Dist. Mehsana, Gujarat during July, 2021 to December, 2021. The experiment was laid out by Completely Randomized Design with Factorial concept (FCRD), which included 24 treatments with 3

Comment [N28]: Remove this information because it is old and out dated.

Comment [N29R28]:

Comment [N30]: Include the source.

Comment [N31]: *et al.*,

Comment [N32]: Cover up the space.

Comment [N33]: Include the source.

Comment [N34]: Include the source.

Comment [N35]: Include the source.

Comment [N36]: Include the source.

Comment [N37]:

Comment [N38]: *et al.*,

Comment [N39]: Include the source.

Comment [N40]: Cover up the space.

Comment [N41]: Include the source.

Comment [N42]: Include the source.

Comment [N43]: Include the source.

Comment [N44]: enough

Comment [N45]: Include the source.

Comment [N46]: Include the source.

... [1]

Comment [N47]: Include the source.

... [2]

Comment [N48]: Include the source.

... [3]

Comment [N49]: aim

Comment [N50]: The aim was not well stated.

Comment [N51]: Use capital letters for writing the materials and methos.

Comment [N52]: Cover up the space.

Comment [N53]: Write in full.

replications, there were two factors which comprised six levels of seed soaking viz., S₁ (Water), S₂ (GA₃ @ 50 ppm), S₃ (GA₃ @ 100 ppm), S₄ (Cow urine @ 25 %), S₅ (Cow urine @ 50 %) and S₆ (Cow dung slurry; 1:1 w/w) and four levels of different growing media viz., G₁ (Sand + Vermicompost), G₂ (Sand + FYM), G₃ (Sand + Vermicompost + Cocopeat) and G₄ (Sand + FYM + Cocopeat) which were used in 1:1 proportion on volume basis. The seeds of acid lime were soaked before sowing in different seed soaking treatments for 12 hrs in beaker. The seeds were dried for 10 minutes in shade after soaking. The dried seeds were immediately sown in the polythene bags at 1.2 cm depth. The polythene bags were watered by water can.

Comment [N54]: replicates

Comment [N55]: :

The days taken for germination of 50 % seeds from the date of sowing were observed daily. The number of days was counted from the date of initiation of germination up to the date of germination of 50 % seeds. Germination percentage was calculated by dividing the total number of seeds germinated by the total number of seeds sown and multiplying it by 100. Observations were recorded with respect to seedling height, stem diameter and number of leaves at 90, 120 and 150 days after sowing. Seedling height was measured from base of seedling to highest tip of plant. Stem diameter was measured 1 cm above from the base of the stem using Vernier caliper. Fresh and dry weight of shoot as well as fresh and dry weight of root were measured by destructive method of uprooting the plant and taking measurement by standard method.

Comment [N56]: artificially

Comment [N57]: Cover up the space.

Germination percentage was recorded by using following formula:

$$\text{Germination (\%)} = \frac{\text{Number of germinated seeds}}{\text{Total number of seedlings}} \times 100$$

Survival per cent was recorded by using following formula:

$$\text{Survival (\%)} = \frac{\text{Total number of survived seedlings}}{\text{Total number of seedlings}} \times 100$$

DATA ANALYSIS

The recorded data were analyzed statistically using various techniques as described by Panse and Sukhatme (1985). The treatment means were compared with C.D. at 5 per cent level.

Comment [N58]: Cover up the space.

Results and Discussion

Comment [N59]: Use capital letters for writing these.

As per the results, the significant different was observed between the treatment on seed germination and seedling growth attributes.

Comment [N60]: Cover up the space.


Germination parameters

The germination parameters were significantly influenced by application of different seed soaking treatments and growing media combinations as shown in Figures 1-2. In Figure 1, the minimum number of days taken to 50 % germination (16.09) as well as maximum germination percentage at 30 DAS (87.50 %) were recorded with treatment S₃ (GA₃ @ 100 ppm). The minimum number of days taken to 50 % germination (16.49) as well as maximum germination percentage at 30 DAS (85.56 %) were observed when seeds of acid lime were sown in growing media G₃ [Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)]. Interaction of S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded minimum number of days taken to 50 % germination (15.00) as well as maximum germination percentage (93.33 %) at 30 DAS.

Comment [N61]: Cover up the space.

This might be due to the promising effect of GA₃ on seed germination owing to its participation in the activity of alpha-amylase which catalyzed the starch conversion into simple carbohydrates and chemical energy was liberated used in the activation of embryo and as growing media have appropriate cation exchange capacity for retention of nutrients and have the properties of good water holding capacity as well as being sufficient porous for permitting adequate moisture and exchange of gases between media and embryo as reported by Ramteke and others (2015) in papaya. Similar results were also found by Parasana and others (2013) in mango and Khatana and others (2015) in Kagzi Lime.

Comment [N62]: Cover up the space.



Comment [N63]: et al.,

Comment [N64]: The sentence is too long. So, split into parts

Comment [N65]: et al.,

Comment [N66]: et al.,

Fig. 1: Effect of seed soaking and growing media on number of days taken to 50 % germination of acid lime

Comment [N67]: Figure 1

Comment [N68]: Bold the figure title.

Growth parameters

The growth parameters were significantly influenced by application of different seed soaking treatments and growing media combinations as shown in Table 1-3. The various growth parameters such as seedling height, stem diameter, number of leaves, fresh weight of

Comment [N69]: Cover up the space.

shoot, dry weight of shoot, fresh weight of root, dry weight of root and survival percentage of seedling were found significant among different treatments. The **height of seedling (cm) at 90, 120 and 150 DAS** as shown in Table 1.

Among seed soaking treatments, maximum height of seedling (16.68 cm, 23.40 cm and 27.81 cm) at 90, 120 and 150 DAS, respectively was observed in S₃ (GA₃@ 100 ppm). Among growing media, maximum height of seedling (15.79 cm, 22.19 cm and 26.76 cm) at 90, 120 and 150 DAS, respectively was observed in G₃ [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S₃G₃ [GA₃ @ 0 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum seedling height (18.30 cm, 26.13 cm and 30.50 cm) at 90, 120 and 150 DAS, respectively.

Table 1: Effects of seed soaking and growing media on number of days taken to 50 % germination, germination percentage at 30 DAS and seedling height (cm) of acid lime seedling

Treatments	Number of days taken to 50 % germination	Germination percentage at 30 DAS	Seedling height (cm)		
			90 DAS	120 DAS	150 DAS
Factor A: Seed soaking (s)					
S ₁ : Water	20.47	65.83	9.46	14.69	19.20
S ₂ : GA ₃ @ 50 ppm	16.49	84.58	14.35	20.35	25.11
S ₃ : GA ₃ @ 100 ppm	16.09	87.50	16.68	23.40	27.81
S ₄ : Cow urine @ 25 %	19.16	71.67	10.98	16.49	21.31
S ₅ : Cow urine @ 50 %	18.86	73.33	11.35	17.12	21.72
S ₆ : Cow dung slurry (1:1 w/w)	17.83	75.00	12.16	17.94	22.67
S.Em±	0.25	1.09	0.16	0.23	0.31
C.D. at 5 %	0.70	3.10	0.46	0.65	0.89
Factor-B: Growing media (g)					
G ₁ : Sand + Vermicompost (1:1 v/v)	17.94	74.72	11.94	17.52	22.30
G ₂ : Sand + FYM (1:1 v/v)	20.41	67.78	10.04	15.51	20.12
G ₃ : Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)	16.49	85.56	15.79	22.19	26.76
G ₄ : Sand + FYM + Cocopeat (1:1:1 v/v/v)	17.76	77.22	12.22	18.11	22.69
S.Em±	0.20	0.89	0.13	0.19	0.25
C.D. at 5 %	0.57	2.53	0.38	0.53	0.73
Interaction (S x G)					
S.Em±	0.49	2.18	0.33	0.45	0.62
C.D. at 5 %	1.41	6.20	0.93	1.29	1.78
C.V. %	4.72	4.94	4.51	4.30	4.71

This might due to the fact that the effect of gibberellic acid in increasing the osmotic uptake of nutrients, and thereby causing cell multiplication and cell elongation reflects in greater intermodal length, ultimately resulting in increase in plant height as reported by Deb and others (2014) in papaya and conducive effect of the media composition have low bulk density and higher water holding capacity resulting in better drainage of water and aeration

Comment [N70]: Unbold.

Comment [N71]: Cover up the space.

Comment [N72]: Close up the space.

Comment [N73]: Bold the Table title.

Comment [N74]: Cover up the space.

Comment [N75]: et al.,

which might have sustained good root and shoot growth leading to increase in seedling height (Abirami and others 2011 in nutmeg). Synergistic effect of both factors which help in increasing seedling height.

The Stem diameter (mm) at 90, 120 and 150 DAS as shown in Table 2.

Among seed soaking treatments, maximum stem diameter (16.68 cm, 23.40 cm and 27.81 cm) at 90, 120 and 150 DAS, respectively was observed in S₃ (GA₃ @ 100 ppm). Among growing media, maximum stem diameter (15.79 cm, 22.19 cm and 26.76 cm) at 90, 120 and 150 DAS, respectively was observed in G₃ [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum stem diameter (2.09 mm, 2.90 mm and 3.49 mm) at 90, 120 and 150 DAS, respectively.

This might be due to GA₃ increases somatic uptake of nutrients causing cell multiplication, cell expansion and elongation in the cambium tissue thus it increases stem diameter of seedling (Feucht and Watson, 1958) in pea and synergistic combinations of these growing media attributed proper aeration, high water holding capacity and better nutrient availability that enhance the physical condition of growing media leading to higher production of photosynthetically functional leaves in this treatment resulting in better diameter of seedling (Borah and others (1994) in silk cotton). These findings are in close accordance with the result of Ramteke and others (2015) in papaya, Rahangdale (2019) in custard apple.

The number of leaves at 90, 120 and 150 DAS as shown in Table 2.

Among seed soaking treatments, maximum number of leaves (17.82, 24.10 and 28.78) at 90, 120 and 150 DAS, respectively was observed in S₃ (GA₃ @ 100 ppm). Among growing media, maximum number of leaves (16.94, 23.49 and 27.66) at 90, 120 and 150 DAS, respectively was observed in G₃ [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum number of leaves (19.87, 27.07 and 32.27) at 90, 120 and 150 DAS, respectively.

This might be due to the activity of GA₃ at apical meristem resulting in more synthesis of nucleoprotein which is responsible for increasing leaf number and expansion and a appropriate media mixture provides better root environment to the plant leading to better nutrient availability to the photo synthetically functional leaves that ultimately utilized for more number of leaves (Ramteke and others 2015). Similar findings were also reported by Meena and others (2012) and Anjanawe and others (2013) in Papaya and Parasana and others (2014) in mango.

Comment [N76]: *et al.,*

Comment [N77]: The sentence is too long. So, split into parts

Comment [N78]: Unbold.

Comment [N79]: Cover up the space.

Comment [N80]: Cover up the space.

Comment [N81]: Remove this information because it is old and outdated.

Comment [N82]: *et al.,*

Comment [N83]: Delete.

Comment [N84]: *et al.,*

Comment [N85]: The sentence is too long. So, split into parts

Comment [N86]: Unbold

Comment [N87]: Cover up the space.

Comment [N88]: Cover up the space.

Comment [N89]: *et al.,*

Comment [N90]: *et al.,*

Comment [N91]: *et al.,*

Comment [N92]: *et al.,*

... [4]

Comment [N93]: The sentence is too long. So, split into parts

... [5]

The fresh and dry weight of shoot (gm) at 150 DAS as shown in Table 3.

Among seed soaking treatments, maximum fresh and dry weight of shoot (6.90 g and 2.60 g, respectively) at 150 DAS was observed in S₃ (GA₃ @ 100 ppm). Among growing media, maximum fresh and dry weight of shoot (6.60 g and 2.57 g, respectively) at 150 DAS was observed in G₃ [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum fresh and dry weight of shoot (7.57 g and 3.05 g, respectively) at 150 DAS.

This might be due to the combination containing many macro and micronutrients, humid acid, which maintaining proper aeration and porosity and GA₃, which increases cell division and uptake of nutrients, thus increasing the growth of seedling. The increase in fresh weight and dry weight might be because the combination of GA₃ and media increases the water and nutrient transportation to aerial parts, which leads to production of photosynthetic product and translocation of various plant parts, resulting in a higher fresh and dry weight of shoot. Similar findings were obtained by Kumawat and others (2014) and Dayeshwari and others (2017) in papaya.

Table 2: Effects of seed soaking and growing media on seedling diameter (cm) and number of leaves of acid lime seedling

Treatments	Seedling diameter (cm)			Number of leaves		
	90 DAS	120 DAS	150 DAS	90 DAS	120 DAS	150 DAS
Factor A: Seed soaking (s)						
S ₁ : Water	1.44	1.87	2.27	10.15	15.52	19.42
S ₂ : GA ₃ @ 50 ppm	1.80	2.45	2.79	15.63	21.58	25.83
S ₃ : GA ₃ @ 100 ppm	1.94	2.66	3.11	17.82	24.10	28.78
S ₄ : Cow urine @ 25 %	1.53	2.03	2.40	12.18	17.55	21.87
S ₅ : Cow urine @ 50 %	1.60	2.10	2.44	12.63	18.07	22.45
S ₆ : Cow dung slurry (1:1 w/w)	1.66	2.18	2.56	13.43	19.13	23.33
S.Em±	0.02	0.03	0.03	0.19	0.20	0.24
C.D. at 5 %	0.06	0.08	0.09	0.54	0.56	0.67
Factor-B: Growing media (g)						
G ₁ : Sand + Vermicompost (1:1 v/v)	1.63	2.14	2.48	13.08	18.52	22.94
G ₂ : Sand + FYM (1:1 v/v)	1.47	1.95	2.33	11.20	16.24	20.41
G ₃ : Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)	1.90	2.57	3.02	16.94	23.49	27.66
G ₄ : Sand + FYM + Cocopeat (1:1:1 v/v/v)	1.64	2.20	2.55	13.34	19.04	23.44
S.Em±	0.02	0.02	0.02	0.16	0.16	0.19
C.D. at 5 %	0.05	0.07	0.07	0.44	0.46	0.55
Interaction (S x G)						
S.Em±	0.04	0.06	0.06	0.38	0.40	0.47
C.D. at 5 %	0.12	0.16	0.18	1.08	1.13	1.34
C.V. %	4.35	4.50	4.14	4.84	3.55	3.45

Comment [N94]: Unbold

Comment [N95]: Cover up the space.

Comment [N96]: Cover up the space.

Comment [N97]: et al.,

Comment [N98]: et al.,

Comment [N99]: This explanation is for Table 3, so move and place it directly under Figure 2.

Comment [N100]: Bold the Table title.




Fig. 2: Effect of seed soaking and growing media on stem diameter (mm) at 90 DAS, 120 DAS and 150 DAS of acid lime

The fresh and dry weight of root (gm) at 150 DAS as shown in Table 3.

Among seed soaking treatments, maximum fresh and dry weight of root (1.90 g and 0.62 g, respectively) at 150 DAS was observed in S₃ (GA₃ @ 100 ppm). Among growing media, maximum fresh and dry weight of root (1.83 g and 0.59 g, respectively) at 150 DAS was observed in G₃ [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum fresh and dry weight of root (2.59 g and 0.81 g, respectively) at 150 DAS.

The increased root growth is attributed to the fact that the combination of GA₃ and media provides a better root environment for the plant, resulting in better nutrient availability to the photo synthetically functional leaves and it also increases the translocation of food into the root zone, which leads to increase in fresh weight of root thereby increased dry weight of root. A similar result was recorded by Ramteke and others (2015) in papaya, Kaur (2017) in mango, Parab and others (2017) in papaya.

Comment [N101]: Figure

Comment [N102]: Bold the figure title.

Comment [N103]: Unbold

Comment [N104]: Cover up the space.

Comment [N105]: Cover up the space.

Comment [N106]: *et al.*,

Comment [N107]: *et al.*,

Table 3: Effects of seed soaking and growing media on fresh weight of shoot, dry weight of shoot, fresh weight of root, dry weight of root and survival percentage of acid lime seedling

Treatments	Fresh weight of shoot (g)	Dry weight of shoot (g)	Fresh weight of root (g)	Dry weight of root (g)	Survival percentage
Factor A: Seed soaking (s)					
S ₁ : Water	4.33	1.54	0.84	0.21	62.50
S ₂ : GA ₃ @ 50 ppm	6.02	2.33	1.61	0.52	80.83
S ₃ : GA ₃ @ 100 ppm	6.90	2.60	1.90	0.62	88.75
S ₄ : Cow urine @ 25 %	4.92	1.83	1.00	0.29	69.58
S ₅ : Cow urine @ 50 %	5.14	1.99	1.06	0.31	71.67
S ₆ : Cow dung slurry (1:1 w/w)	5.22	2.02	1.18	0.36	73.75
S.Em±	0.06	0.02	0.01	0.01	0.95
C.D. at 5 %	0.18	0.06	0.04	0.01	2.70
Factor-B: Growing media (g)					
G ₁ : Sand + Vermicompost (1:1 v/v)	5.24	2.01	1.14	0.34	73.33
G ₂ : Sand + FYM (1:1 v/v)	4.59	1.59	0.91	0.25	65.00
G ₃ : Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)	6.60	2.57	1.83	0.59	86.11
G ₄ : Sand + FYM + Cocopeat (1:1:1 v/v/v)	5.26	2.04	1.19	0.37	73.61
S.Em±	0.05	0.02	0.01	0.00	0.77
C.D. at 5 %	0.15	0.05	0.03	0.01	2.20
Interaction (S x G)					
S.Em±	0.13	0.04	0.03	0.01	1.89
C.D. at 5 %	0.36	0.12	0.08	0.03	5.39
C.V. %	4.10	3.48	3.96	4.35	4.40

Comment [N108]: Bold the Table title.

The Survival percentage of seedling at 150 DAS as shown in Table 3.

Comment [N109]: Unbold

Among seed soaking treatments, maximum survival percentage of seedling (88.75 %) at 150 DAS was observed in S₃ (GA₃ @ 100 ppm). Among growing media, maximum survival percentage (86.11 %) at 150 DAS, respectively was observed in G₃ [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum survival percentage of seedling (96.67 %) at 150 DAS.

Comment [N110]: Cover up the space.

This may be due to GA₃ helps in cell expansion and its elongation resulting better root and shoot growth, which supports and encourage better survival of the seedlings (Rahangdale 2019) and media containing vermicompost and cocopeat as most of the constituents provided a start for establishment of seedlings which further got supplemented by PGPR's. Good physical and biological conditions in media combination had positive effect on root and shoot growth which also helps in better survival. Similar results were obtained by Ramteke and

Comment [N111]: Cover up the space.

others (2015) for papaya. These findings are in agreement with the results obtained by Gupta (1989) and Khatana and others (2015) in Kagzi Lime.

Conclusion

From the present investigation, it can be concluded that seed soaking with GA₃ 100 ppm and growing media Sand + Vermicompost + Cocopeat (1:1:1) significantly influenced the seed germination, seedling growth and survivability of acid lime. The treatment combination S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] was found superior and most effective for better physiological development of acid lime seedlings over the rest of the treatment combinations.

References

- Abirami, K., Rema, J. R., Mathew, P. A. and Sril, V. (2011). Response of nutmeg seeds to different nursery media. *Indian J Hortic*, 67(4):584-586.
- Anjanawe, S. R., Kanpure, R. N., Kachouli, B. K. and Mandloi, D. S. (2013). Effects of plant growth regulators and growth media on seed germination and growth vigour of papaya. *Ann Plant Soil Res*, 15(1):31-34.
- Anonymous, (2020-21) 2nd Advance Estimates, National Horticulture Database. National Horticulture Board, Ministry of Agriculture and Farmers Welfare. Horticulture Statistics Division, Department of Agriculture, Cooperation and Farmers Welfare, Government of India, New Delhi
- Borah, A. S., Ray, A. N., Bhat, A. K., Maheswarappa, R., Subramanian, H. P., Dileep, P. M., Sudhakara, K., Santhoshkumar, A. V., Nazeema, K. K. and Ashokan, P. K. (1994). Effects of seed size, rooting medium and fertilizers on the growth of seedlings of silk cotton (*Ceiba pentandra* Linn). *Indian J For*, 17(4):293-300.
- Cheema, GS, Bhat SS and Naik KC (1954) Commercial fruits of India with special reference to Western India. Macmillan and Co, London, pp 153-156
- Dayeshwari, D., Rayaprolu, S. and Jone, A. (2017). Effects of potting media on seed germination, seedling growth and vigour in papaya cv. CO-8. *Int J Pure Appl Biosci*, 3(5):505-512.
- Deb, P., Das, A., Ghosh, S. K. and Suresh, C. P. (2014). Improvement of seed germination and seedling growth of papaya (*Carica papaya* L.) through different pre-sowing seed treatments. *Acta Hortic*, 8(51):313-316.
- Feucht, JR and Watson, DP (1958) The effect of gibberellins on internal tissues of pea (*Phaseolus vulgaris* L.). *Am J Bot* 45:520-522.
- Gupta, O. P. (1989). Effect of gibberellic acid on seed germination in acid lime (*Citrus aurantifolia* Swingle) cv. Kagzi Lime. *Progress Hortic*, 21(3-4):246-248.
- Kaur, S. (2017). Effects of growing mixtures on seed germination and seedling growth of different mango (*Mangifera indica* L.) cultivars under submountainous conditions of Punjab. *Chem Sci Rev Lett*, 6(23):1599-1603.
- Khan, M. M., Usman, M., Waseem, R. and Ali, M. A. (2002). Role of gibberellic acid (GA₃) on citrus seed germination and study of some morphological characteristics. *Pak J Agric Sci*, 39(2):113-118.
- Khatana, K. J., Jadav, R. G. and Nehete, D. S. (2015). Influence of GA₃ on germination and growth of acid lime cv. Kagzi Lime seed (*Citrus aurantifolia* Swingle) under field as well as net house conditions. *J Asian Hortic*, 10(1):11-16.

Comment [N112]: et al.,

Comment [N113]: et al.,

Comment [N114]: This explanation is for Table 3, so move and place it directly on top of Table 3.

Comment [N115]: Capitals

Comment [N116]: Cover up the space.

Comment [N117]: Capitals

Comment [N118]: Write in full and italicize.

Comment [N119]: Write in full and italicize.

Comment [N120]: Write in full and italicize.

Comment [N121]: Old remove.

Comment [N122]: Write in full and italicize.

Comment [N123]: Write in full and italicize.

Comment [N124]: Old so remove.

Comment [N125]: Write in full and italicize.

Comment [N126]: Write in full and italicize.

Comment [N127]: Write in full and italicize.

Comment [N128]: Write in full and italicize.

- Kumawat, R., Maji, S., Govind, and Meena, D. C. (2014). Studies on seed germination and seedling growth of papaya (*Carica papaya* L.) cv. Coorg Honey Dew as influenced by media and chemicals. *J Crop Weed*, 10(2):281-286.
- Meena, R. R., Jain, M. C. and Mukerjee, S. (2012). Effect of pre-sowing dip treatment with gibberellic acid on germination and survivability of papaya. *Ann Plant Soil Res*, 5(1):120-121.
- Panse, V. G. and Sukhatme, P. V. (1985). *Statistical methods for agricultural workers*, 4th Edition., ICAR, New Delhi, pp 87-89.
- Parab, A. M., Mathad, T. C. and Malshe, K. V. (2017). Effects of pre-soaking chemicals on germination and subsequent seedling growth of papaya (*Carica papaya* L.) cv. Solo. *Int J Chem Stud*, 5(4):1812-1816.
- Parasana, J. S., Leua, H. N. and Ray, N. R. (2013). Effect of different growing media mixture on germination and seedling growth of mango (*Mangifera indica* L.) cultivars under net house condition. *Bioscan*, 8(3):897-900.
- Rahangdale, P. (2019). Effects of GA₃ and date of sowing on seed germination, growth and survival of custard apple (*Annona squamosa* L.) seedlings. *Thesis M.Sc.* (Horti.), submitted to JNKV, Jabalpur
- Raj, A., Jhariya, M. and Toppo, P. (2014). Cow dung for eco-friendly and sustainable productive farming. *Int J Sci Res*, 3(8):201-202.
- Ramteke, V., Paithankar, D. H., Ningot, E. P. and Kurrey, V. K. (2015). Effects of GA₃ and propagation media on germination, growth and vigour of papaya cv. Coorg Honey Dew. *The Bioscan*, 10(3):1011-1016.

Comment [N129]: Write in full and italicize.

Comment [N130]: Write in full and italicize.

Comment [N131]: Italicize.

Comment [N132]: Write in full and italicize.

Comment [N133]: Delete s

Comment [N134]: Write in full and italicize.

Comment [N135]: Italicize.

Comment [N136]: Write in full and italicize.

11/29/2022 10:32:00 AM

11/29/2022 10:21:00 AM