

Original Research Article

Effects of seed soaking and growing media on germination and survivability of acid lime (*Citrus aurantifolia* Swingle)

ABSTRACT

Seed germination is the most important aspect for raising the nursery for successful seedling production of acid lime. The present study was carried out to explore the effect of different seed soaking treatment with growing media on seed germination and survivability of acid lime (*Citrus aurantifolia* Swingle) under net house condition. The experiments were laid out as a Completely Randomized Design with Factorial concept comprising twenty four treatment combinations with three replicates. The seeds of acid lime were treated with different seed soaking: water, GA₃ @ 50 ppm, GA₃ @ 100 ppm, cow urine @ 25 %, cow urine @ 50 % and cow dung slurry (1:1, cow dung:water w/w) and sowed in different growing media combinations: Sand + Vermicompost (1:1 v/v), Sand + Farmyard manure (1:1 v/v), Sand + Vermicompost + Cocopeat (1:1:1 v/v/v), Sand + Farmyard manure + Cocopeat (1:1:1 v/v/v). The result indicated that GA₃ @ 100 ppm was found to be the most effective for better germination parameters, growth parameters and survival percentage of acid lime seedlings. Among the different growing media, Sand + Vermicompost + Cocopeat (1:1:1 v/v/v) recorded higher values for germination parameters, growth parameters and survival percentage of acid lime seedlings. The treatment combination of GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v) showed better germination parameters, growth parameters and survival percentage. In conclusion, the combination of GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v) was found most suitable for better growth of acid lime seedling.

KEYWORDS: Acid lime, seed soaking, growing media, germination, survivability

INTRODUCTION

Citrus is one of the most important fruit crops of tropical and subtropical regions (Jaiswal and others 2018). It occupies 3rd rank after mango and banana in India (Dilip and others 2017). It is grown in over 100 countries and often regarded as golden fruit. The total area under cultivation of acid lime and lemon in India is 316 thousand hectares with production of around 3628 thousand MT (Anon., 2020-21). In Gujarat, area under citrus cultivation is 48503 hectares with 625833 MT productions covering districts of Ahmedabad, Kheda, Mehsana, Bhavnagar, Gandhinagar and Junagadh districts of Gujarat (Anon., 2020-21).

Propagation of acid lime is generally done by grafting, budding, air layering and by seed (Khatana and others 2015). The commercial practice of propagation is by seed (Khatana and others 2015). It produces true to type, because of high degree (39-60 %) of nucellar embryony (Jaiswal and others 2018). The resultant seedlings are known to be free from tristeza virus and thus perform better (Jaiswal and others 2018). Some of the problems faced by acid lime growers are slow, erratic and incomplete germination with high initial seedling

mortality (Khan and others 2002). The possible reason of slow germination is presence of the growth inhibitors and physical resistance of seed coat to radical protrusion (Elza, 1949). The growth of acid lime seedling is very slow in nursery as well as in the field. The increasing germination percentage and producing healthier seedling are major challenges for farmers.

So, the seeds of acid lime cv. Kagzi Lime are soaked with plant growth regulator such as gibberellic acid (GA₃) as well as with organic substance such as cow urine, cow dung slurry and water for better seed germination. GA₃ controls mobilization of starch which acts as a respiratory substrate leading to immediate enhancement in cell elongation (Tandon and others 2019). Gibberellins also help in enhancing the availability of reserved mineral elements which promote the germination process (Tandon and others 2019). Cow urine contains iron, urea, uric acid, estrogens and progesterone which affect the inhibitory responses to seed germination, shoot growth and seedling vigour (Tandon and others 2019). Cow dung slurry also contains some growth promoting substances (auxin), N, P, K, micronutrients and biodigestible enzymes which have been cause for softening of seed coat and thereby enhances seed germination and growth of seedlings (Raj and others 2014). Water soaking of seeds is done to modify hard seed coats, remove inhibitors, soften seeds and reduce the time of germination (Wagner-Junior and others 2005).

Media not only act as a growing place but also as a source of nutrient for plant growth (Suthar, 2019). It is a substrate that provides the required elements and physical support to the growing plants (Suthar, 2019). Vermicompost provides sufficient levels of oxygen to roots, adequate storage of water and nutrient for the plants (Atefe and others 2012). FYM is having good water holding capacity as well as enough porosity (Anjanawe and others 2013). Cocopeat provides excellent pore space (25-30 %) and fine structure required for proper growth (Desai and others 2017). It is a rich source of nutrients and can easily mixed with other growing media used in seed germination and growth of seedling (Desai and others 2017). Sand is used as a rooting media or adding the coarser texture needed to induce proper drainage and aeration (Yadav and others 2012).

Producing plants from seeds are the most important propagation method. Seedling propagation involves careful management of seeds, seed storage period, germination conditions and knowledge of requirements of seed for germination as well as the overall growth. Growing media and seed soaking both are an integral part for most of the

horticultural production system. Keeping in view above, the aims of the study was to identify suitable treatment combinations for better seed germination and seedling growth of acid lime.

MATERIALS AND METHODS

The experiment was carried out under a shade net house at College Farm, College of Horticulture, Sardarkrushinagar Dantiwada Agricultural University, Jagudan, District Mehsana, Gujarat during July, 2021 to December, 2021. The experiment was laid out by Completely Randomized Design with Factorial concept (FCRD), which included 24 treatments with 3 replicates, there were two factors which comprised six levels of seed soaking: S₁ (Water), S₂ (GA₃ @ 50 ppm), S₃ (GA₃ @ 100 ppm), S₄ (Cow urine @ 25 %), S₅ (Cow urine @ 50 %) and S₆ (Cow dung slurry; 1:1 w/w) and four levels of different growing media: G₁ (Sand + Vermicompost), G₂ (Sand + FYM), G₃ (Sand + Vermicompost + Cocopeat) and G₄ (Sand + FYM + Cocopeat) which were used in 1:1 proportion on volume basis. The seeds of acid lime were soaked before sowing in different seed soaking treatments for 12 hrs in beaker. The seeds were dried for 10 minutes in shade after soaking. The dried seeds were immediately sown in the polythene bags at 1.2 cm depth. The polythene bags were watered artificially.

The days taken for germination of 50 % seeds from the date of sowing were observed daily. The number of days was counted from the date of initiation of germination up to the date of germination of 50 % seeds. Germination percentage was calculated by dividing the total number of seeds germinated by the total number of seeds sown and multiplying it by 100. Observations were recorded with respect to seedling height, stem diameter and number of leaves at 90, 120 and 150 days after sowing. Seedling height was measured from base of seedling to highest tip of plant. Stem diameter was measured 1 cm above from the base of the stem using Vernier caliper. Fresh and dry weight of shoot as well as fresh and dry weight of root was measured by destructive method of uprooting the plant and taking measurement by standard method.

Germination percentage was recorded by using following formula:

$$\text{Germination (\%)} = \frac{\text{Number of germinated}}{\text{Total number of seedlings}} \times 100$$

Survival per cent was recorded by using following formula:

$$\frac{\text{Total number of survived seedlings}}{\text{Total number of seedlings}} \times 100$$

$$\text{Survival (\%)} = \frac{\quad}{\text{Total number of seedlings}}$$

DATA ANALYSIS

The recorded data were analyzed statistically using various techniques as described by Panse and Sukhatme (1985). The treatment means were compared with C.D. at 5 per cent level.

RESULTS AND DISCUSSION

As per the results, the significant different was observed between the treatment on seed germination and seedling growth attributes.

Germination parameters

The germination parameters were significantly influenced by application of different seed soaking treatments and growing media combinations as shown in Figures 1-2. In Figure 1, the minimum number of days taken to 50 % germination (16.09) as well as maximum germination percentage at 30 DAS (87.50 %) were recorded with treatment S₃ (GA₃ @ 100 ppm). The minimum number of days taken to 50 % germination (16.49) as well as maximum germination percentage at 30 DAS (85.56 %) were observed when seeds of acid lime were sown in growing media G₃ [Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)]. Interaction of S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded minimum number of days taken to 50 % germination (15.00) as well as maximum germination percentage (93.33 %) at 30 DAS.

This might be due to the promising effect of GA₃ on seed germination owing to its participation in the activity of alpha-amylase which catalyzed the starch conversion into simple carbohydrates and chemical energy was liberated used in the activation of embryo. Growing media have appropriate cation exchange capacity for retention of nutrients and have the properties of good water holding capacity as well as being sufficient porous for permitting adequate moisture and exchange of gases between media and embryo as reported by Ramteke and others (2015) in papaya. Similar results were also found by Parasana and others (2013) in mango and Khatana and others (2015) in Kagzi Lime.




Figure 1: Effect of seed soaking and growing media on number of days taken to 50 % germination of acid lime

Growth parameters

The growth parameters were significantly influenced by application of different seed soaking treatments and growing media combinations as shown in Table 1-3. The various growth parameters such as seedling height, stem diameter, number of leaves, fresh weight of shoot, dry weight of shoot, fresh weight of root, dry weight of root and survival percentage of seedling were found significant among different treatments.

The height of seedling (cm) at 90, 120 and 150 DAS as shown in Table 1.

Among seed soaking treatments, maximum height of seedling (16.68 cm, 23.40 cm and 27.81 cm) at 90, 120 and 150 DAS, respectively was observed in S₃ (GA₃@ 100 ppm). Among growing media, maximum height of seedling (15.79 cm, 22.19 cm and 26.76 cm) at 90, 120 and 150 DAS, respectively was observed in G₃ [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum seedling height (18.30 cm, 26.13 cm and 30.50 cm) at 90, 120 and 150 DAS, respectively.

This might be due to the fact that the effect of gibberellic acid in increasing the osmotic uptake of nutrients, and thereby causing cell multiplication and cell elongation reflects in greater intermodal length, ultimately resulting in increase in plant height as reported by Deb and others (2014) in papaya. Conducive effect of the media composition have low bulk density and higher water holding capacity resulting in better drainage of water and aeration which might have sustained good root and shoot growth leading to increase in seedling height (Abirami and others 2011 in nutmeg). Synergistic effect of both factors which help in increasing seedling height.

Table 1: Effect of seed soaking and growing media on number of days taken to 50 % germination, germination percentage at 30 DAS and seedling height (cm) of acid lime seedling

Treatments	Number of days taken to 50 % germination	Germination percentage at 30 DAS	Seedling height (cm)		
			90 DAS	120 DAS	150 DAS
Factor A: Seed soaking (s)					
S ₁ : Water	20.47	65.83	9.46	14.69	19.20
S ₂ : GA ₃ @ 50 ppm	16.49	84.58	14.35	20.35	25.11
S ₃ : GA ₃ @ 100 ppm	16.09	87.50	16.68	23.40	27.81
S ₄ : Cow urine @ 25 %	19.16	71.67	10.98	16.49	21.31
S ₅ : Cow urine @ 50 %	18.86	73.33	11.35	17.12	21.72
S ₆ : Cow dung slurry (1:1 w/w)	17.83	75.00	12.16	17.94	22.67
S.Em±	0.25	1.09	0.16	0.23	0.31
C.D. at 5 %	0.70	3.10	0.46	0.65	0.89
Factor-B: Growing media (g)					
G ₁ : Sand + Vermicompost (1:1 v/v)	17.94	74.72	11.94	17.52	22.30
G ₂ : Sand + FYM (1:1 v/v)	20.41	67.78	10.04	15.51	20.12
G ₃ : Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)	16.49	85.56	15.79	22.19	26.76
G ₄ : Sand + FYM + Cocopeat (1:1:1 v/v/v)	17.76	77.22	12.22	18.11	22.69
S.Em±	0.20	0.89	0.13	0.19	0.25
C.D. at 5 %	0.57	2.53	0.38	0.53	0.73
Interaction (S x G)					
S.Em±	0.49	2.18	0.33	0.45	0.62
C.D. at 5 %	1.41	6.20	0.93	1.29	1.78
C.V. %	4.72	4.94	4.51	4.30	4.71

The stem diameter (mm) at 90, 120 and 150 DAS as shown in Table 2.

Among seed soaking treatments, maximum stem diameter (16.68 cm, 23.40 cm and 27.81 cm) at 90, 120 and 150 DAS, respectively was observed in S₃ (GA₃ @ 100 ppm). Among growing media, maximum stem diameter (15.79 cm, 22.19 cm and 26.76 cm) at 90, 120 and 150 DAS, respectively was observed in G₃ [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum stem diameter (2.09 mm, 2.90 mm and 3.49 mm) at 90, 120 and 150 DAS, respectively.

This might be due to fact that the GA₃ caused greater cell division and elongation at the stem portion resulting into the greater stem diameter. The above results are conformity with Meena and Jain (2012) and Anjanawe and others (2013) in papaya, Parvin and others (2015) in black

walnut seeds. Synergistic combinations of these growing media attributed proper aeration, high water holding capacity and better nutrient availability that enhance the physical condition of growing media leading to higher production of photosynthetically functional leaves in this treatment resulting in better diameter of seedling (Borah and others 1994 in silk cotton). These findings are in close accordance with the result of Ramteke and others (2015) in papaya, Rahangdale (2019) in custard apple.

Table 2: Effect of seed soaking and growing media on seedling diameter (cm) and number of leaves of acid lime seedling

Treatments	Seedling diameter (cm)			Number of leaves		
	90 DAS	120 DAS	150 DAS	90 DAS	120 DAS	150 DAS
Factor A: Seed soaking (s)						
S ₁ : Water	1.44	1.87	2.27	10.15	15.52	19.42
S ₂ : GA ₃ @ 50 ppm	1.80	2.45	2.79	15.63	21.58	25.83
S ₃ : GA ₃ @ 100 ppm	1.94	2.66	3.11	17.82	24.10	28.78
S ₄ : Cow urine @ 25 %	1.53	2.03	2.40	12.18	17.55	21.87
S ₅ : Cow urine @ 50 %	1.60	2.10	2.44	12.63	18.07	22.45
S ₆ : Cow dung slurry (1:1 w/w)	1.66	2.18	2.56	13.43	19.13	23.33
S.Em±	0.02	0.03	0.03	0.19	0.20	0.24
C.D. at 5 %	0.06	0.08	0.09	0.54	0.56	0.67
Factor-B: Growing media (g)						
G ₁ : Sand + Vermicompost (1:1 v/v)	1.63	2.14	2.48	13.08	18.52	22.94
G ₂ : Sand + FYM (1:1 v/v)	1.47	1.95	2.33	11.20	16.24	20.41
G ₃ : Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)	1.90	2.57	3.02	16.94	23.49	27.66
G ₄ : Sand + FYM + Cocopeat (1:1:1 v/v/v)	1.64	2.20	2.55	13.34	19.04	23.44
S.Em±	0.02	0.02	0.02	0.16	0.16	0.19
C.D. at 5 %	0.05	0.07	0.07	0.44	0.46	0.55
Interaction (S x G)						
S.Em±	0.04	0.06	0.06	0.38	0.40	0.47
C.D. at 5 %	0.12	0.16	0.18	1.08	1.13	1.34
C.V. %	4.35	4.50	4.14	4.84	3.55	3.45




Figure 2: Effect of seed soaking and growing media on stem diameter (mm) at 90 DAS, 120 DAS and 150 DAS of acid lime

The number of leaves at 90, 120 and 150 DAS as shown in Table 2.

Among seed soaking treatments, maximum number of leaves (17.82, 24.10 and 28.78) at 90, 120 and 150 DAS, respectively was observed in S₃ (GA₃ @ 100 ppm). Among growing media, maximum number of leaves (16.94, 23.49 and 27.66) at 90, 120 and 150 DAS, respectively was observed in G₃ [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum number of leaves (19.87, 27.07 and 32.27) at 90, 120 and 150 DAS, respectively.

This might be due to the activity of GA₃ at apical meristem resulting in more synthesis of nucleoprotein which is responsible for increasing leaf number and expansion. An appropriate media mixture provides better root environment to the plant leading to better nutrient availability to the photo synthetically functional leaves that ultimately utilized for more number of leaves (Ramteke and others 2015). Similar findings were also reported by Meena and others (2012) and Anjanawe and others (2013) in Papaya and Parasana and others (2014) in mango.

The fresh and dry weight of shoot (gm) at 150 DAS as shown in Table 3.

Among seed soaking treatments, maximum fresh and dry weight of shoot (6.90 g and 2.60 g, respectively) at 150 DAS was observed in S₃ (GA₃ @ 100 ppm). Among growing media, maximum fresh and dry weight of shoot (6.60 g and 2.57 g, respectively) at 150 DAS was observed in G₃ [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S₃G₃ [GA₃ @ 100

ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum fresh and dry weight of shoot (7.57 g and 3.05 g, respectively) at 150 DAS.

This might be due to the combination containing many macro and micronutrients, humid acid, which maintaining proper aeration and porosity and GA₃, which increases cell division and uptake of nutrients, thus increasing the growth of seedling. The increase in fresh weight and dry weight might be because the combination of GA₃ and media increases the water and nutrient transportation to aerial parts, which leads to production of photosynthetic product and translocation of various plant parts, resulting in a higher fresh and dry weight of shoot. Similar findings were obtained by Kumawat and others (2014) and Dayeshwari and others (2017) in papaya.

The fresh and dry weight of root (gm) at 150 DAS as shown in Table 3.

Among seed soaking treatments, maximum fresh and dry weight of root (1.90 g and 0.62 g, respectively) at 150 DAS was observed in S₃ (GA₃ @ 100 ppm). Among growing media, maximum fresh and dry weight of root (1.83 g and 0.59 g, respectively) at 150 DAS was observed in G₃ [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum fresh and dry weight of root (2.59 g and 0.81 g, respectively) at 150 DAS.

The increased root growth is attributed to the fact that the combination of GA₃ and media provides a better root environment for the plant, resulting in better nutrient availability to the photo synthetically functional leaves and it also increases the translocation of food into the root zone, which leads to increase in fresh weight of root thereby increased dry weight of root. A similar result was recorded by Ramteke and others (2015) in papaya, Kaur (2017) in mango, Parab and others (2017) in papaya.

The survival percentage of seedling at 150 DAS as shown in Table 3.

Among seed soaking treatments, maximum survival percentage of seedling (88.75 %) at 150 DAS was observed in S₃ (GA₃ @ 100 ppm). Among growing media, maximum survival percentage (86.11 %) at 150 DAS, respectively was observed in G₃ [Sand + Vermicompost + Cocopeat (1:1:1)]. Interaction of S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] recorded maximum survival percentage of seedling (96.67 %) at 150 DAS.

This may be due to GA₃ helps in cell expansion and its elongation resulting better root and shoot growth, which supports and encourage better survival of the seedlings (Rahangdale 2019) and media containing vermicompost and cocopeat as most of the constituents provided a start for establishment of seedlings which further got supplemented by PGPR's. Good physical and biological conditions in media combination had positive effect on root and shoot growth which also helps in better survival. Similar results were obtained by Ramteke and others (2015) for papaya. These findings are in agreement with the results obtained by Gupta (1989) and Khatana and others (2015) in Kagzi Lime.

Table 3: Effects of seed soaking and growing media on fresh weight of shoot, dry weight of shoot, fresh weight of root, dry weight of root and survival percentage of acid lime seedling

Treatments	Fresh weight of shoot (g)	Dry weight of shoot (g)	Fresh weight of root (g)	Dry weight of root (g)	Survival percentage
Factor A: Seed soaking (s)					
S ₁ : Water	4.33	1.54	0.84	0.21	62.50
S ₂ : GA ₃ @ 50 ppm	6.02	2.33	1.61	0.52	80.83
S ₃ : GA ₃ @ 100 ppm	6.90	2.60	1.90	0.62	88.75
S ₄ : Cow urine @ 25 %	4.92	1.83	1.00	0.29	69.58
S ₅ : Cow urine @ 50 %	5.14	1.99	1.06	0.31	71.67
S ₆ : Cow dung slurry (1:1 w/w)	5.22	2.02	1.18	0.36	73.75
S.Em±	0.06	0.02	0.01	0.01	0.95
C.D. at 5 %	0.18	0.06	0.04	0.01	2.70
Factor-B: Growing media (g)					
G ₁ : Sand + Vermicompost (1:1 v/v)	5.24	2.01	1.14	0.34	73.33
G ₂ : Sand + FYM (1:1 v/v)	4.59	1.59	0.91	0.25	65.00
G ₃ : Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)	6.60	2.57	1.83	0.59	86.11
G ₄ : Sand + FYM + Cocopeat (1:1:1 v/v/v)	5.26	2.04	1.19	0.37	73.61
S.Em±	0.05	0.02	0.01	0.00	0.77
C.D. at 5 %	0.15	0.05	0.03	0.01	2.20
Interaction (S x G)					
S.Em±	0.13	0.04	0.03	0.01	1.89
C.D. at 5 %	0.36	0.12	0.08	0.03	5.39
C.V. %	4.10	3.48	3.96	4.35	4.40

CONCLUSION

From the present investigation, it can be concluded that seed soaking with GA₃ 100 ppm and growing media Sand + Vermicompost + Cocopeat (1:1:1) significantly influenced the seed germination, seedling growth and survivability of acid lime. The treatment combination S₃G₃ [GA₃ @ 100 ppm and Sand + Vermicompost + Cocopeat (1:1:1 v/v/v)] was found superior and most effective for better physiological development of acid lime seedlings over the rest of the treatment combinations.

REFERENCES

- Abirami, K., Rema, J. R., Mathew, P. A. and Sril, V. (2011). Response of nutmeg seeds to different nursery media. *Indian Journal of Horticulture*, 67(4):584-586.
- Anjanawe, S. R., Kanpure, R. N., Kachouli, B. K. and Mandloi, D. S. (2013). Effects of plant growth regulators and growth media on seed germination and growth vigour of papaya. *Annals of Plant and Soil Research*, 15(1):31-34.
- Anonymous, (2020-21) 2nd Advance Estimates, National Horticulture Database. National Horticulture Board, Ministry of Agriculture and Farmers Welfare. Horticulture Statistics Division, Department of Agriculture, Cooperation and Farmers Welfare, Government of India, New Delhi.
- Atefe, A., Ali, T., Shoor, M. and Davarynejad, G. H. (2012). Study of the effect of vermicompost as one of the substrate constituents on yield indexes of strawberry. *Journal of Horticultural Science and Ornamental Plants*, 4(3):241-246.
- Bhutiya, N. H. (2017). Effect of growing media and gibberellic acid on seed germination and growth of acid lime. *Thesis M.Sc. (Horti.)* submitted to JAU, Junagadh.
- Borah, A. S., Ray, A. N., Bhat, A. K., Maheswarappa, R., Subramanian, H. P., Dileep, P. M., Sudhakara, K., Santhoshkumar, A. V., Nazeema, K. K. and Ashokan, P. K. (1994). Effects of seed size, rooting medium and fertilizers on the growth of seedlings of silk cotton (*Ceiba pentandra* Linn). *The Indian Journal of forestry*, 17(4):293-300.
- Dayeshwari, D., Rayaprolu, S. and Jone, A. (2017). Effects of potting media on seed germination, seedling growth and vigour in papaya cv. CO-8. *International Journal of Pure and Applied Bioscience*, 3(5):505-512.
- Deb, P., Das, A., Ghosh, S. K. and Suresh, C. P. (2014). Improvement of seed germination and seedling growth of papaya (*Carica papaya* L.) through different pre-sowing seed treatments. *Acta Horticulturae*, 8(51):313-316.

- Desai, A. B.; Trivedi, A.; Panchal, B. and Desai, V. (2017). Effect of GA₃ and cow urine with different media on seed germination and seedling growth of papaya (*Carica papaya* L.) cv. Madhubindu under net house condition. *International Journal of Current Microbiology and Applied Sciences*, 6(9):828-834.
- Dilip, W. S., Singh, D., Moharana, D., Rout, S. and Patra, S. S. (2017). Effect of gibberellic acid (GA₃) different concentrations at different time intervals on seed germination and seedling growth of rangpur lime. *Journal of Agroecology and Natural Resources Management*, 4(2):157-165.
- Elza, D. L. (1949). Germination of citrus seeds and certain properties and their coats. *Palestine Journal of Botany*, 7:69-80.
- Gupta, O. P. (1989). Effect of gibberellic acid on seed germination in acid lime (*Citrus aurantifolia* Swingle) cv. Kagzi Lime. *Progressive Horticulture*, 21(3-4):246-248.
- Jaiswal, S. B., Nainwal, R. V., Supekar, S. J. and Manel, S. B. (2018). The effect of growth regulators and chemicals on growth of Kagzi Lime (*Citrus aurantifolia* Swingle). *International Journal of Current Microbiology and Applied Sciences*, 6(5):940-944.
- Kaur, S. (2017). Effects of growing mixtures on seed germination and seedling growth of different mango (*Mangifera indica* L.) cultivars under submountainous conditions of Punjab. *Chemical Science Review and Letters*, 6(23):1599-1603.
- Khan, M. M., Usman, M., Waseem, R. and Ali, M. A. (2002). Role of gibberellic acid (GA₃) on citrus seed germination and study of some morphological characteristics. *Pakistan Journal of Botany*, 39(2):113-118.
- Khatana, K. J., Jadav, R. G. and Nehete, D. S. (2015). Influence of GA₃ on germination and growth of acid lime cv. Kagzi Lime seed (*Citrus aurantifolia* Swingle) under field as well as net house conditions. *The Asian Journal of Horticulture*, 10(1):11-16.
- Kumawat, R., Maji, S., Govind, and Meena, D. C. (2014). Studies on seed germination and seedling growth of papaya (*Carica papaya* L.) cv. Coorg Honey Dew as influenced by media and chemicals. *Journal of Crop and Weed*, 10(2):281-286.
- Meena, R. R., Jain, M. C. and Mukerjee, S. (2012). Effect of pre-sowing dip treatment with gibberellic acid on germination and survivability of papaya. *Annals of Plant and Soil Research*, 5(1):120-121.
- Panse, V. G. and Sukhatme, P. V. (1985). *Statistical methods for agricultural workers*. 4th Edition., ICAR, New Delhi, pp 87-89.

- Parab, A. M., Mathad, T. C. and Malshe, K. V. (2017). Effects of pre-soaking chemicals on germination and subsequent seedling growth of papaya (*Carica papaya* L.) cv. Solo. *International Journal of Chemical Studies*, 5(4):1812-1816.
- Parasana, J. S., Leua, H. N. and Ray, N. R. (2013). Effect of different growing media mixture on germination and seedling growth of mango (*Mangifera indica* L.) cultivars under net house condition. *The Bioscan*, 8(3):897-900.
- Parvin, P., Khezri, M., Tavosolian, I. and Hosseini, H. (2015). The effect of gibberellic acid and chilling stratification on seed germination of eastern black walnut (*Juglans nigra* L.). *Journal of nuts*, 6(1):67-76.
- Rahangdale, P. (2019). Effects of GA₃ and date of sowing on seed germination, growth and survival of custard apple (*Annona squamosa* L.) seedlings. *Thesis M.Sc. (Horti.)*, submitted to JNKV, Jabalpur.
- Raj, A., Jhariya, M. and Toppo, P. (2014). Cow dung for eco-friendly and sustainable productive farming. *International Journal of Scientific Research*, 3(8):201-202.
- Ramteke, V., Paithankar, D. H., Ningot, E. P. and Kurrey, V. K. (2015). Effects of GA₃ and propagation media on germination, growth and vigour of papaya cv. Coorg Honey Dew. *The Bioscan*, 10(3):1011-1016.
- Suthar, D. P. (2019). Effect of growing media and gibberellic acid on seed vigour, growth and survival of acid lime (*Citrus aurantifolia* Swingle) cv. Kagzi lime. *Thesis M.Sc. (Horti.)* submitted to AAU, Anand.
- Tandon, K., Gurjar, P. K. S., Lekhi, R. and Soni, D. (2019). Effect of organic substances and plant growth regulators on seed germination and survival of Tamarind (*Tamarindus indica* L.) seedlings. *International Journal of Current Microbiology and Applied Sciences*, 8(2):2270-2274.
- Wagner-Junior, A., Alexandre, R. S., Negreiros, J. R. S., Parizzotto, A. and Bruckner, C. H. (2005). Influence of seed scarification and imbibition time on passion fruit (*Passiflora edulis flavicarpa* Degener) seed germination. *Revista Ceres*, 52(301):369-378.
- Yadav, R. K., Jain, M. C. and Jhakar, R. P. (2012). Effect of media on growth and development of acid lime (*Citrus aurantifolia* Swingle) seedling with or without *Azotobacter*. *African Journal of Agricultural Research*, 7(48):6421-6426.