

Histological studies on mango cultivars

Abstract

The present study was conducted on ten mango cultivars namely Alphonso, Bangalora, Kalepad, Himayuddin, Sendura, Mulgoa, Neelum, Rumani, Banganapalli and Swarnarekha and seasons are main and off-season at Kanyakumari and Tenkasi. The stem along with bud sample was collected during the month of October and November and conducted to examine the relation between internal structure of leafy and reproductive bud. During main season. Among ten mango cultivars, Bangalora and Swarnarekha have well developed at kanyakumar during main season for stage I, whereas Tenkasi, Alphonso and Bangalora well developed buds during the main season in stage I was observed. During off-season observed at kanyakumari Neelum and Alphonso had well developed for vegetative bud in stage I. At Tenkasi cultivars Kalapad and Bangalora had well developed for vegetative bud in stage I. These stages has distinguished buds might be quiescent and undifferentiated and the well development route of axillary buds of vegetative buds prevailing to climatic and physiological factors or cultural factors adopted (Moens, 1963; Kumar, 1982). The climatic factors prevailing at Kanyakumari during the main and off seasons for the months of August to September low temperature (19°C to 21°C) and high humidity (85 to 90%) and main season for the month of October to November, low temperature (21°C to 23°C) and humidity (80 to 85%). Whereas reproductive stage observed mango cultivars, Alphonso, Bangalora, Kalapad, Neelum and Swarnarekha had well developed reproductive buds in stage II & III during main season at kanyakumari. During off-season, cultivars of Himampasand, Kalapad, Alphoso, Neelum and Swarnarekha had well developed reproductive bud for stag II & III was favoured different stages of bud differentiation in the ten cultivars of mango.

Key words: mango cultivars, microtome, vegetative, reproductive

Introduction

Mango (*Mangifera indica* L.) belonging to the family Anacardiaceae occupies a pre-eminent place among the fruit crops grown in India and christened as the 'King of fruits' owing to its delicious flavor and taste. Mango is among the most important fruits produced in India. They are grown in an area of 1.23 million hectares with an annual production of almost 11.0 million

tonnes, which accounts for more than 55 percent of the world's total production. Indian mangoes are famous for their flavour, aroma, and taste in different shapes, sizes, and colours. Mangoes are very popular among people because of their wide range of adaptability, high nutritional value, richness in variety, delicious taste, and excellent flavour. It is a rich source of Vitamin A and Vitamin C. Mangoes can be eaten raw and ripe. Raw fruits of the local varieties are used to make products like raw pieces in brine, pickle, chutney, and many others. During the present time, raw mangoes are used to prepare pickles and raw slices. Alphonso is used in preparing squash in the Coastal Western zone. It is a general consideration that mango flowering is a complex phenomenon. Besides, favorable climate conditions that favours off - season flowering, genetic potential of the varieties, physiological and biochemical variations and better management interventions could also play the vital role in promoting off season flowering. The current research was to study the histological characteristics of some mango cultivars which may be associated with their flowering behavior.

Materials and Methods

An experiment was conducted at State Horticultural Farm, Kanyakumari and farmers field of Tenkasi was undertaken by the Horticultural College and Research Institute, Tamil Nadu Agricultural University, Periyakulam. The experiment was laid out in a Factorial Randomized Block Design (FRBD), with two seasons and ten varieties and replicated twice. Ten year old trees of mango cultivars were selected for this study. Ten mango cultivars selected for this study are Alphonso, Bangalora, Kalepad, Himampasand, Sendura, Mulgoa, Neelum, Rumani, Banganapalli and Swarnarekha and seasons are main and off-season. The weather parameters viz., Maximum and minimum temperature, relative humidity, average rainfall in Kanyakumari and Tenkasi were recorded in experimental locations. The stem along with bud sample was collected both the location during the month of October and November and immediately fixed in F.A.A., dehydration of the ethyl and butyl alcohol and then the samples were sectioned by rotary microtome and stained with Safranin (Staining in red the lignified cell walls) and light green (staining in blue green cellulose walls) and examined under light microscope (Trifilo, *et al.*2007 and Bacelar, *et al.*2007). Images were captured by light microscope supplement with camera (Panasonic WV-CP 220, Japan). The obtained images were subjected to analysis by image analysis software (Digimizer software package) and the following parameters

Result and discussion

Vegetative shoot

The present study was microtome analysis and observed in ten mango cultivars both locations for vegetative and reproductive shoots. The images were captured by light microscope supplement with camera, and distinguished the images for Vegetative shoot and reproductive shoot. Among ten mango cultivars, Bangalora and Swarnarekha mango cultivars well developed at kanyakumar during main season for stage I, whereas Tenkasi mango cultivars of Alphonso and Bangalora well developed buds during the main season in stage I. During off-season observed at kanyakumari cultivars Neelum and Alphonso had well developed for vegetative bud in stage I. At Tenkasi cultivars Kalapad and Bangalora had well developed for vegetative bud in stage I. The anatomy of mango vegetative shoot development has been described by (Singh, 1958; Chaikiattiyos *et al.*, 1994). Vegetative shoots may arise either from axillary buds, if no apical bud exists due to flowering in the previous flush, or from the apical bud when present. The latter is considered as extension growth or addition of an intercalary unit on the existing stem, but the developmental events during shoot formation from either apical or lateral buds are basically the same. Cells in the leaf primordia of initiating buds begin to form individual leaves in the proximal portion of the vegetative shoot of the mango cultivars.

Reproductive shoot

The reproductive shoot was observed both the locations during main and off-season for ten mango cultivars. Among ten mango cultivars, Alphonso, Bangalora, Kalapad, Neelum and Swarnarekha had well developed reproductive buds in stage II & III during main season at kanyakumari. During off-season, cultivars of Himampasand, Kalapad, Alphoso, Neelum and Swarnarekha had well developed reproductive bud for stag II & III. Two types of reproductive shoots typically occur in mango. Generative shoots display only flowers and have floral bracts or

non-developed leaves at the base of each lateral inflorescence. Terminal inflorescences, that is panicles or thyrsoids (Weberling, 1989), develop from dormant apical buds. The anatomy of panicle development has been described (Ravishankar *et al.*, 1979; Scholefield *et al.*, 1986). The complexes of primary to quaternary branching lateral structures of the mango inflorescence each terminate with three cymose flowers. The terminal flower opens first, followed by two subtending lateral flowers. These complexes form the lateral inflorescence structures emerging from the central axis of the panicle. The central axis extension also terminates in a similar fashion. Morphological stages of mango floral buds and panicle development were described by Shu (1981) and Oosthuysen (1991).

Generative shoot development in apical buds initially involves swelling of the lateral meristem tissues and their bud scales. Each axillary meristem develops as an inflorescence on a primary peduncle. The apical meristem then forms new lateral meristems and leaf primordia for the distal portion of panicle development if floral inductive conditions persist (Núñez-Elisea *et al.*, 1996). Panicles may be open or compact in nature, depending upon internodes elongation, which is cultivar dependent (Singh, 1960), but the architecture generally conforms to that in mixed shoots develop under weak floral inductive conditions (i.e. in the low-latitude tropics). Both leaves and primary pedunculate inflorescences develop from the same nodes. Leaf primordia and lateral meristems develop as leaf and floral structures, respectively.

Floral meristems could be easily distinguished from vegetative meristems due to their size difference. In general floral meristems are larger and broader than the vegetative ones. The size difference is caused by a marked increase in the frequency of cell division in the central meristematic zone during the transition period from vegetative stage to floral stage (Fosket, 1994). When the axillary bud follows the reproductive path, the central cells of the apical portion

undergo frequent periclinal and anticlinal divisions resulting in the broadening and enlargement of the apical zone (Moens, 1963).

Conclusion

Buds at stage I & II was observed both locations during main and off-season. During main season. Among ten mango cultivars, Bangalora and Swarnarekha have well developed at kanyakumar during main season for stage I, whereas Tenkasi, Alphonso and Bangalora well developed buds during the main season in stage I was observed. During off-season observed at kanyakumari Neelum and Alphonso had well developed for vegetative bud in stage I. At Tenkasi cultivars Kalapad and Bangalora had well developed for vegetative bud in stage I. These stages has distinguished buds might be quiescent and undifferentiated and the well development route of axillary buds of vegetative buds prevailing to climatic and physiological factors or cultural factors adopted (Moens, 1963; Kumar, 1982). The climatic factors prevailing at Kanyakumari during the main and off seasons for the months of August to September low temperature (19°C to 21°C) and high humidity (85 to 90%) and main season for the month of November to December, low temperature (21°C to 23°C) and humidity (80 to 85%). Whereas reproductive stage observed mango cultivars, Alphonso, Bangalora, Kalapad, Neelum and Swarnarekha had well developed reproductive buds in stage II & III during main season at kanyakumari. During off-season, cultivars of Himampasand, Kalapad, Alphoso, Neelum and Swarnarekha had well developed reproductive bud for stag II & III was favoured different stages of bud differentiation in the ten cultivars of mango.

References

- Bacelar, E.A., J.M. Moutinho-Pereira, B.C. Gonçalves, H.F. Ferreira and C.M. Carlos, 2007. Changes in growth, gas exchange, xylem hydraulic properties and water use efficiency of three olive cultivars under contrasting water availability regimes. *Environ. Exp. Bot.*, 60(2): 183-192.
- Chaikiattiyos, S., C.M. Menzel and T.S. Rasmussen. (1994). Floral induction in tropical fruit trees: effects of temperature and water-supply. *J. of Hort. Sci.* 69 : 397–415.

- Fosket. D.E.1994. Plant growth and development a molecular approach. Academic press.California.
- Kumar, D.1982. Primary investigations into some flowering abnormalities of coffee in Kenya. Kenya coffee, 47: 16 – 24.
- Moens. P.1963. Coffee canephora Pierre, (Morphological and morphogenetica). La cellule 63: 165 – 244.
- Nunez – Eliseas, R., T.L.Davenport and M.L. Caldeire. 1996. Control of bud morphogenesis in mango (*Mangifera indica* L.) by girdling and defoliation and temperature modification. J.Hort Sci. 71 (1): 25 – 40.
- Oosthuysen, S.A. 1991. Stages of development of the mango panicle. South African Mango Growers' Association Yearbook 11: 59–61.
- Ravishankar, H., M.M. Rao, and K.M. Bojappa, 1979. Fruit bud differentiation in mango “Alphonso” and “Totapuri” under mild tropical rainy conditions. Scientia Hort. 10 (1): 95 – 99.
- Scholfield, P. B., D. R. Oag and M. Sedgley. 1986. The relationship between vegetative and reproductive development in northern Australia. Aust. J. Agric. Res., 37:425-433.
- Shu, Z.H. (1981) Flower differentiation of mango (*Mangifera indica*). National Science Council Monograph 9: 865–870.
- Singh, L.B. 1960. The mango. Leonard Hill, London.
- Singh, R.N. 1958. Studies in the differentiation and development of fruit buds in mango (*Mangifera indica* L.) II. Morphological and histological changes. Hort. Adv., 2:37–43.
- Trifilo, P., M.A. Lo Gullo, A. Nardini, F. Pernice and S. Salleo, 2007. Rootstock effects on xylem conduit dimensions and vulnerability to cavitation of *Olea europaea* L. Trees-Struct. Funct., 21: 549-556.

Plate 9. Histological studies on mango varieties during main season at Kanyakumari











Vegetative shoots	Reproductive shoots
 <p align="center">Stage I</p>	 <p align="center">Stage II</p>
 <p align="center">Stage I</p>	 <p align="center">Stage II</p>
 <p align="center">Stage I</p>	 <p align="center">Stage III</p>
 <p align="center">Stage I</p>	 <p align="center">Stage II</p>
 <p align="center">Stage I</p>	 <p align="center">Stage II</p>

Plate 9A. Histological studies on mango varieties during main season at Kanyakumari (cont...)




Plate 8. Histological studies on off season mango varieties at Kanyakumari











Vegetative shoots	Reproductive shoots
	
Stage I	Stage II
Alphonso	
	
Stage I	Stage II
Bangalora	
	
Stage I	Stage II
Banganapali	
	
Stage I	Stage III
Himayuddin	
	
Stage I	stage V
Kalepad	

Plate 8A. Histological studies on off season mango varieties at Kanyakumari (cont...)

Vegetative shoots	Reproductive shoots
	
Stage I	Mulgoa
	
Stage I	Neelum
	
Stage I	Rumani
	
Stage I	Senduram
	
Stage I	Swarnarekhai

**Plate 11. Histological studies during main season
at Tenkasi**




































Vegetative shoots	Reproductive shoots
	
Stage I	Stage II
	
Stage I	Stage II
	
Stage I	Stage II
	
Stage I	Stage II
	
Stage I	Stage II

Plate 11A. Histological studies during main season at Tenkasi (cont...)









Vegetative shoots	Reproductive shoots
	
Stage I	Mulgoa
	
Stage I	Neelum
	
Stage I	Rumani
	
Stage I	Banganapalli
	
Stage I	Swarnarekha

**Plate 10. Histological studies during off season
at Tenkasi**

Vegetative shoots	Reproductive shoots
	
Stagel	Alphonso
	
Stagel	Bangalora
	
Stagel	Kalepad
	
Stagel	Himayuddin
	
Stagel	Sendura



**Plate 10A. Histological studies during off season
at Tenkasi (cont...)**

Vegetative shoots	Reproductive shoots
	
Stagel	Mulgoa
	
Stagel	Neelum
	
Stagel	Rumani
	
Stagel	Banganapalli
	
Stagel	Swarnarekha
Stage I	