

Interaction effect of phosphorous and zinc on yield attributes, yield and quality characteristics of chickpea under central plain zone of Uttar Pradesh

Comment [D1]: characteristics

Comment [D2]:

Comment [D3]: the central

Abstract

The present field experiments were conducted on studies effect of phosphorus and zinc on yield and quality parameter of chickpea was taken up at Student's Instructional Farm, at Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, (U.P.) India, during rabi season 2021-22. The experiment consists of 16 treatments combinations in factorial randomized block design with three replications consisted of different combination of phosphorus and zinc. Chickpea variety RVG-203 was grown with the recommended agronomic practices. On the basis of results emanated from investigation it can be concluded that among the yield components and productivity parameters maximum values in relation to number of pods plant⁻¹ (61.26), number of grains pod⁻¹ (1.73), 100 grain wt. (20.20 gm), grain yield (18.85 q ha⁻¹), straw yield (23.35 q ha⁻¹), biological yield (42.20 q ha⁻¹), harvest index (44.67 %) and protein content in grain (20.93 %) were found in the treatment T₁₅ [P₉₀Zn₆]. Corresponding minimum values in relation to number of pods plant⁻¹ (45.00), number of grains pod⁻¹ (1.20), 100 grain wt. (16.96 gm), grain yield (12.25 q ha⁻¹), straw yield (17.16 q ha⁻¹), biological yield (29.41 q ha⁻¹), harvest index (41.65 %) and protein content in grain (19.56 %) were associated with the treatment T₁[P₀Zn₀].

Comment [D4]: parameters

Comment [D5]:

Comment [D6]: chickpeas

Comment [D7]:

Comment [D8]: remove (was)

Comment [D9]: therabi

Comment [D10]: therabi

Comment [D11]: therabi

Comment [D12]: therabi

Comment [D13]: treatment

Comment [D14]: a factorial

Comment [D15]: a factorial

Comment [D16]: consisting

Comment [D17]: a different

Comment [D18]:

Comment [D19]: emanating

Comment [D20]: the investigation,

Comment [D21]: the number

Comment [D22]: a, and

Comment [D23]:

Key Words: Chickpea, Phosphorous, Protein, Yield and Zinc.

Introduction

India is the largest producer and consumer of pulses in the world. Among the pulses, chickpea is the most important grown in every part of India. It is largest produced food legume in South Asia. Chickpea (*Cicer arietinum* L.) is a major legume crop cultivated for its edible seeds legume of the genus *Cicer*, Tribe *Cicereae*, family *Fabaceae* (*leguminaceae*), and subfamily *Papilionaceae*. It provide protein rich diet to the vegetarian of the Indian and complement the stable cereals in the diets with proteins, essential amino acids, vitamins and minerals (Pingoliya et al. 2013). They contain carbohydrate (61.51%), fat (4.5%) and relatively free from anti nutritional factors. Chickpea is rich in protein content (20.47g/100g), carbohydrate (62.95g/100g), fibre (12.2g/100g), phosphorous (252mg/100g), high amount of minerals such as calcium

Comment [D24]:

Comment [D25]: the largest

Comment [D26]: provides

Comment [D27]: vegetarian Indian

Comment [D28]: complements

Comment [D29]:

Comment [D30]: carbohydrates

Comment [D31]: and fat

Comment [D32]: are relatively

Comment [D33]: fiber

(57mg/100g), magnesium (79mg/100g), iron (4.31mg/100g) and zinc (15mg/100g), low in fat content and most of it is polyunsaturated (Wallace *et al.* 2016).

India is the largest producer (25% of global production), consumer (27% of world consumption) and importer (14%) of pulses in the world. India ranks first in the world in terms of pulse production (25% of total worlds production) (FAOSTAT 2017). In India chickpea occupies 10.17 million ha area, with a production of 11.35 million tonnes registering the productivity of 1116 kg/ha. In Uttar Pradesh, chickpea crop occupied 0.62 million hectares area, 0.85 million tonnes production and 1371 kg/ha productivity (Anonymous, 2021).

Phosphorus deficiency can limit nodule number, leaf area, and biomass and grain development in legumes. Symbiotic nitrogen fixation has a high P demand because the process consumes large amounts of energy (Schulze *et al.*, 2006) and energy generating metabolism strongly depends upon the availability of P (Plaxton, 2004). Singh and Sale (2000) reported that P fertilization stimulates root growth, photosynthesis and increases hydraulic conductivity of roots. Phosphorus is used in numerous molecular and biochemical plant processes, particularly in energy acquisition, storage and utilization (Epstein and Bloom, 2005). The phosphorus content per unit dry weight is usually considerably higher in the nodules than in the roots and shoots, particularly at low external phosphorus supply. Nitrogen fixing plants have an increased requirement for P over dose receiving direct nitrogen fertilization, probably due to need for nodule development and signal transduction, and to P-lipids in the large number of bacterioids (Graham and Vance, 2000). It acts as a catalyst in several biochemical reactions occurred in plants. It plays an important role in capturing and converting the solar energy into useful plant compounds. These compounds help in the general health and vigor of plants (Griffith, B. 2010). Legumes are heavy feeder of phosphorus and less responsive to nitrogen because of their capacity to meet their own nitrogen requirement through symbiotic fixation (Kumar *et al.* 2016). Phosphorus is connected with some particular plant growth factors that are root development, vigorous stem, enhanced flower formation and seed production, earlier and more uniform crop maturity, increase nitrogen fixing capacity of legumes, improvement in crop quality and resistance to plant diseases (Rehan *et al.* 2018). It is required for higher and sustainable production of grain legumes. Generally, legumes have higher P requisites due to more consumption of energy in the process of symbiotic nitrogen fixation (Islam *et al.* 2012).

Comment [D34]: the consumer

Comment [D35]: the importer

Comment [D36]: remove (s) only world

Comment [D37]:

Comment [D38]: a productivity

Comment [D39]: the chickpea

Comment [D40]: remove (area)

Comment [D41]: of production, and

Comment [D42]: biomass, and

Comment [D43]:

Comment [D44]: energy-generating

Comment [D45]:

Comment [D46]: the hydraulic

Comment [D47]:

Comment [D48]: the need

Comment [D49]:

Comment [D50]: a large

Comment [D51]:

Comment [D52]: bacterioids

Comment [D53]: remove (the)

Comment [D54]: feeders

Comment [D55]: are (remove) that

Comment [D56]: more sustainable

Comment [D57]: more energy consumption in the symbiotic nitrogen fixation process (Islam *et al.* 2012).

Phosphorus enhances the activity of rhizobia and increases the formation of root nodules thereby helps in fixing more of atmospheric nitrogen in root nodules. Phosphorus is also an important fertilizer in chickpea production; it is a very important chemical fertilizer that can raise the water holding capacity of soil (Dotaniya et al., 2014).

Comment [D58]: helping in fixing more atmospheric

Comment [D59]: water-holding capacity of the soil

Zinc is required for the proper functioning of various metabolic processes. It's necessary for chlorophyll and carbohydrate production. Several enzyme systems, auxin and protein synthesis, seed formation and maturity rate all require zinc, either directly or indirectly. Zinc is known to help with RNA synthesis, which is required for protein production. In the plant, zinc is not translocated. As a result, symptoms occur first on the plant's younger leaves and other sections. Stunted growth, the formation of light green yellowish patches and chlorotic bands on either side of the midrib in the plants are all common symptoms of zinc deficiency. The sustainable production needs balanced supply from soil along with suitable physical and biological properties to attain a better growth of roots and efficient utilization of nutrient from the rhizosphere.

Comment [D60]:

Comment [D61]: Sustainable production needs a balanced supply of soil

Comment [D62]: nutrients

The soils of Kanpur are alkaline in nature, low in organic content in generally low in fertility status. All these factors lead to the deficiency of zinc in soils. As such, it is possible that the application of zinc may be helpful in increasing the yields of chickpea under agro climatic condition of Kanpur.

Comment [D63]: remove (it is possible that) the applicationcondition of Kanpur

Comment [D64]:

Comment [D65]: chickpeas

Comment [D66]: the agro-climatic conditions

Comment [D67]:

Comment [D68]:

Resources and Methods

Experimental Site

The experiment was conducted during rabi season of 2021-22 at student's Instructional farm, C.S.A. University of Agriculture and Technology, Kanpur Nagar (U.P.). The field was well leveled and irrigated by tube well. The farm is situated at main campus of the university, in the west northern part of Kanpur city under sub-tropical zone in vth agroclimatic zone (central plain zone).

Comment [D69]:

Comment [D70]: therabi

Comment [D71]: the student's

Comment [D72]: a tube

Comment [D73]: the main

Comment [D74]:

Comment [D75]: west-northern part of Kanpur city under the sub-tropical

Edaphic Condition

The soil was moist, well drained with uniform plane topography. The soil of the experimental field was alluvial in origin, sandy loam in texture and slightly alkaline in reaction having pH 7.9 (1:2.5 soil: water suspension method given by Jackson, 1973), electrical conductivity 0.30 dSm⁻¹ (1:2.5 soil: water suspension method given by Jackson, 1973), Organic carbon percentage in soil

is 0.45 per cent (Walkley and Black's rapid titration method given by **Walkley and Black, 1934**), with available nitrogen 210 kg ha⁻¹ (Alkaline permanganate method given by **Subbiah and Asija, 1956**), available phosphorus as sodium bicarbonate-extractable P was 12.8 kg ha⁻¹ (Olsen's calorimetrically method, **Olsen et al., 1954**) available potassium was 198 kg ha⁻¹ (Flame photometer method given by **Hanwey and Heidel, 1952**), available and available zinc was 0.55 kg ha⁻¹ (DTPA extraction method given by (**Lindsay and Norvell, 1978**)).

Comment [D76]:

Comment [D77]: the soil is 0.45 percent

Detail of treatments and design

The 16 treatments combination of nutrient management practices of inorganic fertilizer (DAP and zinc sulphate). Experiment was laid out in factorial randomized block design with three replications.

Comment [D78]: The experiment was laid out in a factorial

Table -1: detail of the treatment combinations:

S. No.	Symbol	Treatment combination
1.	T ₁	P ₀ Zn ₀
2.	T ₂	P ₀ Zn ₃
3.	T ₃	P ₀ Zn ₆
4.	T ₄	P ₀ Zn ₉
5.	T ₅	P ₃₀ Zn ₀
6.	T ₆	P ₃₀ Zn ₃
7.	T ₇	P ₃₀ Zn ₆
8.	T ₈	P ₃₀ Zn ₉
9.	T ₉	P ₆₀ Zn ₀
10.	T ₁₀	P ₆₀ Zn ₃
11.	T ₁₁	P ₆₀ Zn ₆
12.	T ₁₂	P ₆₀ Zn ₉
13.	T ₁₃	P ₉₀ Zn ₀
14.	T ₁₄	P ₉₀ Zn ₃
15.	T ₁₅	P ₉₀ Zn ₆
16.	T ₁₆	P ₉₀ Zn ₉

Crop Husbandry

A pre-sowing irrigation (Paleva) was done in the experimental field with an object to get optimum moisture conditions for attaining good germination. At proper tillage, one ploughing with tractor drawn mould board plough was done followed by two ploughings by cultivator. Full dose of Phosphorus, and Zinc were applied as basal at the time of sowing in the form of DAP and Zinc sulphate respectively. The sowing of seeds of Chickpea variety RGV-203 was done by line sowing by hand at 4-5 cm depth of soil and with line to line spacing of 45 cm to maintain uniform plant population.

Comment [D79]: Pre-sowing

Harvesting and threshing: The crop was harvested at maturity and was allowed to dry in sun. Separate bundles were made for each plot and weighed. The after drying harvest was threshed manually.

Data Collection

Grain yield

After threshing the grain yield from each plot was separately weighed and recorded after converting into quintals per hectare.

Comment [D80]: threshing,

Comment [D81]: it into

Straw yield

After subtracting the grain yield per plot from the total biological yield. After converting the yields into quintals per hectare, yields were recorded.

Biological yield (q ha^{-1})

Seed yield and Stover yield together were regarded as biological yield. The biological yield was calculated with the following formula:

Comment [D82]: the following

$$\text{Biological yield} = \text{Seed yield} + \text{Stover yield}$$

Harvest index(%):

The recovery of grains in total dry matter was considered as harvest index, expressed in percentage.

Comment [D83]:

Comment [D84]: the harvest

It has been calculated by following formula:

$$\text{Harvest Index (\%)} = [\text{Seed Yield (q ha}^{-1}) / \text{Biological Yield (q ha}^{-1})] \times 100$$

Protein content (%):

Nitrogen content (%) in grains was determined by Kjeldahl's method. Protein content of chickpea seed was determined by multiplying the N content of chickpea seed with factor 6.25 ($\text{N} \% \times 6.25$). (A.O.A.C., 1970).

Comment [D85]: The protein

Comment [D86]: the chickpea

Comment [D87]:

Comment [D88]: the chickpea

Comment [D89]: a factor of 6.25

Statistical analysis: The growth parameters and yields were recorded and analysed as per Gomez and Gomez (1984) the tested at 5% level of significance to interpret the significant differences.

Comment [D90]: a 5%

Result and Discussion

Yield Components

At a glance over the data given in the Table-2 and depicted in Fig.-1 clearly shows that among the yield attributing characters of chickpea such as number of pods plant⁻¹, number of grain pod⁻¹ and 100 grain weight (gm) significantly increase due to the application of Phosphorus and Zinc. Significantly response on yield components was recorded with T₁₅ [P₉₀ Zn₆] over other treatments. The number of pods plant⁻¹, number of grain pod⁻¹ and 100 grain weight (gm) increased to the magnitude of 45.00 to 61.26, 1.20 to 1.73 and 16.96 to 20.20, respectively. Maximum number of pods plant⁻¹ (61.26), number of grain pod⁻¹ (1.73) and 100 grain weight (20.20 gm) was associated with the treatment T₁₅ [P₉₀ Zn₆]. Minimum number of pods plant⁻¹ (45.00), number of grain pod⁻¹ (1.20) and 100 grain weight (16.96 gm) was associated with the treatment T₁ [Control]. The results of the present investigation are also in agreement with the findings of Yadav *et al.* (2022), Singh *et al.* (2022), Kumar *et al.* (2022), Yadav *et al.* (2022) and Sachanet *et al.* (2022)

Comment [D91]: A glance

Comment [D92]:

Comment [D93]:

Comment [D94]: Table 2

Comment [D95]: the number

Comment [D96]:

Comment [D97]: the number

Comment [D98]: Significant

Comment [D99]:

Comment [D100]: , and 100-grain

Comment [D101]: The maximum


Comment [D102]:

Comment [D103]: , and 100-grain

Table-2: Effect of different treatment combinations on yield components of chickpea

Treatment	No of pods plant ⁻¹	No of grains pod ⁻¹	100 Grains weight (g)
T ₁	45.00	1.20	16.96
T ₂	47.21	1.27	17.42
T ₃	52.21	1.36	17.65
T ₄	48.43	1.31	17.53
T ₅	50.12	1.32	17.61
T ₆	54.31	1.48	17.71
T ₇	55.18	1.49	17.87
T ₈	53.42	1.43	17.68
T ₉	56.21	1.52	17.94
T ₁₀	59.87	1.62	18.12
T ₁₁	60.89	1.71	19.82
T ₁₂	58.13	1.57	18.10
T ₁₃	57.32	1.55	18.03
T ₁₄	60.38	1.69	19.11
T ₁₅	61.26	1.73	20.20
T ₁₆	60.12	1.65	18.42

S.E (m) ±	P	0.055	0.34	0.010
	Zn	0.055	0.49	0.010
	P × Zn	0.110	0.59	0.019
CD at 5%	P	0.159	1.03	0.028
	Zn	0.159	1.48	0.028
	P × Zn	NS	NS	NS



Comment [D104]:

Comment [D105]: Check the word (characters)

Fig.-1: Effect of different treatment combinations on yield components of chickpea Productivity Parameters

It is visualized from the data given in Table-3 and depicted in Fig.-2 clearly indicate that among the productivity parameters viz. grain yield (q ha^{-1}), straw yield (q ha^{-1}), biological yield (q ha^{-1}) and harvest index (%) significantly increase due to the application of Phosphorus and Zinc. Grain yield varied from 12.25 to 18.85 q ha^{-1} , straw yield varied from 17.16 to 23.35 q ha^{-1} , biological yield varied from 29.41 to 42.20 q ha^{-1} and harvest index varied from 41.65 to 44.67 %. The maximum grain yield (18.85 q ha^{-1}), straw yield (23.35 q ha^{-1}), biological yield (42.20 q ha^{-1}) and harvest index (44.67 %) was associated with the treatment T₁₅ [P₉₀Zn₆] during the experimentation. The minimum grain yield (12.25 q ha^{-1}), straw yield (17.16 q ha^{-1}), biological yield (29.41 q ha^{-1}) and harvest index (41.65 %) was under the treatment T₁ [control] during the experimentation. The surge in seed and straw yields under adequate nutrients supply might be attributed to mainly to the collective effect of a greater number of number of podplant⁻¹, number of grains pod⁻¹ and higher test weight, which was the result of improved translocation of photosynthates from source to sink ultimately yield is increased. The increase in grain yield

Comment [D106]: were

Comment [D107]: were

Comment [D108]: remove (to)

Comment [D109]: number of pod plant-1, the number of grains

under adequate nutrients supply mainly due to more yield attributes ultimately resulted more grain yield. Grain, straw yield, biological yield and harvest index of chickpea significantly increased due to application of P 90 (kg ha⁻¹) and Zinc 6(kg ha⁻¹) over their controls. These results also confirms the findings of Singhet *et al.* (2021), Pal *et al.* (2021), Yadav *et al.* (2022), Kumar *et al.* (2022) and Sachan *et al.* (2022)

Table-3: Effect of different treatment combinations on productivity parameters of chickpea

Treatment	Grain yield (qha ⁻¹)	Straw yield (qha ⁻¹)	Biological yield (qha ⁻¹)	Harvest index (%)	
T ₁	12.25	17.16	29.41	41.65	
T ₂	12.52	17.35	29.87	41.91	
T ₃	13.76	18.72	32.48	42.36	
T ₄	12.87	17.81	30.68	41.95	
T ₅	13.24	18.25	31.49	42.05	
T ₆	14.86	19.65	34.51	43.05	
T ₇	15.34	20.12	35.46	43.26	
T ₈	14.25	19.10	33.35	42.73	
T ₉	15.77	20.46	36.23	43.53	
T ₁₀	16.96	21.78	38.74	43.78	
T ₁₁	18.24	22.96	41.20	44.27	
T ₁₂	16.71	21.34	38.05	43.91	
T ₁₃	16.10	20.95	37.05	43.45	
T ₁₄	17.65	22.54	40.19	43.92	
T ₁₅	18.85	23.35	42.20	44.67	
T ₁₆	17.31	22.24	39.55	43.76	
S.E (m) ±	P	0.022	0.021	0.043	0.010
	Zn	0.022	0.021	0.043	0.010
	P×Zn	0.044	0.043	0.087	0.019
CD at 5%	P	0.064	0.062	0.126	0.028
	Zn	0.064	0.062	0.126	0.028
	P×Zn	NS	NS	NS	NS

Comment [D110]: nutrient

Comment [D111]: resulted in more grain

Comment [D112]: chickpeas

Comment [D115]:

Comment [D116]: the application

Comment [D117]:

Comment [D118]: confirm

Comment [D119]:

Comment [D120]: Table 3

Comment [D113]:

Comment [D114]: I did look but must've missed it! So q stands for a *quintale* then or 100kg. Would I be right in thinking that *quintal* is not very common in modern English though?




Fig.-2:Effect of different treatment combinations on productivity parameters of chickpea

Quality parameter

Protein

It is visualized from the data given in Table-4 clearly indicate that among the quality parameters viz. protein content increase due to the application of Phosphorus and Zinc. Protein content varied 19.56 to 20.93 (%). The maximum protein content 20.93 % was associated with the treatment T₁₅ [P₉₀Zn₆]. The minimum protein content 19.53% was under the treatment T₁ [control]. Similar findings were reported by Ahmed *et al.* (2003), Vikrant *et al.* (2005) and Tripathy *et al.* (2019)

- Comment [D121]:
- Comment [D122]: Table 4 clearly indicates
- Comment [D123]:
- Comment [D124]: from 19.56

Table-4:Effect of different treatment combinations on quality parameters of chickpea

Treatment	Treatment combination	Protein content in grain (%)
T ₁	P ₀ Zn ₀	19.56
T ₂	P ₀ Zn ₃	19.68
T ₃	P ₀ Zn ₆	19.94
T ₄	P ₀ Zn ₉	19.81
T ₅	P ₃₀ Zn ₀	19.87
T ₆	P ₃₀ Zn ₃	20.18
T ₇	P ₃₀ Zn ₆	20.25

T ₈	P ₃₀ Zn ₉	20.06
T ₉	P ₆₀ Zn ₀	20.37
T ₁₀	P ₆₀ Zn ₃	20.62
T ₁₁	P ₆₀ Zn ₆	20.87
T ₁₂	P ₆₀ Zn ₉	20.56
T ₁₃	P ₉₀ Zn ₀	20.44
T ₁₄	P ₉₀ Zn ₃	20.75
T ₁₅	P ₉₀ Zn ₆	20.93
T ₁₆	P ₉₀ Zn ₉	20.68
S.E (m) ±	P	0.34
	Zn	0.29
	P×Zn	0.49
CD at 5%	P	1.03
	Zn	1.03
	P×Zn	NS

Conclusion

The current study demonstrates the benefit of application of phosphorus 90 kg ha⁻¹ and zinc 6 kg ha⁻¹ significantly increased yield attributes, productivity and quality parameter like number of pod plant⁻¹, number of grains pod⁻¹, test weight of 100 seeds, grain yield, straw yield, biological yield, harvest index and protein quality. Finally it can be concluded that the treatment T₁₅ [P₉₀Zn₆] is a best option for improving productivity, yields and quality parameter of chickpeacrop.

Comment [D125]: the application

Comment [D126]:

Comment [D127]: , and quality parameters like the number

Comment [D128]: the best

Comment [D129]:

Comment [D130]: crops.

References

Ahmed, M. K. A., Afifi, M. H. and Mohamed, M. F. (2003). Effect of biofertilizers, chemical and organic fertilizers on growth, yield and quality of some leguminous crops. *Egyptian Journal of Agronomy*. **25**: 45-52.

Anonymous. (2021). Agricultural Statistics at a Glance 2020. Directorate of Economics & Statistics, Department of Agriculture, Cooperation and Farmers Welfare, Ministry of Agriculture & Farmers Welfare, Govt. of India, New Delhi, p. 63.

Dotaniya ML, Datta SC, Biswas DR, Kumar K. (2014), Effect of organic sources on P₂O₅ fraction and available P₂O₅ in typical Haplustept. *J. Indian Soc. Soil Sci.* 62(1):80-83.

Epstein, E. and Bloom, A.J. (2005). *Mineral nutrition of plants: Principles and perspectives*, 2nd ed. Sunderland, Massachusetts: Sinauer Associates.

Gomez, K. A., & Gomez, A. A. (1984). *Statistical procedures for agricultural research*. John Wiley & sons.

Graham, P.H. and Vance, C.P. (2000). Nitrogen fixation in perspective, an overview of research and extension needs. *Field Crops Research* 65: 93-106.

Griffith B (2010). Efficient Fertilizer Use Phosphorus, 1-7.

Hanway, J.J; and Heidel, H. (1952). Soil analysis methods as used in Iowa State College, Soil Testing Laboratory. *Iowa Agriculture* 54: 1-31.

Islam M, Mohsan S, Ali S, Khalid R & Afzal S (2012). Response of chickpea to various levels of phosphorus and sulphur under rainfed conditions in Pakistan. *Romanian AgricRes.* 29: 175-183.

Jackson, M.L. (1973). Soil chemical analysis. Prentice Hall of India Pvt. Ltd, New Delhi.

Kumar, K., Pyare, R., Niwas, R., Tiwari, K., Sachan, R., Pal, R. K., Patel, V. K. & Ranjan, A. R., (2022). Studies on the Root Architecture with Nodulation of the Chickpea (*Cicer arietinum* L.) as Influence by Different Moisture Management Practices along with Seed Inoculation and Level of Zinc. *International Journal of Environment and Climate Change*, 2896-2904.

Kumar, K., Pyare, R., Verma, V. K., Sachan, R., Niwas, R., Yadav, A., Pal, R. K. & Ranjan, A. R., (2022). Impact of Moisture Conservation Practices, Seed Inoculation and Zinc Level on Growth and Yield of Chickpea (*Cicer arietinum* L.). *International Journal of Plant & Soil Science*, 34(23), 546-556.

Comment [D131]:

Comment [D132]: Wiley

Kumar, S., Tripathi, D.K., Bharose, R., Kumar, M. and Kumar, R. (2016). Effect of different fertility level and micronutrients on nodulation and nutrient uptake by chickpea. *An Asian Journal of Soil Sci.* **11**(1): 62-66.

Comment [D133]:

Comment [D134]: levels

Lindsay, W. L., & Norvell, W. (1978). Development of a DTPA soil test for zinc, iron, manganese, and copper. *Soil science society of America journal*, **42**(3), 421-428.

Olsen, S.R, Cole, C.V., Watanable, F. S. and Dean, L. A. (1954). Estimation of available phosphorous in soil by extraction with sodium bicarbonate. *USDA, Cric.* **930**:19- 23

Pal, S., Pandey, S. B., Singh, A., Singh, S., Sachan, R., & Yadav, A. (2021). Effect of Phosphorus, Boron and *Rhizobium* inoculation on productivity and profitability of chickpea. *The Pharma Innovation Journal* 2021; 10(12): 1810-1814

Pingoliya, K. K., Dotaniya, M. L., Mathur, A. K. (2013). Role of phosphorus and iron in chickpea (*Cicer arietinum* L.). *Lap Lambert Academic Publisher, Germany.*

Plaxton, W.C. (2004). Plant response to stress: biochemical adaptations to phosphate deficiency. In: Goodman R (eds.) *Encyclopedia of Plant and Crop Science*. New York:Marcel Dekker. pp. 976-980.

Rehan, W., Jan, A., Liaqat, W., Jan, F.M., Ahmadzai, M.D., Ahmad, H., Haroon, J., Anjum, M.M. and Ali, N. (2018). Effect of phosphorous, rhizobium inoculation and residue types on chickpea productivity. *Pure Appl. Biol.*, **7**(4): 1203-1213.

Sachan, R., Pandey, S. B., Kumar R., Singh D., Sharma, S., Kumar. K., Singh, A., Pal, S., Verma, A. K. and Sachan, H. (2022)Effect of Phosphorous, Chloropyriphos and *Rhizobium* Inoculation on Production and Economics of Chickpea (*Ciceraretinum* L.). *International Journal of Plant & Soil Science*, 1390-1398.

Sachan, R., Pandey, S. B., Kumar, A., Pathak, R. K., Pandey, H. P., Singh, A., ... & Kumar, M. (2022). Interaction effect of Phosphorous, Chloropyriphos and *Rhizobium* Inoculation on Growth Characteristics, Yield Components and Productivity Parameters of Chickpea under Central Plain Zone of Uttar Pradesh. *AMA ISSN: 00845841* Volume 53, Issue 08, August, 2022

Comment [D135]: characteristics,

Comment [D136]: August 2022

Schulze, J., Temple, G., Temple, S.J., Beschow, H., and Vance, C.P. (2006). Nitrogen fixation by white lupin under phosphorus deficiency. *Annals of Botany* **98**: 731-740.

Singh, A. K., Dimree, S., Kumar, A., Sachan, R., Sirohiya, A., & Nema, S. (2022). Effect of *rhizobium* inoculation with different levels of inorganic fertilizers on yield, nutrient content & uptake of chickpea (*Cicer arietinum* L.). *Int J Plant Soil Sci*, 34(22), 262-8.

Singh, A., Singh, D., Kumar, R., Pal, S., Sachan, R. & Yadav, A. (2021). Study the effect of organic, inorganic and biofertilizers on nutrients content and uptake of chickpea (*Cicer arietinum* L.). *The Pharma Innovation Journal* 2021; 10(10): 418-423

Singh, D.K. and Sale, P.W.G. (2000). Growth and potentially conductivity of white clover roots in dry soil with increasing phosphorus supply and defoliation frequency. *Agronomy Journal* 92: 868-874.

Subbiah, B.V. and Asija, C.L. (1956). A rapid procedure for the estimation of available N in Soil. *Curr. Sci.* 25:259-260.

Tripathy, D.K., Kumar, S. and Zaidy, S.F.A. (2019). Effect of phosphorus, sulphur and micronutrients (Zinc and Boron) levels on performance of chickpea (*Cicerarietinum* L.). *Natl. Acad. Sci. Lett.* 10.1007/s40009-019-00802-4.

Vikrant, Singh, H., Malik, C.V.S. and Singh, B.P. (2005). Grain yield and protein content of cowpea as influenced by farm yard manure and phosphorus application. *Indian Journal of Pulses Research* 18 (2): 250-251.

Walkley, A. and Black, C. S.A. (1934). Old piper, S.S. soil and plant analysis. *Soil Sci.* 37:29-38.

Wallace, T. C., Murray, R., Kathleen, M. and Zelman, K. (2016). The nutritional value and health benefits of chickpeas and humus. *Nutrients.* 8(12): 766.

Yadav, A., Singh, D., Kumar, R., Sachan, R., Kumar, K., Singh, A., & Singh, K. K. (2022). Response of different level of phosphorus, zinc and Rhizobium inoculation on growth yield attributes and yield of Chickpea (*Cicerarietinum* L.). *International Journal of Environment and Climate Change*, 12(11), 1954-1964.

Yadav, P., Yadav, D. D., Pandey, H. P., Yadav, A., Sachan, R., & Yadav, S. (2022). Effect of Fertility Levels and Biofertilizers on Growth Parameters, Root Architecture and Quality of Chickpea (*Cicer arietinum* L.). *International Journal of Plant & Soil Science* 34(17): 61-67, 2022; Article no.IJPSS.86643 ISSN: 2320-7035

Comment [D137]: potential

Yadav, S., Yadav, D.D., Kumar, A., Sachan, R. & Yadav, S. (2022). Effect of Fertility Levels and Bio-fertilizers Application on Yield, Yield Attributes, and Economics of Chickpea (*Cicer arietinum* L.). *International Journal of Plant & Soil Science* 34(13): 65-69, 2022; Article no.IJPSS.85129 ISSN: 2320-7035

UNDER PEER REVIEW