

Interaction effect of phosphorous and zinc on yield attributes, yield and quality characteristics of chickpea under the central plain zone of Uttar Pradesh

Abstract

The present field experiments were conducted on studies effect of phosphorus and zinc on yield and quality parameters of chickpeas taken up at Student's Instructional Farm, at Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, (U.P.) India, during the *rabi* season 2021-22. The experiment consists of 16 treatment combinations in a factorial randomized block design with three replications consisting of a different combination of phosphorus and zinc. Chickpea variety *RVG-203* was grown with the recommended agronomic practices. On the basis of results emanating from the investigation, it can be concluded that among the yield components and productivity parameters maximum values in relation to the number of pods plant⁻¹ (61.26), number of grains pod⁻¹ (1.73), 100 grain wt. (20.20 gm), grain yield (18.85 q ha⁻¹), straw yield (23.35 q ha⁻¹), biological yield (42.20 q ha⁻¹), harvest index (44.67 %) and protein content in grain (20.93 %) were found in the treatment T₁₅ [P₉₀Zn₆]. Corresponding minimum values in relation to number of pods plant⁻¹ (45.00), number of grains pod⁻¹ (1.20), 100 grain wt. (16.96 gm), grain yield (12.25 q ha⁻¹), straw yield (17.16 q ha⁻¹), biological yield (29.41 q ha⁻¹), harvest index (41.65 %) and protein content in grain (19.56 %) were associated with the treatment T₁ [P₀Zn₀].

Key Words: Chickpea, Phosphorous, Protein, Yield and Zinc.

Introduction

“India is the largest producer and consumer of pulses in the world. Among the pulses, chickpea is the most important grown in every part of India. It is the largest produced food legume in South Asia. Chickpea (*Cicer arietinum* L.) is a major legume crop cultivated for its edible seeds legume of the genus *Cicer*, Tribe *Cicereae*, family *Fabaceae* (*leguminaceae*), and subfamily *Papilionaceae*. It provides protein rich diet to the vegetarian Indian and complements the stable cereals in the diets with proteins, essential amino acids, vitamins and minerals” (Pingoliya *et al.* 2013). “They contain carbohydrates (61.51%) and fat (4.5%) are relatively free from anti nutritional factors. Chickpea is rich in protein content (20.47g/100g), carbohydrate (62.95g/100g), fiber (12.2g/100g), phosphorous (252mg/100g), high amount of minerals such as

calcium (57mg/100g), magnesium (79mg/100g), iron (4.31mg/100g) and zinc (15mg/100g), low in fat content and most of it is polyunsaturated” (Wallace *et al.* 2016).

“India is the largest producer (25% of global production), the consumer (27% of world consumption) and the importer (14%) of pulses in the world. India ranks first in the world in terms of pulse production (25% of total world production) (FAOSTAT 2017). In India chickpea occupies 10.17 million ha area, with a production of 11.35 million tonnes registering a productivity of 1116 kg/ha. In Uttar Pradesh, the chickpea crop occupied 0.62 million hectares, 0.85 million tonnes of production, and 1371 kg/ha productivity” (Anonymous, 2021).

“Phosphorus deficiency can limit nodule number, leaf area, biomass and grain development in legumes. Symbiotic nitrogen fixation has a high P demand because the process consumes large amounts of energy” (Schulze *et al.*, 2006) and “energy-generating metabolism strongly depends upon the availability of P” (Plaxton, 2004). Singh and Sale (2000) reported that “P fertilization stimulates root growth, photosynthesis and increases the hydraulic conductivity of roots”. “Phosphorus is used in numerous molecular and biochemical plant processes, particularly in energy acquisition, storage and utilization” (Epstein and Bloom, 2005). “The phosphorus content per unit dry weight is usually considerably higher in the nodules than in the roots and shoots, particularly at low external phosphorus supply. Nitrogen fixing plants have an increased requirement for P over dose receiving direct nitrogen fertilization, probability due to the need for nodule development and signal transduction, and to P-lipids in a large number of bacteroids” (Graham and Vance, 2000). “It acts as a catalyst in several biochemical reactions occurred in plants. It plays an important role in capturing and converting solar energy into useful plant compounds. These compounds help in the general health and vigor of plants” (Griffith, B. 2010). “Legumes are heavy feeders of phosphorus and less responsive to nitrogen because of their capacity to meet their own nitrogen requirement through symbiotic fixation” (Kumar *et al.* 2016). “Phosphorus is connected with some particular plant growth factors are root development, vigorous stem, enhanced flower formation and seed production, earlier and more uniform crop maturity, increase nitrogen fixing capacity of legumes, improvement in crop quality and resistance to plant diseases” (Rehan *et al.* 2018). “It is required for higher and more sustainable production of grain legumes. Generally, legumes have higher P requisites due to more energy consumption in the symbiotic nitrogen fixation process” (Islam *et al.* 2012).

“Phosphorus enhances the activity of rhizobia and increases the formation of root nodules thereby helping in fixing more atmospheric nitrogen in root nodules. Phosphorus is also an important fertilizer in chickpea production; it is a very important chemical fertilizer that can raise the water-holding capacity of the soil” (Dotaniya *et al.*, 2014).

Zinc is required for the proper functioning of various metabolic processes. It's necessary for chlorophyll and carbohydrate production. Several enzyme systems, auxin and protein synthesis, seed formation and maturity rate all require zinc, either directly or indirectly. Zinc is known to help with RNA synthesis, which is required for protein production. In the plant, zinc is not translocated. As a result, symptoms occur first on the plant's younger leaves and other sections. Stunted growth, the formation of light green yellowish patches and chlorotic bands on either side of the midrib in the plants are all common symptoms of zinc deficiency. Sustainable production needs a balanced supply of soil along with suitable physical and biological properties to attain a better growth of roots and efficient utilization of nutrients from the rhizosphere.

The soils of Kanpur are alkaline in nature, low in organic content in generally low in fertility status. All these factors lead to the deficiency of zinc in soils. As such, that the application of zinc may be helpful in increasing the yields of chickpeas under agro-climatic conditions of Kanpur.

Resources and Methods

Experimental Site

The experiment was conducted during the *rabi* season of 2021-22 at the student's Instructional farm, C.S.A. University of Agriculture and Technology, Kanpur Nagar (U.P.). The field was well leveled and irrigated by a tube well. The farm is situated at the main campus of the university, in the west-northern part of Kanpur city under the sub-tropical zone in vth agroclimatic zone (central plain zone).

Edaphic Condition

The soil was moist, well drained with uniform plane topography. The soil of the experimental field was alluvial in origin, sandy loam in texture and slightly alkaline in reaction having pH 7.9 (1:2.5 soil: water suspension method given by Jackson, 1973), electrical conductivity 0.30 dSm⁻¹ (1:2.5 soil: water suspension method given by Jackson, 1973), Organic carbon percentage in the soil is 0.45 per cent (Walkley and Black's rapid titration method given by Walkley and Black,

1934), with available nitrogen 210 kg ha⁻¹ (Alkaline permanganate method given by **Subbiah and Asija, 1956**), available phosphorus as sodium bicarbonate-extractable P was 12.8 kg ha⁻¹ (Olsen's calorimetrically method, **Olsen et al., 1954**) available potassium was 198 kg ha⁻¹ (Flame photometer method given by **Hanwey and Heidel, 1952**), available and available zinc was 0.55 kg ha⁻¹ (DTPA extraction method given by (**Lindsay and Norvell, 1978**).

Detail of treatments and design

The 16 treatments combination of nutrient management practices of inorganic fertilizer (DAP and zinc sulphate). The experiment was laid out in a factorial randomized block design with three replications.

Table -1: detail of the treatment combinations:

S. No.	Symbol	Treatment combination
1.	T ₁	P ₀ Zn ₀
2.	T ₂	P ₀ Zn ₃
3.	T ₃	P ₀ Zn ₆
4.	T ₄	P ₀ Zn ₉
5.	T ₅	P ₃₀ Zn ₀
6.	T ₆	P ₃₀ Zn ₃
7.	T ₇	P ₃₀ Zn ₆
8.	T ₈	P ₃₀ Zn ₉
9.	T ₉	P ₆₀ Zn ₀
10.	T ₁₀	P ₆₀ Zn ₃
11.	T ₁₁	P ₆₀ Zn ₆
12.	T ₁₂	P ₆₀ Zn ₉
13.	T ₁₃	P ₉₀ Zn ₀
14.	T ₁₄	P ₉₀ Zn ₃
15.	T ₁₅	P ₉₀ Zn ₆
16.	T ₁₆	P ₉₀ Zn ₉

Crop Husbandry

Pre-sowing irrigation (Paleva) was done in the experimental field with an object to get optimum moisture conditions for attaining good germination. At proper tilth, one ploughing with tractor drawn mould board plough was done followed by two ploughings by cultivator. Full dose of Phosphorus, and Zinc were applied as basal at the time of sowing in the form of DAP and Zinc sulphate respectively. The sowing of seeds of Chickpea variety RGV-203 was done by line sowing by hand at 4-5 cm depth of soil and with line to line spacing of 45 cm to maintain uniform plant population.

Harvesting and threshing: The crop was harvested at maturity and was allowed to dry in sun. Separate bundles were made for each plot and weighted. The after drying harvest was threshed manually.

Data Collection

Grain yield

After threshing the grain yield from each plot was separately weighed and recorded after converting it into quintals per hectare.

Straw yield

After subtracting the grain yield per plot from the total biological yield. After converting the yields into quintals per hectare, yields were recorded.

Biological yield ($q\ ha^{-1}$)

Seed yield and Stover yield together were regarded as biological yield. The biological yield was calculated with the following formula:

$$\text{Biological yield} = \text{Seed yield} + \text{Stover yield}$$

Harvest index (%):

The recovery of grains in total dry matter was considered as the harvest index, expressed in percentage.

It has been calculated by following formula:

$$\text{Harvest Index (\%)} = [\text{Seed Yield (q ha}^{-1}\text{)} / \text{Biological Yield (q ha}^{-1}\text{)}] \times 100$$

Protein content (%):

Nitrogen content (%) in grains was determined by Kjeldahl's method. The Protein content of the chickpea seed was determined by multiplying the N content of the chickpea seed with a factor of 6.25 ($N\ \% \times 6.25$). (A.O.A.C., 1970).

Statistical analysis: The growth parameters and yields were recorded and analysed as per Gomez and Gomez (1984) the tested at a 5% level of significance to interpret the significant differences.

Result and Discussion

Yield Components

At glance over the data given in the Table 2 and depicted in Fig.-1 clearly shows that among the yield attributing characters of chickpea such as the number of pods plant⁻¹, the number of grain pod⁻¹ and 100 grain weight (gm) significant increase due to the application of Phosphorus and Zinc. Significantly response on yield components was recorded with T₁₅ [P₉₀ Zn₆] over other treatments. The number of pods plant⁻¹, number of grain pod⁻¹ and 100-grain weight (gm) increased to the magnitude of 45.00 to 61.26, 1.20 to 1.73 and 16.96 to 20.20, respectively. The maximum number of pods plant⁻¹ (61.26), number of grain pod⁻¹ (1.73) and 100-grain weight (20.20 gm) was associated with the treatment T₁₅ [P₉₀ Zn₆]. Minimum number of pods plant⁻¹ (45.00), number of grain pod⁻¹ (1.20) and 100 grain weight (16.96 gm) was associated with the treatment T₁ [Control]. The results of the present investigation are also in agreement with the findings of Yadav *et al.* (2022), Singh *et al.* (2022), Kumar *et al.* (2022), Yadav *et al.* (2022) and Sachan *et al.* (2022)

Table-2: Effect of different treatment combinations on yield components of chickpea

Treatment	No of pods plant ⁻¹	No of grains pod ⁻¹	100 Grains weight (g)
T ₁	45.00	1.20	16.96
T ₂	47.21	1.27	17.42
T ₃	52.21	1.36	17.65
T ₄	48.43	1.31	17.53
T ₅	50.12	1.32	17.61
T ₆	54.31	1.48	17.71
T ₇	55.18	1.49	17.87
T ₈	53.42	1.43	17.68
T ₉	56.21	1.52	17.94
T ₁₀	59.87	1.62	18.12
T ₁₁	60.89	1.71	19.82
T ₁₂	58.13	1.57	18.10
T ₁₃	57.32	1.55	18.03
T ₁₄	60.38	1.69	19.11
T ₁₅	61.26	1.73	20.20
T ₁₆	60.12	1.65	18.42

S.E (m) ±	P	0.055	0.34	0.010
	Zn	0.055	0.49	0.010
	P × Zn	0.110	0.59	0.019
CD at 5%	P	0.159	1.03	0.028
	Zn	0.159	1.48	0.028
	P × Zn	NS	NS	NS




Fig.-1: Effect of different treatment combinations on yield components of chickpea Productivity Parameters

It is visualized from the data given in Table-3 and depicted in Fig.-2 clearly indicate that among the productivity parameters viz. grain yield (q ha^{-1}), straw yield (q ha^{-1}), biological yield (q ha^{-1}) and harvest index (%) significantly increase due to the application of Phosphorus and Zinc. Grain yield varied from 12.25 to 18.85 q ha^{-1} , straw yield varied from 17.16 to 23.35 q ha^{-1} , biological yield varied from 29.41 to 42.20 q ha^{-1} and harvest index varied from 41.65 to 44.67 %. The maximum grain yield (18.85 q ha^{-1}), straw yield (23.35 q ha^{-1}), biological yield (42.20 q ha^{-1}) and harvest index (44.67 %) were associated with the treatment T₁₅ [P₉₀Zn₆] during the experimentation. The minimum grain yield (12.25 q ha^{-1}), straw yield (17.16 q ha^{-1}), biological yield (29.41 q ha^{-1}) and harvest index (41.65 %) were under the treatment T₁ [control] during the experimentation. “The surge in seed and straw yields under adequate nutrients supply might be attributed mainly to the collective effect of a greater number of pod plant⁻¹, the number of grains pod⁻¹ and higher test weight, which was the result of improved translocation of photosynthates from source to sink ultimately yield is increased. The increase in grain yield under adequate

nutrient supply mainly due to more yield attributes ultimately resulted in more grain yield. Grain, straw yield, biological yield and harvest index of chickpeas significantly increased due to the application of P 90 (kg ha⁻¹) and Zinc 6 (kg ha⁻¹) over their controls” [Pal, et al.2021]. These results also confirms the findings of **Singh et al. (2021)**, **Pal et al. (2021)**, **Yadav et al. (2022)**, **Kumar et al. (2022)** and **Sachan et al. (2022)**

Table 3: Effect of different treatment combinations on productivity parameters of chickpea

Treatment		Grain yield (q ha ⁻¹)	Straw yield (q ha ⁻¹)	Biological yield (q ha ⁻¹)	Harvest index (%)
T₁		12.25	17.16	29.41	41.65
T₂		12.52	17.35	29.87	41.91
T₃		13.76	18.72	32.48	42.36
T₄		12.87	17.81	30.68	41.95
T₅		13.24	18.25	31.49	42.05
T₆		14.86	19.65	34.51	43.05
T₇		15.34	20.12	35.46	43.26
T₈		14.25	19.10	33.35	42.73
T₉		15.77	20.46	36.23	43.53
T₁₀		16.96	21.78	38.74	43.78
T₁₁		18.24	22.96	41.20	44.27
T₁₂		16.71	21.34	38.05	43.91
T₁₃		16.10	20.95	37.05	43.45
T₁₄		17.65	22.54	40.19	43.92
T₁₅		18.85	23.35	42.20	44.67
T₁₆		17.31	22.24	39.55	43.76
S.E (m) ±	P	0.022	0.021	0.043	0.010
	Zn	0.022	0.021	0.043	0.010
	P×Zn	0.044	0.043	0.087	0.019
CD at 5%	P	0.064	0.062	0.126	0.028
	Zn	0.064	0.062	0.126	0.028
	P×Zn	NS	NS	NS	NS




Fig.-2: Effect of different treatment combinations on productivity parameters of chickpea

Quality parameter

Protein

It is visualized from the data given in Table 4 clearly indicates that among the quality parameters viz. protein content increase due to the application of Phosphorus and Zinc. Protein content varied from 19.56 to 20.93 (%) The maximum protein content 20.93 % was associated with the treatment T₁₅ [P₉₀Zn₆]. The minimum protein content 19.53% was under the treatment T₁ [control]. Similar findings were reported by Ahmed *et al.* (2003), Vikrant *et al.* (2005) and Tripathy *et al.* (2019)

Table-4: Effect of different treatment combinations on quality parameters of chickpea

Treatment	Treatment combination	Protein content in grain (%)
T ₁	P ₀ Zn ₀	19.56
T ₂	P ₀ Zn ₃	19.68
T ₃	P ₀ Zn ₆	19.94
T ₄	P ₀ Zn ₉	19.81
T ₅	P ₃₀ Zn ₀	19.87
T ₆	P ₃₀ Zn ₃	20.18
T ₇	P ₃₀ Zn ₆	20.25

T₈	P₃₀Zn₉	20.06
T₉	P₆₀Zn₀	20.37
T₁₀	P₆₀Zn₃	20.62
T₁₁	P₆₀Zn₆	20.87
T₁₂	P₆₀Zn₉	20.56
T₁₃	P₉₀Zn₀	20.44
T₁₄	P₉₀Zn₃	20.75
T₁₅	P₉₀Zn₆	20.93
T₁₆	P₉₀Zn₉	20.68
S.E (m) ±	P	0.34
	Zn	0.29
	P×Zn	0.49
CD at 5%	P	1.03
	Zn	1.03
	P×Zn	NS

Conclusion

The current study demonstrates the benefit of the application of phosphorus 90 kg ha⁻¹ and zinc 6 kg ha⁻¹ significantly increased yield attributes, productivity and quality parameters like the number of pod plant⁻¹, number of grains pod⁻¹, test weight of 100 seeds, grain yield, straw yield, biological yield, harvest index and protein quality. Finally it can be concluded that the treatment T₁₅ [P₉₀Zn₆] is the best option for improving productivity, yields and quality parameter of chickpea crops.

References

Ahmed. M. K. A., Afifi, M. H. and Mohamed, M. F. (2003). Effect of biofertilizers, chemical and organic fertilizers on growth, yield and quality of some leguminous crops. *Egyptian Journal of Agronomy*. **25**: 45-52.

Anonymous. (2021). Agricultural Statistics at a Glance 2020. Directorate of Economics & Statistics, Department of Agriculture, Cooperation and Farmers Welfare, Ministry of Agriculture & Farmers Welfare, Govt. of India, New Delhi, p. 63.

Dotaniya ML, Datta SC, Biswas DR, Kumar K. (2014), Effect of organic sources on P_2O_5 fraction and available P_2O_5 in typical Haplustept *J. Indian Soc. Soil Sci.* 62(1):80-83.

Epstein, E. and Bloom, A.J. (2005). *Mineral nutrition of plants: Principles and perspectives*, 2nd ed. Sunderland, Massachusetts: Sinauer Associates.

Gomez, K. A., & Gomez, A. A. (1984). *Statistical procedures for agricultural research*. John Wiley & sons.

Graham, P.H. and Vance, C.P. (2000). Nitrogen fixation in perspective, an overview of research and extension needs. *Field Crops Research* 65: 93-106.

Griffith B (2010). Efficient Fertilizer Use Phosphorus, 1-7.

Hanway, J.J; and Heidel, H. (1952). Soil analysis methods as used in Iowa State College, Soil Testing Laboratory. *Iowa Agriculture* 54: 1-31.

Islam M, Mohsan S, Ali S, Khalid R & Afzal S (2012). Response of chickpea to various levels of phosphorus and sulphur under rainfed conditions in Pakistan. *Romanian Agric Res.* 29: 175-183.

Jackson, M.L. (1973). Soil chemical analysis. Prentice Hall of India Pvt. Ltd, New Delhi.

Kumar, K., Pyare, R., Niwas, R., Tiwari, K., Sachan, R., Pal, R. K., Patel, V. K. & Ranjan, A. R., (2022).. Studies on the Root Architecture with Nodulation of the Chickpea (*Cicer arietinum* L.) as Influence by Different Moisture Management Practices along with Seed Inoculation and Level of Zinc. *International Journal of Environment and Climate Change*, 2896-2904.

Kumar, K., Pyare, R., Verma, V. K., Sachan, R., Niwas, R., Yadav, A., Pal, R. K. & Ranjan, A. R., (2022). Impact of Moisture Conservation Practices, Seed Inoculation and Zinc Level on Growth and Yield of Chickpea (*Cicer arietinum* L.). *International Journal of Plant & Soil Science*, 34(23), 546-556.

Kumar, S., Tripathi, D.K., Bharose, R., Kumar, M. and Kumar, R. (2016). Effect of different fertility levels and micronutrients on nodulation and nutrient uptake by chickpea. *An Asian Journal of Soil Sci.* **11**(1): 62-66.

Lindsay, W. L., & Norvell, W. (1978). Development of a DTPA soil test for zinc, iron, manganese, and copper. *Soil science society of America journal*, **42**(3), 421-428.

Olsen, S.R, Cole, C.V., Watanable, F. S. and Dean, L. A. (1954). Estimation of available phosphorous in soil by extraction with sodium bicarbonate. *USDA, Cric.* **930**:19- 23

Pal, S., Pandey, S. B., Singh, A., Singh, S., Sachan, R., & Yadav, A. (2021). Effect of Phosphorus, Boron and *Rhizobium* inoculation on productivity and profitability of chickpea. *The Pharma Innovation Journal* 2021; 10(12): 1810-1814

Pingoliya, K, K., Dotaniya, M, L., Mathur, A, K. (2013). Role of phosphorus and iron in chickpea (*Cicer arietinum* L.). *Lap Lambert Academic Publisher, Germany.*

Plaxton, W.C. (2004). Plant response to stress: biochemical adaptations to phosphate deficiency. *In: Goodman R (eds.) Encyclopedia of Plant and Crop Science.* New York: Marcel Dekker. pp. 976-980.

Rehan, W., Jan, A., Liaqat, W., Jan, F.M., Ahmadzai, M.D., Ahmad, H., Haroon, J., Anjum, M.M. and Ali, N. (2018). Effect of phosphorous, rhizobium inoculation and residue types on chickpea productivity. *Pure Appl. Biol.*, **7**(4): 1203-1213.

Sachan, R., Pandey, S. B., Kumar R., Singh D., Sharma, S., Kumar. K., Singh, A., Pal, S., Verma, A. K. and Sachan, H. (2022) Effect of Phosphorous, Chloropyriphos and *Rhizobium* Inoculation on Production and Economics of Chickpea (*Cicer aretinum* L.). *International Journal of Plant & Soil Science*, 1390-1398.

Sachan, R., Pandey, S. B., Kumar, A., Pathak, R. K., Pandey, H. P., Singh, A., ... & Kumar, M. (2022). Interaction effect of Phosphorous, Chloropyriphos and *Rhizobium* Inoculation on Growth Characteristics, Yield Components and Productivity Parameters of Chickpea under Central Plain Zone of Uttar Pradesh. *AMA ISSN: 00845841* Volume 53, Issue 08, August, 2022

Schulze, J., Temple, G., Temple, S.J., Beschow, H., and Vance, C.P. (2006). Nitrogen fixation by white lupin under phosphorus deficiency. *Annals of Botany* **98**: 731-740.

Singh, A. K., Dimree, S., Kumar, A., Sachan, R., Sirohiya, A., & Nema, S. (2022). Effect of *rhizobium* inoculation with different levels of inorganic fertilizers on yield, nutrient content & uptake of chickpea (*Cicer arietinum* L.). *Int J Plant Soil Sci*, 34(22), 262-8.

Singh, A., Singh, D., Kumar, R., Pal, S., Sachan, R. & Yadav, A. (2021). Study the effect of organic, inorganic and biofertilizers on nutrients content and uptake of chickpea (*Cicer arietinum* L.). *The Pharma Innovation Journal* 2021; 10(10): 418-423

Singh, D.K. and Sale, P.W.G. (2000). Growth and potential conductivity of white clover roots in dry soil with increasing phosphorus supply and defoliation frequency. *Agronomy Journal* **92**: 868-874.

Subbiah, B.V. and Asija, C.L. (1956). A rapid procedure for the estimation of available N in Soil. *Curr. Sci.* **25**:259-260.

Tripathy, D.K., Kumar, S. and Zaidy, S.F.A. (2019). Effect of phosphorus, sulphur and micronutrients (Zinc and Boron) levels on performance of chickpea (*Cicer arietinum* L.). *Natl. Acad. Sci. Lett.* 10.1007/s40009-019-00802-4.

Vikrant, Singh, H., Malik, C.V.S. and Singh, B.P. (2005). Grain yield and protein content of cowpea as influenced by farm yard manure and phosphorus application. *Indian Journal of Pulses Research* **18** (2): 250-251.

Walkley, A. and Black, C. S.A. (1934). Old piper, S.S. soil and plant analysis. *Soil Sci.* 37:29-38.

Wallace, T. C., Murray, R., Kathleen, M. and Zelman, K. (2016). The nutritional value and health benefits of chickpeas and humus. *Nutrients.* **8**(12): 766.

Yadav, A., Singh, D., Kumar, R., Sachan, R., Kumar, K., Singh, A., & Singh, K. K. (2022). Response of different level of phosphorus, zinc and Rhizobium inoculation on growth yield attributes and yield of Chickpea (*Cicer aretinum* L.). *International Journal of Environment and Climate Change*, 12(11), 1954-1964.

Yadav, P., Yadav, D. D., Pandey, H. P., Yadav, A., Sachan, R., & Yadav, S. (2022). Effect of Fertility Levels and Biofertilizers on Growth Parameters, Root Architecture and Quality of Chickpea (*Cicer arietinum* L.). *International Journal of Plant & Soil Science* 34(17): 61-67, 2022; Article no.IJPSS.86643 ISSN: 2320-7035

Yadav, S., Yadav, D.D., Kumar, A., Sachan, R. & Yadav, S. (2022). Effect of Fertility Levels and Bio-fertilizers Application on Yield, Yield Attributes, and Economics of Chickpea (*Cicer arietinum* L.). *International Journal of Plant & Soil Science* 34(13): 65-69, 2022; Article no.IJPSS.85129 ISSN: 2320-7035

Pal S, Pandey SB, Kumar R, Singh D, Singh A, Singh S. Response of phosphorus, boron and rhizobium inoculation on growth attributes and productivity of chickpea. *The Pharma Innovation Journal* 2021; 10(10): 255-260

UNDER PEER REVIEW