

SERO-PREVALENCE OF SYPHILIS AMONG PREGNANT WOMEN ATTENDING ANTENATAL CLINIC IN DOGON AGOGO PRIMARY HEALTH CARE CENTRE JOS NORTH, PLATEAU STATE, NIGERIA

ABSTRACT

Background: Syphilis remains a public health problem in many parts of the world, including Nigeria, where it is common during pregnancy.

Design and Duration: This is a cross-sectional study involving pregnant women attending antenatal clinic in Dogon Agogo Primary Health Centre, Jos North, from March 2017 to November, 2021.

Aim: The aim of this study is to seroprevalence of Syphilis among pregnant women attending Primary health care centre in Jos North,

Materials and Methods: A total of one thousand, two hundred and nineteen blood samples were screened from 2017 to 2021, using rapid diagnostic tests. Patient's demographic variables, clinical profiles and information considered to be risk factors were also collected using structured questionnaires. Patient's demographic details, clinical profiles and information considered to be risk factors were also collected using structured questionnaires.

Results: The results showed that only 33 (2.7%) were positive. This includes 14(42.4%) patients between the age of 20 – 34 years as the highest frequency, followed by 12(36.4%) in 35 – 49 years of age. Most of the syphilis-positive patients are married 21(63.6%) and had attended tertiary level of education 13(39.4%), but mostly unemployed/Housewives 11(33.3%). The clinical profile of the participants revealed that the majority 14(42.5%) of the women that attended the facility throughout this period are in their third pregnancy and above (Multigravida). They were screened during the first-time visit (72.7%) and first trimester (33.3%). The pregnancy outcome among the seropositive patients showed that many 14(42.4%) undergo normal delivery, but 10(30.3%) had miscarriages. Blood transfusion (81.8%), multiple sex partners (75.8%), unprotected sexual intercourse with a person of unknown status (57.6%), and previous exposure to sexually transmitted infections (84.8%) were the most prevalent risks factors observed. The overall prevalence of syphilis across the five years of sampling was 2.71% (33 of 1219), where the highest number of cases 57.6% was recorded in the year 2020, followed by 18.2% in 2018, with 3.0% in 2019 as the least.

Conclusion: This study found that the prevalence of syphilis over the five-year period among pregnant women in this area is low. However, the study advocates the improvement of screening programs during pregnancy as part of routine antenatal care.

Keywords: Syphilis, Pregnancy, Antenatal, Multigravida, Miscarriage. Blood transfusion

INTRODUCTION

“Syphilis is a sexually transmitted disease caused by the bacterium *Treponema pallidum* [1]. The signs and symptoms of syphilis vary depending in which of the four stages it presents (primary, secondary, latent, and tertiary)” [1]. “The primary stage classically presents with a single chancre (a firm, painless, non-itchy skin ulceration usually between 1 cm and 2 cm in diameter) though there may be multiple sores” [1,2]. “In secondary syphilis, a diffuse rash occurs, which frequently involves the palms of the hands and soles of the feet [2]. There may also be sores in the mouth or vagina” [1]. “In latent syphilis, which can last for

years, there are few or no symptoms” [1]. “In tertiary syphilis, there are gummas (soft, non-cancerous growths), neurological problems, or heart symptoms” [3]. “Syphilis has been known to cause symptoms similar to many other diseases” [1,4].

“Syphilis is most commonly spread through sexual activity” [5]. “It may also be transmitted from mother to baby during pregnancy or at birth, resulting in congenital syphilis” [5,6]. “Common risk factors for syphilis include unprotected sexual activity involving contact with oral, genital mucosa or anus” [2]. “Sexual contact with a known case of syphilis, sex with someone from a country/region with a high prevalence of syphilis, previous syphilis, HIV infection or other STDs, born to a person diagnosed with infectious syphilis in pregnancy, member of a vulnerable population. Other activities that may increase potential exposure to STIs include anonymous sexual partnering, street involvement and drug abuse” [2].

“Syphilis can be transmitted by direct contact with an infectious lesion or through vertical transmission during pregnancy. Primary, secondary and early latent stages are considered infectious, with an estimated risk of transmission per partner of 51% to 64%” [3]. “Early latent syphilis is considered infectious because of the 25% chance of relapse to the second stage” [4]. “The primary mode of syphilis transmission is by vaginal, anal and oral sexual contact. Other routes of transmission (e.g. kissing and needle-sharing) are rare” [5].

Transplacental transmission can occur as early as nine (9) weeks gestation [6] and throughout pregnancy. “Most infants with congenital syphilis are infected in utero, but they can also be infected by contact with an active genital lesion at the time of delivery. The risk of transmission is related to both stages of syphilis as well as gestational age at the time of diagnosis and treatment, with the greatest risk of transmission in pregnant people acquiring syphilis near term” [7]. “The risk of transmission to the foetus is 70% to 100% in untreated pregnant people with primary or secondary syphilis and 40% with early latent syphilis” [8,9]. “In a recent systematic review, adverse pregnancy outcomes occurred in 76.8% of untreated pregnancies compared to 13.7% of uninfected pregnancies” [10]. “The clinical manifestations of syphilis are usually described according to the stage of the disease: primary, secondary, latent, and congenital, neurosyphilis [which may occur at tertiary stage of syphilis](#). Medical practitioners should consider syphilis in people presenting with rashes or genital ulcerative disease and/or proctitis” [11].

“Congenital syphilis is considered a severe public health problem because it accounts for approximately 40% of the perinatal mortality rates, 25% of stillbirths, and 14% of neonatal deaths, in addition to causing severe consequences for the foetus. Syphilis treatment is easy to access and simple to perform and therefore, its management during pregnancy should be straightforward. The diagnosis and treatment should be performed in a timely manner in the early period of pregnancy. However, being a sexually transmitted infection (STI) makes it a difficult situation to approach, especially when experienced during pregnancy” [12] (Rêgo *et al.*, 2020).

“Syphilis is transmitted via sexual contact or from mother to child during pregnancy or at delivery” [9]. “In pregnant women syphilis might also increase the risk of mother-to-child transmission of HIV in cases where mothers are coinfecting” [10]. “Syphilis remains a major cause of reproductive morbidity and poor pregnancy outcomes in developing countries. Syphilis in pregnant women can result in adverse outcomes of pregnancy in up to 80% of cases, such as stillbirth and spontaneous abortion (40%), perinatal death (20%), and serious neonatal infections and low-birth weight babies (20%)” [as reported by Shazia *et al.* \[13\]](#).

“It was reported that more than ten million people are infected with syphilis worldwide; with majority found in sub-Saharan Africa and Asia” [14]. “Every year around two million pregnant

women are estimated to have active syphilis where only few were diagnosed and treated. The prevalence of syphilis in pregnant women was estimated to be 2.7% in sub-Saharan Africa, which represents nearly 1 million pregnancies to be at risk annually” [15]. “Poor pregnancy outcomes were reported more than four times higher in untreated cases [16], where about 90% of these infections occur in areas with poor social amenities” [17].

“Information regarding syphilis infection in pregnancy in Nigeria shows a wide geographical variation in seroprevalence. Several models have been proposed to estimate adverse pregnancy outcomes in women infected with syphilis, with resulting estimates ranging from 50% to 80%” [18]. “Transmission occurs more commonly in the last two trimesters, but the spirochete can cross the placenta at any time during pregnancy” [19]. Foetal death and morbidity due to congenital syphilis are preventable if the infected mother is identified and treated appropriately by the middle of the second trimester. Because of the serious complications of syphilis in pregnancy, WHO has recommended universal antenatal screening and recommended screening for syphilis at the first antenatal visit, as early as possible in pregnancy, repeating in the third trimester if resources permit, to detect infection acquired during pregnancy. The effectiveness of such antenatal syphilis screening and its treatment for the prevention of adverse pregnancy outcomes is of paramount importance.

Prevalence of syphilis infections among pregnant mothers in Nigeria differs between regions depending on a number of factors such as the overall HIV prevalence, culture of the general population, level of awareness and associated risk factors. To date there are few studies conducted in [Nigeria \[1\]](#) as a result, there is scarcity of information about the true burden and determinant factors of syphilis among pregnant women in the study area. Therefore, this study was aimed to determine the seroprevalence of Syphilis among pregnant women attending the Antenatal clinic at Dogon Agogo PHC, in Jos North.

MATERIALS AND METHODS

Study Design and Population

This study is [cross-sectional](#) survey research involving a total of One thousand two hundred and nineteen (1219) pregnant women that attended the Antenatal clinic in Dogon Agogo PHC Centre from 2017 to 2021.

Sample and Data collection

A fingerstick blood specimen was aseptically collected from each pregnant woman. An information sheet was used to record the participant’s name, age, time and date of collection/screening test. Data on patient’s demographic variables, clinical profiles and information considered to be risk factors were also collected using structured questionnaires after their consent were sought, with ethical approval from the health centre’s management.

Screening test for Syphilis

A hanging drop of fingerstick whole blood specimen was allowed to fall into the centre of the specimen well (S), then 2 drops of buffer was added on it. The test kit was laid flat on a clean, dry surface and waited for the coloured band to appear, within 15-20 minutes to read the results [20].

Statistical Analysis

The chi-square test was employed to determine the relationship between the patient’s socio-demographic variables and clinical profile/risk factors with syphilis infection. P values of < 0.05 were considered to be statistically significant.

RESULTS

The results of seroprevalence of Syphilis among pregnant women attending Primary health care centre Dogon Agogo showed that out of one thousand two hundred and nineteen blood samples tested from 2017 to 2021, only 33 (2.7%) were positive. This includes 14 (42.4%) patients between the aged of 20 – 34 years as the highest frequency, followed by 12 (36.4%) in 35 – 49 years of age. Most of the syphilis-positive patients in our study are married 21(63.6%), attended tertiary level of education 13 (39.4%), but mostly unemployed/Housewives 11 (33.3%).

Table 1: Prevalence of syphilis among pregnant women according to demographic detail Patients

| Demographic profile | Number of Women Screened (N=1219) | Number of Reactive cases (n=33) | Prevalence (%) |
|-------------------------------|------------------------------------------|----------------------------------------|-----------------------|
| Age (Years) | | | |
| 15 – 19 | 143 | 04 | 12.1 |
| 20 – 34 | 463 | 14 | 42.4 |
| 35 - 49 | 486 | 12 | 36.4 |
| 50 and above | 127 | 03 | 9.1 |
| Marital status | | | |
| Married | 1115 | 21 | 63.6 |
| Single | 104 | 12 | 36.4 |
| Divorced | 00 | 00 | 0.0 |
| Widowed | 00 | 00 | 0.0 |
| Educational background | | | |
| Primary | 402 | 11 | 33.3 |
| Secondary | 657 | 09 | 27.3 |
| Tertiary | 160 | 13 | 39.4 |
| Occupation | | | |
| Civil servant | 132 | 09 | 27.3 |
| Business | 82 | 06 | 18.2 |
| House wife | 384 | 11 | 33.3 |
| Private | 161 | 01 | 3.0 |
| Daily labourer | 460 | 06 | 18.2 |

Table 2: Distribution of syphilis among pregnant women according to clinical profile

| Clinical profile | Number of Women Screened (N=1219) | Number of Reactive cases (n =33) | Prevalence (%) |
|-------------------------|------------------------------------------|-----------------------------------------|-----------------------|
| Parity | | | |
| Primigravida | 395 | 11 | 33.3 |
| Biogravida | 102 | 08 | 24.2 |
| Multigravida | 722 | 14 | 42.5 |
| Antenatal visit | | | |
| First visit | 729 | 24 | 72.7 |
| Second visit | 286 | 07 | 21.2 |

| | | | |
|---------------------------|-----|----|------|
| Third visit | 117 | 02 | 6.1 |
| Fourth visit | 62 | 00 | 00 |
| Fifth and above visit | 25 | 00 | 00 |
| Gestational Period | | | |
| 1 st trimester | 402 | 13 | 39.4 |
| 2 nd trimester | 621 | 09 | 27.3 |
| 3 rd trimester | 196 | 11 | 33.3 |
| Pregnancy outcome | | | |
| Normal delivery | 893 | 14 | 42.4 |
| Caesarean section | 87 | 04 | 12.1 |
| Low birth weight | 65 | 02 | 6.1 |
| Pre-term delivery | 26 | 00 | 0.0 |
| Still birth | 13 | 00 | 0.0 |
| Perinatal death | 22 | 03 | 9.1 |
| Miscarriage | 113 | 10 | 30.3 |

The clinical profile of the participants in this study (Table 2) revealed that the majority 14 (42.5%) of the pregnant women that attended the facility throughout this period are in their third pregnancy and above (Multigravida). They were screened during the first-time visit (72.7%) and first trimester (33.3%). Analysis of pregnancy outcomes among the seropositive patients showed that the majority of 14 (42.4%) undergo normal delivery, but 10 (30.3%) had a miscarriage.

Table 3: Distribution of syphilis among pregnant women according to Risk factors

| Clinical profile | Number of Women Interviewed (N=1219) | Number of Reactive cases (n =33) | Prevalence (%) |
|------------------------------------------------------|--------------------------------------|----------------------------------|----------------|
| History of blood transfusion | | | |
| Yes | 395 | 27 | 81.8 |
| No | 824 | 06 | 18.2 |
| Multiple sex partner | | | |
| Yes | 00 | 25 | 75.8 |
| No | 1219 | 08 | 24.2 |
| Unprotected sex with person of unknown status | | | |
| Yes | 26 | 19 | 57.6 |
| No | 1193 | 14 | 42.4 |
| History of STIs | | | |
| Yes | 196 | 28 | 84.8 |

| | | | |
|----|------|----|------|
| No | 1023 | 05 | 15.2 |
|----|------|----|------|

The distribution of syphilis based on risk factors in this study (Table 3) indicated that blood transfusion (81.8%), association with multiple sex partners (75.8%), unprotected sexual intercourse with person of unknown status (57.6%) and previous exposure to sexually transmitted infections (84.8%) are the most prevalent risk factors observed among the pregnant women interviewed.

Table 4: Distribution of syphilis among pregnant women base on the year of sampling

| Year | Number of Women Screened (N=1219) | Number of Syphilis cases (n=33) | Prevalence (%) |
|------|-----------------------------------|---------------------------------|----------------|
| 2017 | 78 | 02 | 6.0 |
| 2018 | 173 | 06 | 18.2 |
| 2019 | 231 | 01 | 3.0 |
| 2020 | 285 | 19 | 57.6 |
| 2021 | 452 | 03 | 15.2 |

In this study (Table 4), the overall prevalence of syphilis across the five years of sampling was 2.71% (33 of 1219), where the highest number of cases 57.6% (19 of 33) was recorded in the year 2020, followed by 18.2% (6 of 33) in 2018, with 3.0% (1 of 33) in 2019 as the least.

DISCUSSION

Syphilis is one of the most significant venereal diseases of public health importance, associated with poor pregnancy outcomes, such as miscarriage, low birth weight and stillbirth. In Nigeria, the prevalence of syphilis among pregnant women varies and depends on geographical locations [21]. The disease has continued to exert a high burden worldwide especially in sub-Saharan Africa, despite that its diagnosis, prevention, and control of pregnancy is highly recommended. Timely ante-natal visits and testing play a vital role in the early diagnosis of syphilis [22]. However, the actual burden of syphilis infection during pregnancy is not well known, especially in developing countries where adequate health facilities are poor or unavailable, and most pregnant women attended antenatal clinics late [23, 24].

In this study, the results showed that the overall prevalence of Syphilis among pregnant women across the five years of sampling was 2.7% which is higher than the rate (2.0%) obtained by Lamidi and Gbemisola (2021) [25] in Kogi, Middle of Nigeria. The rate of infection among women was found in the active age group of 20 years and above, even in those who are single. This may be due to the fact that they have more than one sexual partner and are not aware of the preventive measures. This frequency was higher than the National Average of 0.3% for Syphilis among pregnant women in Nigeria [25]. These cases might result in a severe impact on pregnancy outcomes, such as spontaneous abortion, stillbirth and vertical transmission resulting in congenital syphilis.

The prevalence of 2.7% in our study was also higher than 0.4% reported by Olokoba *et al.* (2008) [26] in Northeastern, Nigeria; 0.13% by Ozumba *et al.* (1999) [27] in Enugu, Southeastern, Nigeria; 1.7% obtained by Aboyeji and Nwaburi (2003) [28] in Ilorin, North

Central, Nigeria. A lower prevalence of 0.10% and 0.98% was reported by Shazia *et al.* (2012) [29] among pregnant women in a rural area of India and Mbamara *et al.* (2011) [30] respectively, A 1.9% was also found by Isa *et al.* (2014) [31] in Maiduguri and 0.2% by Gao *et al.* (2009) [32] in China.

The higher rate of syphilis reported in this study may be associated with a lack of awareness among pregnant women. In contrast, the prevalence of 2.7% reported in this study is lower than the 2.97% reported by Taiwo *et al.* (2007) [33] in Osogbo, Southwestern, Nigeria; 5.8% reported by Buseri *et al.* (2010) [34] in South-South, Nigeria; 4.3% reported by Creek *et al.* (2005) [35] in Botswana; 2.2% reported by Kirakaya, *et al.* (2010) [36], 8.0% in Ouagadougou; 12.5% reported by Ratnam, *et al.* (1982) [37] among pregnant women in Zambia; 18.3% by Lindstrand *et al.* (1993) [38] 10.0% among antenatal women in Mozambique and 5.0% reported by Kwiek *et al.* (2008) [39] in Malawi. In the present study, the Seropositive prevalence in the active age groups, indicated that these groups are actively reproductive as such have a high risk of infection. Therefore, routine screening for syphilis during an antenatal visit is hereby encouraged.

A 20 years systematic review and meta-analysis of syphilis in sub-Saharan Africa among 175,546 pregnant women, from January 1999 to November 2018 using different data sources was conducted by Hussen and Tedesse (2019) [40]. The pooled prevalence of syphilis among pregnant women in the region was 2.9%, [which slightly agrees with our findings \(2.7%\)](#). East and Southern African regions had a higher syphilis prevalence among pregnant women (3.2%) and (3.6%), respectively) [40]. The prevalence of syphilis among pregnant women in most parts of the region seemed to have decreased over the past 20 years except for the East African region [40]. However, prevalence did not significantly differ by region and time period [41].

In our study, blood transfusion, multiple sex partners, unprotected sexual intercourse with a person of unknown status, and previous exposure to sexually transmitted infections are the most frequent risk factors observed in the area. This can be attributed to cultural practice and diversity in the area, where marriage is not strictly a pre-requisite to bearing pregnancy/child. Some of the women give birth with profuse loss of blood or being anaemic, due to improper prenatal care. This result is contrary to the findings of Isa *et al.* (2014) [31] Maiduguri, where zero prevalence of Syphilis was recorded among pregnant women having a history of blood transfusion.

A study by Lumbiganon *et al.* (2002) [42] in Thailand evaluated the magnitude, risk factors and outcomes of syphilis in pregnancy. All women attending the first antenatal care at each selected clinic were enrolled. The women attending prenatal care for the first time after the date of the start of the study at each of the 53 selected clinics, regardless of their gestational age, medical or obstetrical characteristics or previous ANC, were enrolled in the trial. Screening at the first antenatal visit was routinely performed. All women also had the same syphilis tests after delivery. The initial prevalence, the incidence during pregnancy and the overall prevalence of syphilis at delivery were 0.9%, 0.4% and 1.3% respectively [42]. Risk factors for syphilis during pregnancy were found in younger age for the incidence and older age and those with a history of stillbirth for the prevalence. Women with syphilis during pregnancy had significantly more adverse outcomes [43].

A prospective cross-sectional study in three teaching hospitals in Addis Ababa was carried out by Kebede and Chamiso (2000) [44] on a total of 410 pregnant women attending antenatal care within six months. The main outcome measures considered were seropositivity for syphilis and socio-demographic factors related to it. Among the study population, twelve women (2.9%) positive for the disease were found among those with lesser monthly income, as observed in our

study, where the past history of abortion was significantly associated with seropositive cases ($p < 0.05$). It is, therefore, necessary to conduct a large-scale study to evaluate if screening is cost-effective and establish risk scoring methods.

The rates of sexually transmitted infections have increased since the turn of the millennium in many countries, often in association with HIV/AIDS [45]. This rise has been attributed partly to unsafe sexual practices among homosexuals, increased rate of promiscuity, prostitution, and inconsistent use of condoms. In the present study, the highest cases of 57.6% (19 of 33) of syphilis were recorded in the year 2020. This was ascribed to the COVID-19 pandemic, resulting in lock-down, and posing challenges in securing means of livelihood. During this period, there are many cases of sexual abuse, that can increase the rate of STIs in the area. Adverse effects on the foetus and infant due to maternal syphilis were among the complications reported in many studies [46,47,48]. The incidence of such outcomes is more severe in areas of a high prevalence of the infection, where latent or early syphilis is likely to be common among women.

CONCLUSION

This study revealed that the prevalence of syphilis over the five-year period among pregnant women is low in this area. However, the findings advocate improving the screening program during pregnancy as part of the routine antenatal care schedule. Primary prevention of syphilis in pregnant women should also be highly encouraged. The present study also recommended that in addition to the initial testing, a second routine test for syphilis should be established early even in the third trimester.

Ethical Approval:

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

Consent

As per international standard or university standard, Participants' written consent has been collected and preserved by the author(s).

REFERENCE

1. Olokoba AB, Salawu FK, Danburam A, Desalu OO, Midala JK, Badung LH, Olokoba LB, Abdulkarim A. Syphilis in pregnant Nigerian women: Is it still necessary to screen. *Eur. J. Sci. Res.* 2009; 29(3):315-19.
2. Kent ME, Romanelli F. "Reexamining syphilis: an update on epidemiology, clinical manifestations, and management". *Ann. Pharm.* 2008; 42 (2): 226–36.
- Singh AE, Romanowski B. Syphilis: review with emphasis on clinical, epidemiologic, and some biologic features. *Clin Microbiol Rev.* 1999;12(2):187-209.
3. Choudhri Y, Miller J, Sandhu J, Leon A, Aho J. Infectious and congenital syphilis in Canada, 2010-2015. *Can Commun Dis Rep.* 2018;44(2):43-48.
4. Stoltey JE, Cohen SE. Syphilis transmission: A review of the current evidence. *Sexual health.* 2015;12(2):103-109.
5. Edwards S, Carne C. Oral sex and transmission of non-viral STIs. *Sex Transm Infect.* 1998;74(2):95-100.
6. De Santis M, De Luca C, Mappa I, et al. Syphilis infection during pregnancy: Foetal risks and clinical management. *Infect Dis Obstet Gynecol.* 2012; 20(12), 34-42.
7. Finelli L, Berman SM, Koumans EH, Levine WC. Congenital syphilis. *Bull World Health Organ.* 1998;76 Suppl 2:126-128.

8. Qin J, Yang T, Xiao S, Tan H, Feng T, Fu H. Reported estimates of adverse pregnancy outcomes among women with and without syphilis: A systematic review and meta-analysis. *PLoS One*. 2014;9(7):e102203.
9. Karp G, Schlaeffer F, Jotkowitz A, Riesenber K. Syphilis and HIV co-infection. *Eur. J Intern Med*. 2009;20(1):9-13.
10. Zetola NM, Klausner JD. Syphilis and HIV infection: An update. *Clin. Infect. Dis*. 2007;44(9):1222-1228.
11. Golden MR, Marra CM, Holmes KK. Update on syphilis: Resurgence of an old problem. *JAMA*. 2003; 290(11):1510-1514.
12. Rego AS, Costa LC, Rodrigues LS, Garcia RA, Silva FMA, Junior AD and Rodrigues LS Congenital syphilis in Brazil: distribution of cases notified from 2009 to 2016. *J. Brazilian Soc. Trop. Med*. 2020, 53; e20200338.
13. Woods CR. "Congenital syphilis-persisting pestilence". *Paediatr. Infect. Dis. J*. 2009; 28 (6): 536–37.
14. Newman, L; Rowley, J; Vander Hoorn, S; Wijesooriya, NS; Unemo, M; Low, N; Stevens, G; Gottlieb, S; Kiarie, J; Temmerman, M (2015). "Global Estimates of the Prevalence and Incidence of Four Curable Sexually Transmitted Infections in 2012 Based on Systematic Review and Global Reporting". *PLOS ONE*. 10 (12): e0143304.
15. Franzen C, "Syphilis in composers and musicians – Mozart, Beethoven, Paganini, Schubert, Schumann, Smetana". *Eur. J. Clin. Microbiol. & Infect Dis*. 27 (12): 1151–57.
16. Coffin LS, Newberry A, Hagan H, Cleland CM, Des Jarlais DC, Perlman DC, "Syphilis in Drug Users in Low- and Middle-Income Countries". *The Int. J. Drug Policy*. 2010; 21 (1): 20–27.
17. Gao L, Zhang L, Jin Q. "Meta-analysis: prevalence of HIV infection and syphilis among MSM in China". *Sexually Trans. Infect* 2009; 85 (5): 354–58.
18. Karp G, Schlaeffer F, Jotkowitz A, Riesenber K. "Syphilis and HIV co-infection". *Eur J. Inter. Med*. 2009; 20 (1): 13.
19. Lynn, W. A. and Lightman, S. "Syphilis and HIV: a dangerous combination," *The Lancet Infect. Dis.*, 2004; 4(7) 456–466.
20. Taiwo SS, Adesiji YO, Adekanle DA. Screening for syphilis during pregnancy in Nigeria: A practice that must continue. *Sex Transm. Infect*. 2007; 83:357-58.
21. Arnesen L, Martinez G, Mainero L, Serruya S, Dur'an P. "Gestational syphilis and stillbirth in Latin America and the Caribbean," *Int. J. Gynec and Obst*, 2015; 128 (3), 241–245.
22. Cameron CE and Lukehart SA "Current status of syphilis vaccine development: Need, challenges, prospects," *Vaccine*, 2014, 32 (14),1602–1609.
23. Fleming DT and Wasserheit JN "From epidemiological synergy to public health policy and practice: the contribution of other sexually transmitted diseases to sexual transmission of HIV infection," *Sexually Trans. Infect*, 1999; 75 (1), 3–17.
24. Kojima N, Klausner JD. An Update on the Global Epidemiology of Syphilis. *Curr Epidemiol Rep*. 2018, 5(1): 24–38.
25. Lamidi Y, Gbemisola OM. Seroprevalence of Syphilis among Pregnant Women Attending Antenatal Care in Kogi West, Nigeria. *J. Health Sci. Res*. 2020, 5(1), 24-29.
26. Olokoba AB, Olokoba LB, Salawu FK, Danburam A, Desala OO, Midalla JK. Syphilis and HIV coinfection in North-Eastern Nigeria. *Int. J. Trop. Med*. 2008; 3(3):70-72.
27. Ozumba UC, Oshi DC, Nwokeji CM, Anya SE. Trends in seroreactivity of syphilis among pregnant Nigerians. *Nig. J. Microbiol*. 22 (3), 168-179.

28. Aboyeji A and Nwabuisi C “Prevalence of sexually transmitted diseases among pregnant women in Ilorin, Nigeria, 2003, *J. Obst & Gynaec.*, 23(6), 637–639, 2003.
29. Shazia PS, Rama RMV, Janardhan Rao R. Declining seroprevalence of syphilis among pregnant women in a rural area. *J. Microbiol. Biotech. Res.* 2012; 2(2):305-307.
30. Mbamara SU, Obiechina NJ “Seroprevalence of venereal disease among pregnant women attending antenatal care (ANC) in Onitsha, Anambra State, Southeast, Nigeria,” *Nig. J. Med.*, 2011. 20 (1), 57–60.
31. Isa M, Bello H, Mustapha A, Mangga HK, Abbas MI. Prevalence of syphilis among pregnant women attending State Specialist Hospital Maiduguri, Borno, Nigeria, *Int. J. Res.* 2014; 1(11), 523–530.
32. Gao, L, Zhang, L, Jin, Q. Meta-analysis: Prevalence of HIV infection and syphilis among MSM in China". *Sexually Trans. Infect.* 2009, 85 (5): 354–58.
33. Taiwo SS, Adesiji YO, Adekanle DA “Screening for syphilis during pregnancy in Nigeria: a practice that must continue,” *Sexually Trans Infect.* 2007, 83 (5), 357-358.
34. Buseri F, Seiyaboh E, Jeremiah Z. Surveying infections among pregnant women in the Niger Delta, Nigeria. *J. Global Infect. Dis.* 2010, 2 (3), 203.
35. Creek TL, Thuku H, Kolou B, Ralman M, Kilmax PH. Declining syphilis prevalence among pregnant women in Northern Botswana: An encouraging sign for the HIV epidemic? *Sex. Trans. Dis.* 2005; 81:543-455.36.
36. Kirakaya-Samadoulougu F, Defer M, Yaro S, Fao P, Iibondo F, Lagani Y, Meda N, Robert A, Nagot N. Low seroprevalence of syphilis in Burkina Faso. *Sex Trans Infect.* 2010; 87:35-37.
37. Ratnam AV, Din SN, Hira SK, Bhat GJ, Wacha DS, Rukmini A, et al. Syphilis in pregnant women in Zambia. *Br J Vener Dis.* 1982;58(6):355-358.
38. Lindstrand A, Bergstrom S, Bugalho A, Zanconato G, Helgesson AM, Hederstedt B. Prevalence of syphilis infection in Mozambican women with second-trimester miscarriage and women attending antenatal care in second trimester. *Genitourin Med.* 1993;69(6):421-433.
39. Kwiek JJ, V. Mwapasa AP, Alker AS, Muula HE, Misiri ME, Molyneux, et al. Sociodemographic characteristics associated with HIV and syphilis seroreactivity among pregnant women in Blantyre, Malawi 2000-2004. *Malawi Med. Journal.* 2008;20(3):80-85.
40. Hussen S and Tadesse BT Prevalence of Syphilis among Pregnant Women in Sub-Saharan Africa: A Systematic Review and Meta-Analysis. *BioMed Res. Int.*, 2019, 4, 3-10.
41. Bukar M, Audu BM, Takai UI Ajayi BB, Kullima AA, Is routine antenatal screening for syphilis in Nigeria still justified clinically and economically? *Saudi Med. Journal*, 2009, 30(10), 1311–1315.
42. Lumbiganon P, Piaggio G, Villar J, et al. The epidemiology of syphilis in pregnancy. *Int J STD AIDS.* 2002;13(7):486-494.
43. Olowe, O. A. Makanjuola, O. B. Olowe, R. A. Olaitan, J. O. Ojurongbe, S. O. Prevalence of Syphilis among Pregnant Women in Two Health Care Facilities in South Western Nigeria *British J. Medicine & Med. Res.*, 2014, 4(34): 5431-5438.
44. Kebede E. and Chamiso, B Prevalence of Syphilis in Pregnancy in Addis Ababa, East African Med. J. 2000, 77 (4), 134-139.
45. Ghanem KG, Ram S, Rice PA. "The Modern Epidemic of Syphilis". *N. Engl. J. Med.* 2020, 382 (9): 845–54.
46. Ropper AH. "Neurosyphilis". *N. Engl. J. Med.* 2019, 381 (14): 1358–63.

47. Zinabie S, Belachew E, Yidenek T. et al., “Sero-prevalence of syphilis and HIV and associated factors in pregnant women attending antenatal care clinics in Debre Berhan Public Health Institutions, Ethiopia,” *American J. Biomed. and Life Sci.*, 2018, 6(3), 56–62.
48. Fissehatsion K, Ali I, Getachew A, Seroprevalence and risk factors of sexually transmitted infections (HIV, HBV and Syphilis) among pregnant women provided health care services, Addis Ababa, Ethiopia. *American J Health Res.* 2017, 5(5), 154–161.