

**TRADITIONAL USE, PHYTOCHEMISTRY AND PHARMACOLOGICAL  
ACTIVITIES OF FOUR DALBERGIA SPECIES (DALBERGIA SISSOO,  
DALBERGIA ODORIFERA, DALBERGIA MELANOXYLON AND  
DALBERGIA LACTEA VATKE) (A REVIEW)**

***ABSTRACT***

*This review study concerned to survey traditional use and scientific reports for four Dalbergia species, namely: Dalbergia sissoo, Dalbergia melanoxylon, Dalbergia odorifera and Dalbergia lactea vatke. Genus Dalbergia (Fabaceae or Leguminosae) is an important plant in traditional use. For example, in different part of the world these plants have been used traditionally to treatment a wide range of ailments such as blood diseases, syphilis, stomach problems, dysentery, nausea, eye and nose disorders, ulcers, skin diseases, abdominal pains, anthelmintic and etc.*

*Phytochemical studies on different parts of these plants have indicated the presences of various secondary metabolites except Dalbergia lactea vatke. The widely reported are flavonoids, neoflavonoids, isoflavonoids, terpenoids, and benzofuran among other things. On the other hand the bioassay studies on crude extract and pure isolates have shown significant, anti-inflammatory, anti-cancer, anti-oxidant, anti-microbial, anti-diabetic and anti-analgesic activities. However there is no literature survey performed on Dalbergia lactea vatke.*

**Key word:** *Dalbergia genus; phytochemical constituents; Biological activity*

---

---

## 1. INTRODUCTION

### 1.1 Background of the study

Plants are a valuable source of wide range of secondary metabolites that are used in pharmaceuticals, agrochemicals, flavour, fragrances, colourant, biopesticides and food additives due to that, two thirds of the new chemicals identified yearly were extracted from higher plants (Snafi, 2017). Plants are nature's gift remedies in treating limitless range of diseases from acute to chronic from human and live stocks, as a matter of fact currently medicinal plants are getting attention than ever more, especially in the line of multidrug resistant bacteria and chronic disease like cancer, so that the bioactive phytochemical constituents of plant are being explored worldwide for their broad-spectrum medicinal potencies (Rashed et al., 2018).

Medicinal plants used traditionally, are now moving from fringe to mainstream as people are becoming more aware of therapeutic properties of these medicinal plant resources and their products in maintaining health and preventing diseases. A medicinal plant" is any plant, which in one or more of its organ contains substance that can be used for the

#### 1.1.1 *Dalbergia sissoo*

therapeutic purpose or which, are precursors for the synthesis of use full drugs (Sundar and Habibur, 2018). From large number of medicinal plants genus of *Dalbergia* is the most common plants that are widely used throughout the world. Therefore on this seminar we are going to give detail explanation on these plants that belongs to this genus.

#### 1.1 The genus of *Dalbergia*

The genus *Dalbergia* is placed under the subfamily Faboideae containing 274 species distributed all over the world, especially in the tropical and subtropical regions (Moritsuka et al., 2017). These genus widely used in traditional medicinal system in Pakistan, India, Afghanistan, Bangladesh, Persia, Iraq, Palestine, India, Malaysia, Thailand, Indonesia, Cameroon, Sudan, Zimbabwe, Kenya, Tanzania and china (Saha et al., 1901). The wood of these species has a characteristic color and texture that makes it highly desirable, and they are referred to by the common name of rosewood (McClure et al., 2015). From the genus of *Dalbergia* the following four species are selected and reviewed.

*Dalbergia sissoo* is called Indian Rose wood and belongs to the legume family (Fabaceae). It is a large deciduous perennial tree found in the lowland region throughout India, Pakistan, Afghanistan, china and Nepal (Asif and Kumar, 2009). Traditionally an aqueous extract of the leaves of *Dalbergia sissoo* has been used for



**Fig. 1: Leaves and pods of *Dalbergia sissoo***

the treatment of gonorrhoea in arabic countries due to the presence of various biological activities such as effective against blood diseases, syphilis, stomach problems, dysentery, nausea, eye and nose disorders, ulcers, skin diseases and etc (Asif and Kumar, 2011).



**Fig. 2: Flowers of *Dalbergia sissoo***

### 1.1.2 *Dalbergia Odorifera*

*Dalbergia odorifera* is also known as fragrant rosewood, which belongs to genus of *Dalbergia*, is a semi-deciduous perennial tree. It grows in East Asian countries especially across Hainan and Guangdong province in China (Choi et al., 2009).

Heartwood is used in Chinese traditional medicine in the treatment of ischemia, blood stagnation syndrome swelling, rheumatic pain, epigastria, traumatic injuries and necrosis (Kang et al., 2005).



**Fig. 3 Leaves of *Dalbergia odorifera***



**Fig.4 Seeda of *Dalbergia odorifera***

### 1.1.3 *Dalbergia melanoxylon*

*Dalbergia melanoxylon* is known as African Blackwood, and it is a flowering plant belongs to family of Fabaceae known to be an economically important tree with high-quality wood and being one of the most expensive timbers in the world (Amri and



**Fig.5** leaves of *Dalbergia melanoxylon*

Juma, 2016). Traditionally, stem bark has a diverse local medicinal uses in Africa. For example, the leaves are boiled in soup and drunk to relieve pain in joints (Kareru et al., 2008) and dried leaves smoked as cigarette to treat asthma and bronchitis (Chigora et al., 2007).



**Fig.6** Areal parts of *Dalbergia melanoxylon*

### 1.1.4 *Dalbergia lactea vatke*

*Dalbergia lactea vatke* is one species belonging to the genus of Dalbergia, as reported by Fenetahun and Eshetu, 2017, the crushed leaves of this plant with water is used to treat Mastitis, Internal Parasite, local

Swelling. There is no phytochemical constituents and pharmacological investigation performed on this plant. The areal parts of this plant have been depicted below




**Fig.7** Leaves and flower of *Dalbergia lactea vatke*

## 2. Phytochemistry of *Dalbergia* species

### 2.1 . Flavonoids

Basic structure of neoflavonoid (1), flavonoid (2) and isoflavonoid are depicted below




**Fig.8: Basic structure of neoflavonoids (1), flavonoids (2) and isoflavonoids (3)**

The stem bark of *Dalbergia sissoo* contains Isoflavone-O-glycoside (Sabira et al., 2015). According to Liu, R et al., 2005, the heartwood of *Dalbergia odorifera* and *Dalbergia sissoo* contains neo flavonoids (4-6), flavonoids (7-10), and (15-19) (Zhao et al., 2011), and (23-25) (Zhao, X et al., 2019) from the dried heartwood of

*Dalbergia odorifera* and from the leave of *Dalbergia sissoo* compound (20) (Behera et al., 2013) are also obtained. From the heart wood of *Dalbergia melanoxylon* compound (11-14) is isolated (lin et al., 2019). From the heartwood of *Dalbergia odorifera* flavonoids (26 and 27) are isolated (Wang, H et al., 2014).

### 2.2 .Benzofurans

Basic skeletal structure of aryl benzofuran is depicted below




**Fig.9: The basic structure of benzofuran**

From the heartwood of *Dalbergia odorifera* two aryl benzofurans (27 and 28) were isolated (An et al., 2008). Obtusafuran (29) and melanoxin (30) (Muangnoicharoen and Frahm, 1981), and from the heart wood (31) (Donnelly et al., 1969) compounds are obtained from *Dalbergia melanoxylon*.

## 2.4. Terpenoids

Terpenoids have unsaturated molecules composed of linked isoprene units and its skeletal structure of isoprene unit is depicted below.




**Fig. 10: Skeletal structures of isoprene**

Phytochemical screening of the crude ethanol extract of the root of *Dalbergia sissoo* (Hajare et al., 2001) and methanolic

extracts of *Dalbergia odorifera* showed presence of terpenoids in stem and root (Tao and Wang, 2010). The bark of *Dalbergia melanoxylon* also contains terpenoids (Njeru and Obonyo, 2016). The heartwood of *Dalbergia odorifera* afforded the sesquiterpenes (35-38) (Wang, et al., 2014), and (39-44) (Zhao, X et al., 2019). No phytochemical investigation is performed on *Dalbergia lactea vatke*.

## 2.6 Cinnamylflavan



**Fig.11 Basic structure of Cinnamyl**

Cinnamylflavans (45 and 46) have been isolated in a number of phenol-cinnamyl alcohol condensations from the acetone extract of heartwood of *Dalbergia melanoxylon* (Donnfly et al., 1975). From the heartwood and root heartwood of *Dalbergia odorifera* cinnamylflavan namely Obtustyrene (47), Hydroxyobtustyrene (48), Isomucronustyrene (49) are isolated (Goda et al., 1985). The last compound (50

and 51) is isolated from the bark of *Dalbergia sissoo* (Reddy, R et al., 2008).

### 3. Pharmacological Activities

#### 3.1 Anticancer activity

Crude ethanol bark extract of *Dalbergia sissoo* also shows antiulcer activity (Al-Snafi, 2017). The methanol extract and of the heartwood of *Delbergia odorifera* possessed potent inhibition of human tumor cell. Compound, 2'-*O*-methyl-isoliquiritigenin, isolated from the heartwood of *Dalbergia odorifera*, showed cytotoxic activity against cancer cell (Saha et al., 1901). Flavonoids and phenolic components isolated from the heartwood of *Delbergia odorifera* also shows cytotoxic activity against cancer cell (Choi et al., 2009) and also compounds (33,24,25) have significant anti-tumor effects on human cancer cell (Zhao, X et al., 2019). AS reported by Bhattacharya, et al., (2014), compounds like an isoflavone, biochanin is a potent chemotherapeutic cancer preventive agent.

#### 3.2 Antioxidant Activities

Flavonoids compound isolated from the root of *Dalbergia odorifera* showed very strong antioxidant activities (Cheng et al., 1998). Stem bark extract of *Dalbergia sissoo* has higher antioxidant activity due to the presence of both the polyphenol and the flavonoid (Roy et al., 2011). Tannins and Neoflavanoids obtained

from the roots bark, stem bark and leaves of *Dalbergia melanxylon* are the potent antioxidant and free radical scavenger activities (Amri and Juma, 2016).

#### 3.3 Antimicrobial Activities

The investigation on *Dalbergia melanoxylon* extracts shows antimicrobial activity (Gundidza and Gaza, 1993). A crude extract and 3-Hydroxyisoflavanones from the stem bark of *Dalbergia melanoxylon* shows antibacterial activities (Mutai et al., 2013). Methanolic extracts of *Dalbergia sissoo* shows antibacterial activities against *S.aureus* and *Pseudomonas* (Yadav et al., 2008; Prasad et al., 2014). The isolated sesquiterpenes and flavonoids from the heart wood of *Dalbergia odorifera* also shows antibacterial activity against *C. albicans* and *S. aureus*, *R. solanacearum*. Isolated compound 15, 16, 17 and 18 showed strong antibacterial activity (Zhao et al., 2011).

#### 3.4 Anti-inflammatory activities

The Neoflavonoid Latifolin, compound (12) isolated from MeOH extract of *Dalbergia odorifera* exhibit anti-inflammatory activity (Lee et al., 2014). Flavonoids have been reported to exhibit anti-inflammatory properties (Hodeket al., 2002). As reported by Tao and Wang, (2010), 4, 2', 5'-Trihydroxy-

4'-methoxychalcone from *Dalbergia odorifera* exhibits anti-inflammatory properties. The ethanolic leaves extract of *Dalbergia sissoo* possesses anti-inflammatory activity (**Hajare et al., 2000**). Chalcone [(*E*)-3-(3,4-dihydroxyphenyl)-1-(2,3,4-trihydroxyphenyl) prop-2-en-1-one] or compound (**20**) isolated from the leaves of *Dalbergia sissoo* exhibit anti-inflammatory activity (**Behera et al., 2013**). The isolated compounds like isoflavanones, neoflavone benzofuran and N-cinnamoyl from the heartwood and bark of *Dalbergia melanoxylon* possesses anti-inflammatory activities (**Lin et al., 2019**).

### 3.5 Ant-diabetic activity

The ethanol, ethyl acetate, n-butanol and petroleum ether extracts of leave of *Dalbergia sissoo* showed most potent ant diabetic activities (**Panda et al., 2016**). According to Al-Snafi, (2017), the ethanol leaf extracts of this plant, exhibited high ant diabetic activity which is comparable with the standard drug,

Glibenclamide. Methanol extract (**Ninh, 2017**) and Isolated compounds like 6-dihydroxy-7-methoxyflavanone (**21**) and isoliquiritigenin (**22**) from the heart wood of *Dalbergia odorifera* show antidiabetic activity (**Zhao et al., 2013**). Anti-diabetic activities of *Dalbergia melanoxylon* are not reported. But since different flavonoids are isolated from the different parts of *Dalbergia melanoxylon* it will show anti-diabetic activities.

### 3.6 Analgesic activities

As reported by **Hajare et al., (2000)**, flavonoids are known to inhibit prostaglandin synthetase. Ethanolic extract of *Dalbergia sissoo* leaves (**Vasudeva et al., 2009**) and seeds (**Sehra et al., 2018**) have shown analgesic activity Analgesic activities of *Dalbergia melanoxylon* and *Dalbergia odorifera* are not reported.


## Summery in tabulated form


**Table 1: Compounds isolated from the genus dalbrgia**









No.	Class and compound name	Occurrence /part	References
1	Neoflavonoids (2'- <i>O</i> -methyl-isoliquiritigenin) Flavonoids Isoflavonoids	<i>Dalbergia odorifera</i> (HW) <i>Dalbergia melanoxylon</i> (HW) <i>Dalbergia sissoo</i> (HW)	(Zhao, X et al.,2011) (Zhao, X et al .,2019) (Liu, R et al., 2005)
2	Sisquiterpene	<i>Dalbergia melanoxylon</i> (bark) and <i>Dalbergia odorifera</i> (HW)	(Njeru, S. N. andObonyo, M. A. 2016) (Tao, Y and Wang, Y. 2010)
3	Obtusafuran and melanoxin	<i>Dalbergia odorifera</i> (HW) <i>Dalbergiamelanoxylon</i> (HW)	(An, R. et al., 2008). (Donnelly et al.,1969)
4	Phenolic compounds	<i>Dalbergia odorifera</i> (HW) <i>Dalbergia melanoxylon</i> (stem bark) <i>Dalbergia sissoo</i> (stem bark)	(Wang, H et al., 2014) (Swetha, U. 2017) (Roy, N et al., 2011).
6	Hydroxyobtustyrene and Isomucronustyrene	<i>Dalbergia odorifera</i> (HW) <i>Dalbergia melanoxylon</i> (HW) <i>Dalbergiasissoo</i> (bark)	(Goda et al., 1985) (Donnfly et al., 1975) (Reddy, R et al ., 2008).


**Note: HW: heartwood**


**Fig.12:Structure of isolated compounds**














## Pharmacological studies

**Table 2: A summary of biological activities**

Compound	Activity	References
2'- <i>O</i> -methyl-isoliquiritigenin	cytotoxic activity against cancer cell	(Saha, S et al .,1901)
Isoflavone and biochanin	cytotoxic activity against cancer cell	(Bhattacharya, M et al., 2014)
Flavonoids	Antioxidant activities	(Swetha, U. 2017)
A 3-Hydroxyisoflavanones	Antimicrobial activity	(Mutai, P et al., 2013)
Sesquiterpenes	Antibacterial acyivity	(Zhao, X et al., 2019)
A 4,2',5'-Trihydroxy-4'-methoxychalcone	Anti-inflammatory activities	(Tao, Y., and Wang, Y. 2010)
Chalcone or [( <i>E</i> )-3-(3,4-dihydroxyphenyl)-1-(2,3,4-trihydroxyphenyl) prop-2-en-1-one	Anti-inflammatory activities	(Bharath, M et al., 2013)
isoflavanones, neoflavone	Anti-inflammatory activities	(Lin .,et al 2019)

Compound	Activity	References
and benzofuran		
A 6-dihydroxy-7-methoxyflavanone and isoliquiritigenin	Antidiabetic activity	(Zhao, C et al., 2013).

### 3. Conclusion

Medicinal plants are the bio resources given by nature and are used to heal a group of human diseases and to evaluate the probable sources for new drugs. Among medicinal plants *Dalbergia odorifera*, *Dalbergia sissoo*, *Dalbergia melanoxylon* and *Dalbergia lactea vatke* are the known plants have been used medicinally for thousands of years all over the world. Except *Dalbergia lactea vatke*, phytochemical investigation and pharmacological activity of the bark, leave, heart wood, root and fruits of these plants reveal a number of secondary metabolites that showed good to moderate biological activities. Three of these plants has medicinal values since it contains more secondary metabolites such as terpenoids, flavonoids (including neo and iso flavonoids), cinnamoylflavan and benzofuran and these phytochemicals shows biological activities such as, anti-inflammatory, anti-cancer, anti-oxidant, anti-microbial, ant diabetic and ant analgesic activities.

### Reference

- Al-Snafi, A. E. (2017). Chemical constituents and pharmacological effects of *Dalbergiasissoo* (A review). *Journal of Pharmacy*, 716(12), 9775-9782.
- Amri, E., and Juma, S. (2016). Evaluation of antimicrobial activity and qualitative phytochemical screening of solvent extracts of *Dalbergia melanoxylon* (Guill. and Perr.). *International Journal of Current Microbiology and Applied Sciences*, 5(7), 412-423.
- An, R. B., Jeong, G. S., and Kim, Y. C. (2008). Flavonoids from the heartwood of *Dalbergia odorifera* and their protective effect on glutamate-induced oxidative injury in HT22 cells. *Chemical and Pharmaceutical Bulletin*, 56(12), 1722-1724.

- Asif, M., and Kumar, A. (2011). Phytochemical investigation and evaluation of antinociceptive activity of ethanolic extract of *Dalbergia sissoo* (Roxb.) bark. *Journal of natural science, biology, and medicine*, 2(1), 76.
- Asif, M., and Kumar, A. R. U. N. (2009). Anti-inflammatory activity of ethanolic extract of *Dalbergia sissoo* (roxb.) Bark. *Malaysian Journal of Pharmaceutical Sciences*, 7(1), 39-50.
- Behera, P. C., Verma, S. M., Kumar, P. M., Das, N. B., Mishra, P. M., and Baliarsingh, S. (2013).  
Anti Inflammatory and Anti Microbial Activity of Chalcone from *DalbergiaSissooRoxb* leaves. *American Journal of Phytomedicine and Clinical Therapeutics*, 1(2), 186-194.
- Bharath, M., Tulasi, E. L. R., Sudhakar, K., and Eswaraiyah, M. C. (2013). *Dalbergia sissoo* DC—an important medicinal plant. *International Journal of Research Pharmacognosy Chemistry*, 3(2), 384-388.
- Bhattacharya, M., Singh, A., and Ramrakhyani, C. (2014). *Dalbergia sissoo*-An Important Medical Plant. *Journal of Medicinal Plants*, 2(2), 76-82.
- Chaitra, S., Kumar, N. N., Shalini, P., Sindhu, R., and Raj, K. S. (2015). Phytochemical analysis and antibacterial activity of *Dalbergia paniculata roxb*. *International Journal of Pharmaceutical Sciences and Research*, 6(2), 712-716.
- Cheng, Z. J., Kuo, S. C., Chan, S. C., Ko, F. N., and Teng, C. M. (1998). Antioxidant properties of butein isolated from *Dalbergia odorifera*. *BiochimicaetBiophysicaActa (BBA)-Lipids and Lipid Metabolism*, 1392(2-3), 291-299.
- Choi, C. W., Choi, Y. H., Cha, M. R., Kim, Y. S., hwan Yon, G., Kim, Y. K., and Ryu, S. Y. (2009). Antitumor components isolated from the heartwood extract of *Dalbergia odorifera*. *Journal of the Korean Society for Applied Biological Chemistry*, 52(4), 375-379.
- Chigora, P., Masocha, R., Mutenheri, F. (2007). The role of indigenous medicinal knowledge (IMK) in the treatment of ailments in rural Zimbabwe: the case of Mutirikwi communal lands. *J. Sustainable Development of Africa*, 9, 26-43.

- Donnelly, B. J., Donnelly, D. M. X., O'Sullivan, A. M., and Prendergast, J. P. (1969). The isolation and structure of melanoxin a new dihydrobenzofuran from *Dalbergiamelanoxylon* Gilg. (Leguminosae). *Tetrahedron*, 25(18), 4409-4414.
- Donnelly, D. M., O'Reilly, J., and Whalley, W. B. (1975). Neoflavanoids of *Dalbergia melanoxylon*. *Phytochemistry*, 14(10), 2287-2290.
- Fenetahun, Y., and Eshetu, G. (2017). A review on ethnobotanical studies of medicinal plants use by agro-pastoral communities in, Ethiopia. *Journal of Medicinal Plants Studies*, 5(1), 33-44.
- Gundidza, M., and Gaza, N. (1993). Antimicrobial activity of *Dalbergiamelanoxylon* extracts. *Journal of ethnopharmacology*, 40(2), 127-130.
- Hajare, S. W., Chandra, S., Sharma, J., Tandan, S. K., Lal, J., and Telang, A. G. (2001). Anti-inflammatory activity of *Dalbergiasissoo* leaves. *Fitoterapia*, 72(2), 131-139.
- Hajare, S. W., Chandra, S., Tandan, S. K., Sarma, J., Lal, J., and Telang, A. G. (2000). Analgesic and antipyretic activities of *Dalbergiasissoo* leaves. *Indian Journal of Pharmacology*, 32(6), 357-360.
- Hodek, P., Trefil, P., and Stiborová, M. (2002). Flavonoids-potent and versatile biologically active compounds interacting with cytochromes P450. *Chemico-biological interactions*, 139(1), 1-21.
- Kang TH, Tian YH, Kim YC. (2005). Isoliquiritigenin: A competitive tyrosinase inhibitor from the heartwood of *Dalbergia odorifera*. *Biomolecules and Therapeutics* 13, 32-34.
- Kareru, P.G., Gachanja, A.N., Keriko, J.M., and Kenji, G.M. 2008. Antimicrobial activity of some medicinal plants used by herbalists in eastern province, Kenya. *African J. Traditional, Complementary and alternative med.*, 5(1): 51-55.
- Khalivulla, S. I., Reddy, B. A., Gunasekar, D., Murthy, M. M., Rao, T. P., Blond, A., and Bodo, B. (2007). A new C-geranylatedisoflavone from *Dalbergiapaniculata*. *Natural Product Communications*, 2(11), 1109 - 1111.

- Lee, D. S., and Jeong, G. S. (2014). Arylbenzofuran isolated from *Dalbergiaodorifera* suppresses lipopolysaccharide-induced mouse BV2 microglial cell activation, which protects mouse hippocampal HT22 cells death from neuro inflammation-mediated toxicity. *European journal of pharmacology*, 728, 1-8.
- Lee, D. S., Kim, K. S., Ko, W., Li, B., Keo, S., Jeong, G. S., & Kim, Y. C. (2014). The Neoflavonoid Latifolin Isolated from MeOH Extract of *Dalbergiaodorifera* Attenuates Inflammatory Responses by Inhibiting NF- $\kappa$ B Activation via Nrf2-Mediated Heme Oxygenase-1 Expression. *Phytotherapy research*, 28(8), 1216-1223.
- Lee, D. S., Li, B., Im, N. K., Kim, Y. C., and Jeong, G. S. (2013). 4, 2', 5'-Trihydroxy-4'-methoxychalcone from *Dalbergiaodorifera* exhibits anti-inflammatory properties by inducing heme oxygenase 1 in murine macrophages. *International immunopharmacology*, 16(1), 114-121.
- Lin, Shuai, Rong-Hua Liu, Guang-Qiang Ma, Dan-Yi Mei, Feng Shao, and Lan-Ying Chen.(2019). Two new compounds from the heartwood of *Dalbergiamelanoxylon*." *Natural product research* 1-8.
- Liu, R. X., Wang, Q., Guo, H. Z., Li, L., Bi, K. S., and Guo, D. A. (2005). Simultaneous determination of 10 major flavonoids in *Dalbergiaodorifera* by high performance liquid chromatography. *Journal of pharmaceutical and biomedical analysis*, 39(3-4), 469-476.
- Liu, R., Ye, M., Guo, H., Bi, K., and Guo, D. A. (2005). Liquid chromatography/electrospray ionization mass spectrometry for the characterization of twenty-three flavonoids in the extract of *Dalbergiaodorifera*. *An International Journal Devoted to the Rapid Dissemination of Up-to-the-Minute Research in Mass Spectrometry*, 19(11), 1557-1565.
- Ma, F. Y., Gu, C. B., Li, C. Y., Luo, M., Wang, W., Zu, Y. G., and Fu, Y. J. (2013). Microwave-assisted aqueous two-phase extraction of isoflavonoids from *Dalbergiaodorifera* T. Chen leaves. *Separation and Purification Technology*, 115, 136-144.

- McClure, P. J., Chavarria, G. D., & Espinoza, E. (2015). Metabolic chemotypes of CITES protected *Dalbergia* timbers from Africa, Madagascar, and Asia. *Rapid Communications in Mass Spectrometry*, 29(9), 783-788.
- Muniyappan, N., and Nagarajan, N. S. (2014). Green synthesis of silver nanoparticles with *Dalbergiaspinosa* leaves and their applications in biological and catalytic activities. *Process Biochemistry*, 49(6), 1054-1061.
- Mutai, P., Heydenreich, M., Thoithi, G., Mugumbate, G., Chibale, K., and Yenesew, A. (2013). 3-Hydroxyisoflavanones from the stem bark of *Dalbergiamelanoxylon*: Isolation, antimycobacterial evaluation and molecular docking studies. *Phytochemistry letters*, 6(4), 671-675.
- Moritsuka, E., Chhang, P., Tagane, S., Toyama, H., Sokh, H., Yahara, T., & Tachida, H. (2017). Genetic variation and population structure of a threatened timber tree *Dalbergiacochinchinensis* in Cambodia. *Tree genetics and genomes*, 13(6), 115.
- Muangnoicharoen, N., and Frahm, A. W. (1981). Arylbenzofurans from *Dalbergiaparviflora*. *Phytochemistry*, 20(2), 291-293.
- Ninh S. (2017). A Review on the medicinal plant *Dalbergiaodorifera* species: Phytochemistry and biological activity. *Evidence-Based Complementary and Alternative Medicine*, p27.
- Njeru, S. N., and Obonyo, M. A. (2016). Potency of extracts of selected plant species from Mbeere, Embu County-Kenya against *Mycobacterium tuberculosis*. *Journal of Medicinal Plants Research*, 10(12), 149-157.
- Panda, S. K., Padhy, R. P., Pani, S., and Bal, K. (2016). Phytochemical investigation and antidiabetic activity of Leaf extracts of *Dalbergiasissoo* (Roxb.) in alloxan induced diabetic rats. *The American Journal of science and medical research*, 1(2), 186-189.
- Prasad, N., Nandi, D., Arora, S., & Pandey, A. (2014). In vitro evaluation of antibacterial properties of *Moringaoleifera*, *Dalbergiasissoo* and *Alstoniascholaris* in Vitro. *Journal of Biology, Agriculture and Healthcare* 4(15), 2224-3208.

- Rashed, M. M., Ghaleb, A. D., Li, J., Nagi, A., Hua-wei, Y., Wen-you, Z., and Tong, Q. (2018). Enhancement of Mass Transfer Intensification for Essential Oil Release from *Lavandula pubescence* Using Integrated Ultrasonic-Microwave Technique and Enzymatic Pretreatment. *ACS Sustainable Chemistry and Engineering*, 6(2), 1639-1649.
- Reddy, R. V. N., Reddy, N. P., Khalivulla, S. I., Reddy, M. V. B., Gunasekar, D., Blond, A., and Bodo, B. (2008). O-Prenylated flavonoids from *Dalbergiasissoo*. *Phytochemistry Letters*, 1(1), 23-26.
- Roy, N., Laskar, R. A., Sk, I., Kumari, D., Ghosh, T., and Begum, N. A. (2011). A detailed study on the antioxidant activity of the stem bark of *Dalbergiasissoo*Roxb., an Indian medicinal plant. *Food chemistry*, 126(3), 1115-1121.
- Sabira Sultana, Hafiz Mohammad Asif ,NaveedAkhtar and NaheedAkhtar. (2015) *Dalbergia sissooroxb: Monograph. International journal of Pharmacognosy*, 2(9): 440-443.
- Saha, S., Shilpi, J. A., Mondal, H., Hossain, F.,Anisuzzman, M., Hasan, M. M., and Cordell, G. A. (2013). Ethnomedicinal, phytochemical, and pharmacological profile of the genus *Dalbergia* L.(Fabaceae). *Phytopharmacology* , 4(2), 291-346.
- Sehra, S. Y., and Sharma, J. (2018). Pharmacological effects and medicinal importance of *dalbergiasissoo*-a review. *International Journal of Pharmaceutical, Chemical and Biological Sciences*, 8(2),234-243.
- Sundar, R. A., and Habibur, R. C. (2018). Pharmacognostic, phytochemical and antioxidant studies of *Gardenia latifolia*Aiton: An ethnomedicinal tree plant. *International Journal of Pharmacognosy and Phytochemical Research*, 10(5), 216-228.
- Swetha, U. (2017). Antioxidant Activity of *DalbergiaMelanoxyton* Bark Extract. *International Journal of Applied Pharmaceutical Sciences and Research*, 2(4), 114-120.
- Tao, Y., and Wang, Y. (2010). Bioactive sesquiterpenes isolated from the essential oil of *Dalbergia odorifera* T. Chen. *Fitoterapia*, 81(5), 393-396.
- Upwar, N., Patel, R., Waseem, N., &Mahobia, N. K. (2011). Evaluation of anthelmintic activity of *dalbergia sissooroxb*. *International Journal of Pharmaceutical Sciences and Research*, 2(1), 171-174.

- Vasudeva, N., Vats, M., Sharma, S., and Sardana, S. (2009). Chemistry and biological activities of the genus *Dalbergia*-A review. *Pharmacognosy Reviews*, 3(6), 307.
- Wang, H., Dong, W. H., Zuo, W. J., Liu, S., Zhong, H. M., Mei, W. L., and Dai, H. F. (2014). Five new sesquiterpenoids from *Dalbergiaodorifera*. *Fitoterapia*, 9(5), 16-21.
- Goda, M., Katayama, K., Ichikawa, M., Shibuya, F., Kiuchi, and U. Sankawa. (1985). "Inhibitors of prostaglandin biosynthesis from *dalbergiaodorifera*. *Chemical and Pharmaceutical Bulletin*, 33(12), 5606–5609.
- Yadav, H., Yadav, M., Jain, S., Bhardwaj, A., Singh, V., Parkash, O., and Marotta, F. (2008). Antimicrobial property of a herbal preparation containing *Dalbergia sissoo* and *Daturastramonium* with cow urine against pathogenic bacteria. *International journal of immunopathology and pharmacology*, 21(4), 1013-1020.
- Zhao, C., Liu, Y., Cong, D., Zhang, H., Yu, J., Jiang, Y., and Sun, J. (2013). Screening and determination for potential  $\alpha$ -glucosidase inhibitory constituents from *Dalbergia odorifera* T. Chen using ultrafiltration LC/ESI MSn. *Biomedical Chromatography*, 27(12), 1621-1629.
- Zhao, X., Mei, W., Gong, M., Zuo, W., Bai, H., and Dai, H. (2011). Antibacterial activity of the flavonoids from *Dalbergia odorifera* on *Ralstonia solanacearum*. *Molecules*, 16(12), 9775-9782.
- Zhao, X., Wang, C., Meng, H., Yu, Z., Yang, M., and Wei, J. (2019). *Dalbergiaodorifera*: A review of its traditional uses, phytochemistry, pharmacology, and quality control. *Journal of ethnopharmacology*, 348(11), 23-28.

#### COMPETING INTERESTS DISCLAIMER:

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.