

Original Research Article

EVALUATION OF HEALING AND MICROBIAL RESPONSES TOSURGICAL WOUNDS IN ALLOXAN-INDUCEDDIABETIC RABBITS

Abstract

The wounds of diabetic patients are associated with delayed healing and microbial invasion. This study evaluated the healing and microbial responses to surgical wounds in diabetic rabbits. Sixteen (16) New Zealand White rabbits were used in this study. They were divided into 4 groups of 4 rabbits each, namely; A, B, C and D. A is the control (non-diabetic and non-wounded), B (diabetic and non-wounded), C (wounded and non-diabetic) and D (wounded and diabetic). Three (3) cm² skin wounds were created in a standard aseptic condition while diabetes was induced by intravenous administration of 200 mg/kg of alloxan monohydrate. Blood samples were collected from ear vein of each rabbit on days 0, 3, 7, 14, 21 and 28 post-surgery for analysis of blood glucose level. Wound samples were collected on days 3, 7, 14 and 21 for evaluation of microbial contaminants. The wounds were examined for exudation and recorded as none (0), scant (1), moderate (2) and large (3) and diameter of wound epithelialization and contraction were determined. The blood glucose concentration of groups B and D from days 3 to 28 ranged between 288.00 ± 40.22 mg/dl to 358.65 ± 25.89 mg/dl, and were significantly higher (P < 0.05) than the corresponding values of groups A and C which ranged from 120.60 mg/dl ± 14.29 to 129.60 ± 8.18 mg/dl. The rate of wound contraction in the C group on day 7 (11.02 ± 2.17%) was significantly higher (P < 0.05) than the D group (2.23 ± 4.60%). *Staphylococcus* and *Pseudomonas* subspecies were identified in the wounds of rabbits in groups C and D. It was concluded that alloxan-induced diabetes caused delayed wound healing while *Staphylococcus* and *Pseudomonas* species were the predominant microorganism detected in the acute wounds of diabetic and non-diabetic rabbits.

Key words: Evaluation, wound healing, alloxan-induced diabetes, microbial infection, NZW rabbits.

Introduction

Wound healing is a fundamental, dynamic, coordinated, interdependent and overlapping cellular and immunologic responses of tissues to injury [1, 2, 3]. It is essential to prevent pathogenic invasion of damaged tissues and to partially or completely reform the affected tissues [2, 3]. Experimental wounds prepared under aseptic conditions are expected to heal at a specific range of times. However, when this co-ordinated biologic process that restores tissue continuity after injury are interrupted due to some disease conditions or infections, the wound healing may delay. Such interruption may occur at any of the phases of wound healing such as angiogenesis, epithelialization, wound contraction and remodelling [4, 5].

Diabetes is a metabolic and heterogeneous disorder characterised primarily by hyperglycaemia due to deficiency of insulin or insensitivity of insulin receptors for normal processes of glucose metabolism in the body [6, 7]. The hyperglycaemia is manifested as glycosuria, hyperlipidaemia,

Comment [p1]: Wordy sentences:

to evaluate

Comment [p2]: Hard-to-read text

.At the same time,

Comment [p3]: Text inconsistencies

coordinated: co-ordinated

Comment [p4]: Comma misuse within
Clauses

, and

Comment [p5]: Incorrect noun number

responses → response

Comment [p6]: Confused words

biologic → biological

Comment [p7]: Faulty subject-verb
Agreement

are → is

Comment [p8]: Punctuation in
compound/complex
sentences

, such

Comment [p9]: Punctuation in
compound/complex
sentences

epithelialization,

Comment [p10]: Mixed dialects of English

remodelling → remodeling

Comment [p11]: Mixed dialects of English

characterised → characterized

Comment [p12]: Mixed dialects of English

hyperglycaemia → hyperglycemia

Comment [p13]: Unclear sentences

Comment [p14]: Mixed dialects of English
hyperglycaemia → hyperglycemia

Comment [p15]: Mixed dialects of English
hyperlipidaemia → hyperlipidemia

polyuria and polydipsia [7, 8, 9]. The disease affects man and animals worldwide [10, 11, 12]. Nigeria has the greatest number of people living with diabetes in Africa [13, 14] reported a prevalence of 0.34% of diabetes mellitus among dogs attending first opinion practice in the UK. Also [15] reported a prevalence of 0.22 % of diabetic mellitus cases among dogs presented for Veterinary Care in Warri, Delta State, Nigeria.

It has been established that impaired wound healing is a notable complication of diabetes mellitus [16, 17]. Such impaired wound healing develops due to various factors associated with diabetes including slow metabolic rate, macro and microvascular complications at the edges of the wounds and peripheral neuropathy [17, 18, 19]. Diabetic wound healing in rabbits is characterised by reduced length and density of blood vessels with a greater radial diffusion distance between vessels, a less efficient network for nutrient exchange and an increase in the number of inflammatory cells and fibroblasts at few weeks post-wounding [20].

Infections have been identified as one of the major contributing factors to delayed wound healing in diabetic and normoglycemic wounds. These wound infections may include bacterial, viral and fungal infection [21]. Diabetic wounds develop more sustained and co-invasive infection that delay wound healing [22]. Consequently, wounds of diabetic patients degenerate to chronic wounds, which may defy therapeutic management [6].

Diabetes affects virtually all the stages of wound healing. The impaired healing observed in diabetic wounds correlates with decreased keratinocyte, fibroblast and immune cell migration into the wound, reduced endothelial cell angiogenesis and decreased efferocytosis [18, 23, 24]. Worthy of note is that there are numerous recent advances in therapeutic modalities targeting enhancing wound healing throughout the world. However, a thorough knowledge of association between microbial invasions of diabetic wounds and the rate of diabetic wound healing is of utmost importance in the study of wound physiology, and this poses a clinical challenge [22]. [3] Proposed an ideal intervention for wound care that must involve components that act at different stages in the process of natural wound healing such as blood-clotting, anti-microbial, growth-promoting and analgesic properties. Animal models of diabetes are useful in biomedical studies because they offer the promise of new insights into human diabetes [25]. This study evaluated the healing and microbial responses to surgical wounds in alloxan-induced diabetic rabbits.

Materials and Methods

Experimental animals, feeding and conditioning

Ethical clearance was obtained for this study from Ahmadu Bello University Committee on Animal Use and Care (ABUCAUC), (Reference number: ABUCAUC/2019/028). This study was carried out in the Department of Veterinary Surgery and Radiology, Faculty of Veterinary Medicine, Ahmadu Bello University, Zaria, Kaduna State, Nigeria. Sixteen adult New Zealand White (NZW) rabbits of both sexes aged between 11.6 - 12.4 months and weighing between 1.8 - 3.2 kg were used for this study. They were procured from the National Animal Production and Research Institute (NAPRI), Zaria, Kaduna State, Nigeria.

The rabbits were kept in intensive management system in separate cages and fed morning and evening daily with combined poultry grower feeds, finisher feeds (Grand Cereals Limited, Jos, Plateau State) and miller's bran that were mixed in equal proportion in order to regulate the

Comment [p16]: Punctuation in compound/complex sentences

. Nigeria

Comment [p17]: Word choice
greatest → most signi*cant

Comment [p18]: Punctuation in compound/complex sentences

Also,

Comment [p19]:

Comment [p20]: Mixed dialects of English

characterised → characterized

Comment [p21]: Incorrect noun number
infection → infections

Comment [p22]: Determiner use
(a/an/the/this, etc.)

a more

Comment [p23]: Incorrect noun number
infection → infections

Comment [p24]: Determiner use
(a/an/the/this, etc.)

the association

Comment [p25]: Wordy sentences

and this → which

Comment [p26]: Wordy sentences

offer the promise of →
promise

Comment [p27]: Incorrect noun number

adult → adults

Comment [p28]: Punctuation in compound/complex sentences

, finisher →
; finisher, and finisher. Finisher

Comment [p29]: Faulty subject-verb Agreement

were → was

blood glucose level of the hyperglycaemic groups. *Daucuscarota subsp. sativus* leaves was given to the rabbits twice weekly to avoid gastro-intestinal impaction and constipation while water was given *ad-libitum* throughout the experimental period.

The rabbits were conditioned to the laboratory for two weeks before the commencement of the experiment. During this period, they were administered with anthelmintic Ivermectin (Hebei, Kexing pharmaceutical, China) prophylaxis at dosage of 0.4mg/kg start, and repeated after two weeks to prevent mites and other internal and external parasites. The rabbits were clinically evaluated using body condition scores, vital and haematological parameters and adjudges apparently healthy before they were certified fit for the study.

Experimental Protocols

The rabbits were divided into 4 groups (A, B, C and D) of 4 rabbits each comprising of two males and two females in each group. Group A (control): No Diabetes was induced and No Wound was created (NDNW); Group B: Diabetes was induced in this group, but No Wound was created (DNW); Group C: Wound was created, but Diabetes was Not induced (WND) and Group D: Wound was created and Diabetes was induced (WD), respectively.

Induction of diabetes

The rabbits were first anaesthetized by intramuscular administration of 7 mg/kg xylazine hydrochloride (Bioveta a. s., Czech Republic) and 50 mg/kg ketamine hydrochloride (Laborate pharmaceutical, India). The pinna were then shaved and dabbed with xylene to dilate the marginal ear vein and 200 mg/kg of Alloxan Monohydrate (SIGMA-aldrich, UK) was administered intravenously through the marginal ear vein on two occasions with 100 mg/kg each at an interval of 72 hours [26, 27]. Blood glucose level was estimated in the blood samples collected via the ear vein using a glucometer (ACCU-CHEK^(R), Roche, Mannheim, Germany) daily until hyperglycaemia was established and stabilized.

Wound creation

The dorsum of four rabbits each in the NDW (group C) and WD (group D) were prepared aseptically. The rabbits were anaesthetized by intramuscular injection of xylazine hydrochloride ((Bioveta a. s., Czech Republic) at 7 mg/kg and Ketamine Hydrochloride (Laborate pharmaceutical, India) at 50 mg/kg [28, 29]. A full-thickness skin wounds of 3 cm² (Fig.1) was created at the dorsum of each of the rabbits using a template designed from x-ray film [30]. The wounds were bandaged with sterile gauze and re-dressed only on the days of sample collections (days 0, 3, 7, 14, 21 and 28).

Blood and Microbial Sample Collections

A drop of blood was collected from the ear vein of each of the rabbits on day 0 (before wound creation) and on days 3, 7, 14, 21 and 28 post-surgery for blood glucose evaluation. Sterile swabs were used to collect microbial samples from the wound surfaces in groups C and D on days 3, 7, 14 and 21 post-surgery for qualitative evaluation of microorganisms present in the wound. Subsequently, after the experiment, the diabetic rabbits were nursed back to their normal health using insulin to reverse diabetes.

Comment [p30]: Faulty subject-verb Agreement

was → were

Comment [p31]: Wrong or missing prepositions

to → in

Comment [p32]: Incorrect verb forms

start → started

Comment [p33]: Mixed dialects of English

haematological → hematologica

Comment [p34]: Unclear sentences

Comment [p35]: Mixed dialects of English

anaesthetized → anesthetized

Comment [p36]: Misspelled words

Laborate → Labourite

Comment [p37]: Faulty subject-verb Agreement

were → was

Comment [p38]: Misspelled words

aldrich → Aldrich

Comment [p39]: Mixed dialects of English

hyperglycaemia → hyperglycemia

Comment [p40]: Mixed dialects of English

anaesthetized → anesthetized

Comment [p41]: Misspelled words

Laborate → Labourite

Comment [p42]: Determiner use (a/an/the/this, etc.)

wounds → wound

Wound Assessment

On each day of post-surgery sample collections (days 0, 3, 7, 14, 21 and 28), the blood was placed in ACCU-check glucometer (ACCU-CHEK^(R), Roche, Mannheim, Germany) and the blood glucose estimates recorded. The wounds were clinically assessed for exudation, epithelialization and contraction. The exudation was recorded as none (0), scant (1), moderate (2) and large (3) [30]. The diameter of the wounds and the epithelial region were measured using tracing papers (Fig. 2a) and a meter rule (Fig. 2b), and calculated with graph sheets (Fig. 3) to determine the epithelialization and contraction [31, 32]. The percentage wound contraction and epithelialization were determined using the formulae described by [33].

Microbial Evaluation

The various isolation media and reagents used for this study were purchased from Oxoid® Ltd and were prepared as described by the manufacturers [34]. The swab samples were inoculated in already prepared blood agar and incubated at 37°C for 48 hours and clusters of microbial growth were observed. This microbial growth was further sub-cultured; A newly prepared blood agar was divided into 4 quadrants. A portion of the medium that has the microbial growth was collected using a wire loop sterilized to red hot and cooled for 10 seconds. The loop was streaked at one quadrant several times. The blood agar was turned at 90°C and from the first quadrant, the second quadrant was streaked several times. The same process was repeated for the 3rd and 4th quadrants for isolation of discrete colony. The subculture process was also carried out in MacConkey agar. The discrete colony was transferred to nutrient agar slant and incubated at 37°C for 48 hours for their separation into morphological characteristics. These pure isolates were subjected to gram stains and various biochemical reactions such as: *Catalase test, Oxidase test, Indole test, Voges-Proskauer (VP) Test, Methyl Red, Methyl Red, Citrate Utilization, Motility test, Urease test, TSI test, Urease test, Mannitol test, Coagulase test, Maltose test, Arginine test* [35, 36, 37].

Data Analysis

The data obtained from blood glucose estimation and wound assessment, were expressed as mean ± Standard Error of Mean (SEM), and subjected to statistical packages with Graph-Pad Prism version 5.03 and subsequently analysed using One Way Analysis of Variance (ANOVA) and Friedman test. Values of $P = .05$ were considered significant. The results were presented in tables and figures.

Results

The mean blood glucose level of the rabbits in group B was lowest on pre-surgery day (0), and increased sharply on post-surgery day 3 (329.40 ± 61.56), and remained at high value until day 28 (273.60 ± 37.26). The mean blood glucose level of rabbits in group D was lowest on Pre-surgery day 0 (113.80 ± 12.50), increased sharply and reached the peak on day 3 (344.70 ± 50.29) and remained constantly high until 28 (295.20 ± 42.57). There were no significant difference ($P > 0.05$) in the mean blood glucose level of the experimental rabbits of the different groups before the induction of diabetes (day 0). Following induction of diabetes, the mean blood glucose level

Comment [p43]: Improper formatting

3rd → third

Comment [p44]: Improper formatting

4th → fourth

Comment [p45]: Incorrect noun number

colony → colonies

Comment [p46]: Misspelled words

Proaskauer → Proskauer

Comment [p47]: Mixed dialects of English

analysed → analyzed

Comment [p48]: Faulty subject-verb Agreement

were → was

of groups B and D rabbits rose significantly ($P < 0.05$) and were higher than the levels in groups A and C rabbits from day 3 to 28 (Fig. 4). There was no significant difference ($P > 0.05$) between the mean blood glucose values of groups B and D; and also between the values in groups A and C rabbits.

The mean exudates value in group C was highest on day 3 (1.75 ± 0.25) while that of the value of group D was highest on day 3 (1.75 ± 0.25) and 7 (1.75 ± 0.25) before the values in both groups declined progressively to approximately 0.00 ± 0.00 on day 21 (Fig. 5). There was no significant difference ($P > 0.05$) in the exudate value observed between the two wounded (WND and WD) groups.

The degree of epithelialization(%) in group C rabbits was observed on day 14(29.37 ± 8.17) and 21(73.50 ± 1.37), while that of group D rabbits was on day 14 (31.78 ± 5.20) and 21(74.66 ± 7.01) (Figs 6 and 8). The wounds in both groups have healed completely by day 28 (Fig. 9). There was no significant difference ($P > 0.05$) between the values of percentage wound epithelialization of rabbits in group C and D throughout the period of study. However, the percentage wound epithelialization observed in group (D) rabbits on day 14 and 21 were slightly higher than the corresponding values observed in group C rabbits (Fig. 6).

The percentage wound contraction in rabbits in group C was lowest on day 3 (1.15 ± 3.36), but increased progressively to complete contraction on day 28 (100.00 ± 0.00). Also, the percentage wound contraction in group D rabbits was lowest on day 3 (1.02 ± 2.50) but increased progressively to complete contraction on day 28 (100.00 ± 0.00). The value of the percentage wound contraction observed on day 7 in group C (11.02 ± 2.17) was significantly higher ($P < 0.05$) than that observed in the D group (2.23 ± 4.60) (Fig. 7). There was complete wound contraction (100) in both group C and D rabbits on post-surgery day 28 (Fig. 9)

Four bacterial isolates namely; *Staphylococcus* species, *Pseudomonas* species, *Bacillus* species and *Corynebacterium* were identified from the wounds of the two wounded groups (C and D). However, biochemical test results confirmed the presence of only two bacterial species; *Staphylococcus* spp. and *Pseudomonas* spp (Table 1). Further biochemical tests result based on conventional biochemical identification of *Staphylococcus* isolate [37] revealed the presence of two *Staphylococcus* subspecies namely; *Staphylococcus aureus* and *Staphylococcus epidermidis*(Table 1).

Discussions

Wound healing is a cellular and immunologic responses of tissues to injury essential for repair of the injured tissues [1, 2,3]. The wounds of diabetic patients are associated with delayed healing and microbial invasion of the injured tissues

The observed rise in mean blood glucose level of rabbits in DNW and WD groups, above the normal mean blood glucose level of New Zealand White rabbits (150g/dl) was an indication of hyperglycaemia, a major sign of diabetes mellitus [27]. It was possible that alloxan destroyed the pancreatic β -cells in these rabbits and caused a deficiency of insulin secretion which resulted in the metabolic derangements associated with Type-1 Diabetes Mellitus (T1DM). It could have also generated reactive oxygen species in a cyclic redox reaction with its hydroxyl radicals responsible for the selective necrosis and death of the pancreatic islet β -cells [38]. There was no

Comment [p49]: Hard-to-read text
while →. In comparison,

Comment [p50]:

Comment [p51]: Faulty tense sequence
have → had

Comment [p52]: Incorrect noun number
group → groups

Comment [p53]: Incorrect noun number
day → days

Comment [p54]: Incorrect noun number
responses → response

Comment [p55]: Word choice
normal → average

Comment [p56]: Mixed dialects of English
hyperglycaemia → hyperglycemia

significant difference ($P > 0.05$) between the blood glucose level of the rabbits in groups A and C and between groups B and D.

Scanty to moderate exudates **was** observed in the wounded rabbits in groups C and D. The wound exudates may contain fluid, proteins, serum, fibrin, and white blood cells to **escapes** into a wound defect and combine with necrotic cells and other waste materials in the wound to form wound exudates [39, 40]. It was probable that such components of exudates that facilitate healing such as fibrin, proteins, serum and water and at a therapeutic proportion dominated the exudates observed in both **group** C and D and contributed to the healing of the acute wound. The exudate in WD group remained at the peak up to **psd 7** as **agaist** **psd3** in group C probable because there was a delay in the metabolic processes that facilitate the resorption of the preparatory phases of wound healing in this WD group. Such delay could cause retention of these exudates at the peak value within 7 days duration in this group D when compared with group C. Scant to moderate wound fluids was beneficial to wound healing because **it** provided a moist environment for cell migration and supply of various nutrients needed for the healing [41, 42]. It has been reported that moist wounds heal two to three times faster than dry wounds [40]. The completed wound epithelialization and contraction within 28 days indicated that these defects healed by myofibroblast contraction and **was** in line with expected rate of experimental wound contraction in rabbits [43]. The loose skin at the **rabbit** back and linear edges of the square wounds favoured effective contraction, the linear traction forces of the myofibroblasts.

The mean percentage epithelialization in rabbits in group D which was slightly higher than the value recorded in group C on post-surgery day 14 indicated delayed wound healing in group D. Epithelial tissue covering was observed higher in group D probably because the contraction was delayed, due to delayed centripetal movement of the dermis and epidermis that border the wound defect to reduce the wound size [24]. In group C, the linear traction force of the myofibroblast was at a higher rate which facilitated the loose skin to cover the epithelial tissues within an interval of assessment [43]. On the other hand, the slow body metabolism associated with diabetes delayed these wound healing processes in group D rabbits thereby exposing the epithelial tissues for a longer duration of time for assessment [18, 19].

A shorter duration of wound contraction observed in group C indicated faster granulation and epithelialization which prepared the wound bed for faster contraction in this group when compared with group D. It might also be that there was a delay in the earlier preparatory phase of wound healing such as regression and haemostasis which consequently delayed contraction from **psd 3** to 14 in group D. Hyperglycaemia following diabetes caused excessive thickening of the basement membrane of blood vessels at the edges of skin wounds which resulted in poor vascular network and disorganized stages of wound healing, (18, 19). Hyperglycaemia has also been implicated in the decreased population of endothelial progenitor cells, progression of macro and microvascular **complication**, dysfunctional and loss of extracellular integrity and decreased efferocytosis all of which delay wound angiogenesis (19, 24).

The minimal organism isolated in this work might be due to comprehensive aseptic measures employed both in the surgery theatre, animal house and throughout the period of wound assessment. These two organisms (*Staphylococcus* and *Pseudomonas*) have always been isolated from acute, traumatic and diabetic wounds probably because they are endemic and ubiquitous in our environment, opportunistic in nature and have high affinity in wounds [37]. It has been reported that *Staphylococcus aurous* is the most frequently isolated **organisms** from diabetic foot infection [44, 45]. Samples from the wound of diabetic patients collected from the hospital

Comment [p57]: Faulty subject-verb Agreement

was → were

Comment [p58]: Incorrect verb forms

escapes → escape

Comment [p59]: Incorrect noun number

group → groups

Comment [p60]: Misspelled words

psd → PSD

Comment [p61]: Misspelled words

agaist → against

Comment [p62]: Misspelled words

psd → PSD

Comment [p63]: Pronoun use

it → they

Comment [p64]: Faulty subject-verb Agreement

was → were

Comment [p65]: Incorrect noun number

rabbit → rabbit's

Comment [p66]: Misspelled words

psd → PSD

Comment [p67]: Incorrect noun number

complication → complications

Comment [p68]: Confused words

aurous → aureus

Comment [p69]: Incorrect noun number

organisms → organism

yielded only *Staphylococcus* spp. and *E. coli* [46]. The wounds created in this research healed within twenty-one days which indicated a process of normal acute wound healing, and therefore justified the isolation of the two organisms (*Staphylococcus* and *Pseudomonas*) that have been previously reported to be common in acute wounds [47].

It was concluded that alloxan-induced diabetes causes delayed wound healing at all the phases of wound healing in NZW rabbits and *Staphylococcus* and *Pseudomonas* species are the predominant microorganism isolated in the acute wounds of diabetic and non-diabetic NZW rabbits. It is recommended that other microbial organisms such as fungal, viral and parasitic organisms associated with wounds of diabetic animals should be evaluated.

Comment [p70]: Incorrect noun number

microorganism → microorganisms

Ethical approval:

All authors hereby declare that "Principles of laboratory animal care" (NIH publication No. 85-23, revised 1985) were followed, as well as specific national laws where applicable. All experiments have been examined and approved by Ahmadu Bello University Committee on Animal Use and Care (ABUCAUC), (Reference number: ABUCAUC/2019/028)

References

1. Hosgood G, Wound repair and specific tissue response to injury. In: Slatter D. editor. Textbook of small animal Surgery; (3rd ed. Philadelphia; Saunders Elsevier Sciences. 2002; 66 – 86.
2. Schultz GS, Chin GA, Moldawer L, Diegelmann R.F. Principles of Wound Healing. In: Mechanisms of Vascular Disease (A Reference Book for Vascular Specialists): by Robert Fitridge and Matthew Thompson. University of Adelaide Press. 2011. Available [https://www.jstor. Org /stable /10.20851 /j.ctt1sq5w](https://www.jstor.org/stable/10.20851/j.ctt1sq5w) 94.27. Downloaded; 31st March 2020.
3. Sharma AK., Sunder V, Yashavardhan MH, Shukla SK. Wound healing: current understanding and future prospect. International Journal of Drug discovery. 2017; 8(1): 240-246. Available from: [https:// bioinfopublication. Org /jouarchive. php?opt=&jouid=BP J0000224](https://bioinfopublication. Org /jouarchive.php?opt=&jouid=BP J0000224)
4. Pavletic MM. Introduction to wound healing and wound management. In Proceedings of the American Animal Hospital Association. Lakewood, CO: AAHA; 1985
5. Lawrence WT. Physiology of the acute wound. Clinics in Plastic Surgery. 1998; 25 (3): 321– 340.

6. Pradham L, Nabzdyk CS, Lorgefo FW, Veves A. Expression of neuropeptide and Cytokines in a rabbit model of diabetic neuroischemic wound healing. *Journal of Surgical Research*. 2013;167: 336 – 342.
7. Sharma RV, Dave V, Sharma S, Jain P, Yadav S. Experimental Models on Diabetes: a comprehensive review; *International Journal of Advances in Pharmaceutical Sciences*. 2013; 4: 01-08. <http://www.arjournals.org/index.php/ijaps/index>
8. Nagata M, Suzuki W, Ilzuka S, Tabuchi M, Maruyama H, Takeda S, Aburada M, Miyamoto K. Type-2 diabetes mellitus in obese mouse model induced by monosodium glutamate. *Experimental Animal*. 2006; 55(2): 109 – 115.
9. Oriaghan, EA, Inegbenebor V, Shelu OJ, Obhimon O, Edono EO, Ekhoye I. (2012).The effect of monosodium glutamate on blood glucose in adult rabbits as model. *International Journal of Basic, Applied and Innovative Research*, 1(1): 10 –18.
10. Nelson RW. *Canine diabetes mellitus*. In: *Ettinger, SJ; Feldman, EC, Textbook of Veterinary Internal Medicine, 7th edition* Saunders publication: St. Louis, Saunders. 2010: 1782-1796.
11. Reusch, C. Feline diabetes mellitus. In: *Ettinger SJ, Feldman EC. editors. Textbook of Veterinary Internal Medicine, 7th edition*. St. Louis, Saunders; 2010; 1796 –1816.
12. World Health Organisation. Screening for type-2 diabetes. *Report of the World Health Organization and International Diabetes Federation Meeting*. Geneva. (2003)
13. Chinenye S. and Young K. The state of diabetes care in Nigeria: a review. *The Nigeria Health Journal*. 2011; 11(4): 101 – 106.
14. Mattin MJ, O'Neill DG, Church DB, McGreevy PD, Thomson PC, Brodbelt, DC.(2014). An epidemiological study of diabetes mellitus in dogs attending first opinion practice in the UK. *Veterinary Microbiology*, 174(14): 349.
15. Gani E Ihedioha JI. Evaluation of the prevalence of diabetes mellitus and impaired glucose homeostasis in dogs presented for veterinary care in Warri, Delta State Nigeria. *Book of Abstract, 6th /53rd NVMA Annual Congress of Nigeria Veterinary Medical Association (NVMA)*. 2016; 82.
16. Mirza R, Koh TJ. (2011). Dysregulation of monocyte/ macrophage phenotype in wounds of diabetic mice. *Cytokine* 56 (2): 256 – 264.
17. DeClue CE, Shornick LP. The cytokine milieu of diabetic wounds. *Diabetes Management*. 2015; 5(6): 525-537.

18. Kidman K. Tissue repair and regeneration: The effects of diabetes on wound healing. *The Diabetic Foot Journal*. 2008; 11(2): 73 – 80.
19. Okonkwo UA, DiPietro LA. (2017). Diabetes and wound angiogenesis. *International Journal of Molecular Sciences*, 18, 1419: 1–15.
20. Breena A, Redmand MD, Dockeny P, óBrien P and Pandit, A. Assessment of wound healing in the alloxan induced diabetic rabbit-ear model. *Journal of Investigative Surgery*. 2008; 21(5): 261-269.
21. Oguntibeju OO. (2019). Medicinal plants and their effects on diabetic wound healing. *Veterinary World*, 12(5): 653 – 663.
22. Hirsch T, Malte S, Baraa Z, Till K, Magdalena F, Hans-Ulrich S, Elof E. Enhanced susceptibility to infections in a diabetic wound healing model. *Biomedical Central Surgery*. 2008; 1471– 2482.
23. Brem, H. and Tomic-Canic, M. Cellular and molecular basis of wound healing in diabetes. *Journal of Clinical Investigation*. 2007; 117: 1219 – 1222.
24. Sorg Lipowsky HH, Gao L, Lescanic A. Shedding of the endothelial glycocalyx in arterioles, capillaries, and venules and its effect on capillary hemodynamics during inflammation. *The American Journal of Physiology: Heart and Circulatory Physiology*. 2011; 301(6): H2235 – 45.
25. Singh MP, Pathak K. Animal models for biological screening of anti-diabetic drugs: An overview. *European Journal of Experimental Biology*. 2015; 5(5): 37 – 48.
26. Sun WT, Lei CL, Bi CC, Chen ZL, Zhang L. Effect of alloxan time-administered drug on establishing diabetic rabbit model. *International Journal of Ophthalmology*. 2010; 3(3): 200 – 202.
27. Wang J, Wan R, Mo Y, Zhang Q, Sherwood LC, Chien S. Creating a long-term diabetic rabbit model. *Experimental Diabetes Research*. 2010; 75 – 95.
28. Flecknell PA. Anaesthesia. *In: Manual of Rabbit Medicine and Surgery*. British Small Animal Veterinary Association. 2000; 103-116.
29. Martin M, Kirsipuu V. (2016). *Rabbit anaesthesia*. Cornell University Institutional Animal Care and Use Committee: P. 1-6.

30. Fletcher J. Development of a new wound assessment form. *Wounds*. 2010; 6(1): 92 – 8.
31. Kiliç S, Timurkaan N, Unsaldi S, Günay C, Stek Ö, Yılmaz B. Comparison of effects of some wound healing materials on full thickness skin wounds in rabbits. *Turkish Journal of Veterinary and Animal Sciences*. 2002; 26(2): 263 – 272.
32. Hayat A, Temamogullari F, Yılmaz R, kerabulat O. Effects of Equisetum arvense on wound contraction of full thickness wound in rabbits. *Journal of Animal and Veterinary Advances*. 2011b; 10 (1): 81 – 83.
33. Sadeh F, Saleem R, Ahmed M., Ahmed SI, Navaid-ul-zafer. Healing potential of cream containing extract of *Sphaeranthus indicus* on dermal wounds in guinea pigs. *Journal of Ethnopharmacology*. 2006; 107 (2): 161 – 3.
34. Bridson EY. *The oxoid manual* 9th ed. Oxoid Limited, Hampshire, England; 2006. <http://www.oxoid.com>
35. Cheesbrough, M. *District laboratory practice manual in tropical countries* (part 2). Cambridge University Press, London. 2000; 178 – 179.
36. Ochei and Kolhatkar (2000). *Michael laboratory science, theory and practice* (2nd.ed). Tata McGraw-Hill Publishing Company Ltd, New Delhi, India: Pp. 1 – 349.
37. Markey B., Leonard, F., Archambault, F., Cullinane, M., Cullinane, A., and Maguire, D. *Clinical Veterinary Microbiology* (2nded). MOSBY Elsevier; London; Pp.1 – 857.
38. Lenzen S. The mechanisms of alloxan- and streptozotocin-induced diabetes. *Diabetologia*, 2008; 51 (2): 216 – 226.
39. Lipowsky HH, Gao L, Lescanic A. Shedding of the endothelial glycocalyx in arterioles, capillaries, and venules and its effect on capillary hemodynamics during inflammation. *The American Journal of Physiology: Heart and Circulatory Physiology*. 2011; 301(6): H2235 – 45.
40. World Union of Wound Healing Societies (WUWHS) Consensus Document Wound exudate: effective assessment and management, *Wounds International*. 2019; 1–35.
41. Weller C. Interactive dressings and their role in moist wound management. *Advanced Textiles for Wound Care*. 2009; 97–113.
42. Cutting KF. Wound exudate: composition and functions. *British Journal of Community Nursing*. 2014; 1-9.

43. Pavletic, M. M. Atlas of small animal woundmanagement andreconstructive surgery 3rd ed. Wiley-Blckwell (A John Wiley & Sons, Inc., Publication)Iowa, USA; 2010: 1 – 669.
44. Hayat AS, Khan AH, Masood N, Shailch N. Study for microbiological pattern and in vitro antibiotic susceptibility in patients having diabetic foot infections at tertiary care hospital in Abbottabad. World Applied Scientific Journal. 2011a; 12(2): 123 – 131.
45. Oates AL, Frank JMB, Andrew B, McBain AJ. (2012). Molecular and culture- based assessment of the microbial diversity of diabetic chronic foot wounds and contralateral skin sites. *Journal of Clinical Microbiology*, 50 (7): 2263 – 2271.
46. Daniel JC,Sowmiya SE, Gowthami E. Isolation and identification of bacterial pathogens from wounds of diabetic patients; International Journal of Current Microbiology and Applied Sciences. (2013);2(11): 72 – 77.
47. Bowler PG. and Davies, BJ. The microbiology of acute and chronic wounds. Wounds. 1999; 11(4): 72 – 78.




Fig. 1. The 3 cm² full-thickness wound (arrow) created at the aseptically draped back of the New Zealand White Rabbit.



Fig. 2a. Marking of wound diameter using a transparent tracing paper (arrow).

UNDER PEER REVIEW



Fig. 2b Measurement of wound diameter using a meter rule (arrow).

UNDER PEER REVIEW



Fig 3. Calculation of the size and epithelial tissues in the wound using a tracing paper in a graph sheet.




Fig. 4. The mean \pm SEM blood glucose concentration in rabbits of experimental groups ($P \leq 0.05$).

Key:

Group A = Non-Diabetic and Non-Wounded (NDNW).

Group B = Diabetic and Non-Wounded (DNW).

Group C = Wounded and Non-Diabetic (WND).

Group D = Wounded and Diabetic (WD).




Fig. 5. The Mean \pm SEM wound exudates value inc and d groups of experimental rabbits ($P \leq 0.05$).

Key:

Group C=Wounded and Non-Diabetic (WND).
Group D = Wounded and Diabetic (WD).




Fig. 6. The Mean \pm SEM percent wound epithelialization in the c and d groups of the experimental rabbits ($P \leq 0.05$).

Key:

Group C=Wounded and Non-Diabetic (WND).
Group D = Wounded and Diabetic (WD).




Fig. 7. The Mean \pm SEM percent wound contraction in the c and d groups of experimental rabbits ($P \leq 0.05$).

Key:

Group C=Wounded and Non-Diabetic(WND).

Group D = Wounded and Diabetic (WD).



D4 - 21



Fig. 8. The Wound of the C and D Rabbit Groups on Psd21 at the terminal Stage of Epithelialization and Contraction (sizes of the wounds reduced more than half (arrows))

Key:

Group C=Wounded and Non-Diabetic(WND).

Group D = Wounded and Diabetic (WD)

C₃_21= Rabbit number 3 in group C on psd 21

D₄_21= Rabbit number 4 in group D on psd 21




Fig. 9. The Wound of the C and D Rabbit Groups on Psd28 with Complete Contraction, and Scar Formation (arrows).

Key:

Group C=Wounded and Non-Diabetic(WND).

Group D = Wounded and Diabetic (WD).

C₃_28 = Rabbit number 3 in group C on psd 28

D₃_28 = Rabbit number 3 in group D on psd 28



Biochemical test	<i>Staphylococcus aureus sub. aureus</i>	<i>Staphylococcus epidermidis</i>	<i>Pseudomonas spp</i>
Grams reaction	+ cocci	+ cocci	- rod
Catalase test	+	+	+
TSI test	**	**	-
Indole test	**	**	-
MR test	**	**	-

Table 1.
The biochemical reactions of the bacterial organisms

isolated and identified in the wounds of the rabbits in groups C (WND) and D (WD).

Citrate test	**	**	+
Urease test	-	+	-
Mannitol test	+	-	**
Vp. Test	-	**	-
Oxidase test	-	-	+
Coagulase test	+	+	**
Haemolysis test	+	-	+
Arginine test	+	-	**
Maltose test	+	+	-

Keys:

WND = Wounded and Non-Diabetic

WD = Wounded and Diabetic

+ = positive

- = negative

** = no test carried out

MR = methyl red.

UNDER PEER REVIEW

