

### Post Harvest Treatments and Shelf Life of Some Tropical Fruit.

#### Abstract

*Fruits are perishable horticultural crops grown in tropical and sub-tropical areas of the world. It is a richest source of micro nutrients and phyto chemicals that contributed a significant role in human health. Mango, avocado, papaya, guava and pine apple are popular and economically important tropical fruits due to their essential nutritional properties and appreciated eating quality. However, in developing countries, upto 50% of those fruits are lost after harvest due to improper post-harvest handling practices. Edible coating, dip in calcium chloride solution, hot water treatment and drying are easily applicable post harvest treatments used in the preservation of such tropical fruits. Shelf life of mangoes coated using edible coating material can be extended up to 18 days without rotting and developing unpleasant-flavour relative to coated guava, papaya, avocado and pine apple fruits. Dipping harvested fruits in various concentration of calcium chloride solution exhibited different storage life. Consequently, mango fruits treated with 10% of calcium chloride solution preserved for 30 days at 12° C and 85-90% relative humidity. Guava fruits treated by hot water at 45°C for 3 minute lasts upto 31 days under room temperature. In this review maximum shelf life up to 9 months was found for Avocado fruit dried using freeze drying method.*

**Keywords:** Fruits, Shelf life, Edible coating, Calcium chloride, Hot water treatment, Drying

#### INTRODUCTION

Fruits are Perishable horticultural crops that contain high amount of water (75-90%) by wet basis. Most fruits are grown in tropical and sub-tropical areas of the world and are used of income generation for those who engaged in its production. Global production of tropical fruits were estimated to 100 million tonne since 2018 with increasing more than 8 million

tonne from 2017 production year (Altendorf, 2019). From this total of world tropical fruit production developing countries share estimated to 99 % of production annually.

Most tropical fruits have played a significant role in human diet in terms of nutritive and medicinal values. Fruits are also a richest source of micro nutrients (vitamins and minerals) as well as phytochemicals. Mango, avocado, papaya, guava and pine apple are popular and economically important tropical fruits due to their nutritional composition (vitamins, minerals, fiber and other phytochemical compounds) and appreciated eating quality (attractive colour, sweet taste and pleasant flavour).

Major limitation of these tropical fruits on marketing, storage and transportation are extensive post harvest loss and shorter shelf life. Post harvest losses of fruits are occur at all stages of post harvest operation, starts immediately after the time of harvest to its final destination due to improper post harvest handling practices (Anjichi et al., 2006). Thus, Post harvest losses of fruits are estimated 5-20% in developed countries and 20-50 % in developing countries (Mashav, 2010). Therefore, harvested fruits could be subjected to different post harvest treatments in order to preserve and ensure availability of such tropical fruits during marketing, off season and storage.

In developing countries, application of several advanced post harvest loss reduction mechanisms are important for mitigating food insecurity. Post-harvest treatments will slow down the physiological processes in fresh fruits such as respiration, senescence and ripening to prolong the shelf life of fresh commodities (Sandarani *et al.*, 2018). Plenty of technologies like Hot water treatments, edible coating, post harvest thermal treatments, dipping in food grade chemicals and other non thermal treatments have been applied on several fruit for preserving their physicochemical quality and reducing losses during post harvest life. Therefore, this article is aimed to summarize some post harvest treatments applied on selected tropical fruits and its impacts on their respective shelf life.

### **Post harvest physiology of fruits**

Harvested fruits are considered as a living organism because their metabolic functions continues during post harvest life (Kays, 1991). These fruits under go different physiological

processes like respiration, ripening and senescence that must be controlled to extend products shelf life. Improper management of such metabolic processes of harvested fruits during storage can contribute to a major loss in both quantitative and qualitative. Respiration is the metabolic process which utilizes oxygen to breakdown stored organic materials (carbohydrates, proteins, and fats) into simple end products with a release of energy. Increased respiration rate results in accelerated ripening and senescence that shorten product shelf life.

There are possibilities of extended product storage life by modification of the internal gas composition of storage atmosphere that minimizes the fruit's respiration rate. Ripening is part of the natural senescence of the whole fruits. It is an irreversible process that contributes to changes in chemical composition, skin color, total soluble solids, flavour and texture. It improves the fruit eating quality and shortens the post harvest life. Ripening and natural senescence of fruits can be delayed by lower temperature, elimination of mechanical damage and reducing ethylene production (Yahia *et al.*, 2006a).

### **Shelf life**

Shelf life is the period of time that food products retained its microbial safety, nutritive value and sensorial attributes begins from the time the food is manufactured to its start of deterioration. During this period, a food product remained acceptable without significantly, losing its sensory, chemical, physical, functional, microbiological and nutritional properties under different radiation, gas concentration, redox potential, packaging material and storage condition (Singh, 2005). Proper temperature and relative humidity are critical to obtain an extended shelf-life of stored fruits. When harvested fruits stored at higher than critical temperatures set for recognized produce, the storage life will be shortened. Similarly, if harvested fruits are stored at a lower than critical temperatures set by scientific organization for selected produce, their shelf life could be affected by either freezing or chilling injury.

## Edible Coating

Edible coatings are thin layers of edible materials and environmentally safe technology that can be applied by dipping, spraying or painting on different fruits and vegetables. It controls moisture transfer, oxygen or carbon dioxide migration and oxidation processes to improve the mechanical integrity or handling characteristics of the food (O'Sullivan *et al.*, 2006). In addition, edible coatings help to preserve color, texture and volatile compounds of fresh fruits, maintain the structural integrity and protect against mechanical damages (Dhall, 2013). Edible coatings have a major advantage of extending the shelf life of food products, control material exchange, enhancing safety, improve nutritive value, sensory attributes and attractiveness of food products because of active ingredients incorporated to food matrix during coating (Kokoszka and Lenart, 2007 ; Bodini *et al.*, 2013).

The success of edible coatings for fresh products totally depends on the control of internal gas composition (Dhal, 2013). Several specific requirements described for edible coating by Arvanitoyannis and Gorris, (1999) are manifested that it could have good barrier to water, not deplete oxygen or build up excessive carbon dioxide, A minimum of 1–3% oxygen is required to avoid a shift from aerobic to anaerobic respiration, reduce water vapor permeability, carry active agents (antioxidants, vitamins, etc.) and retain volatile flavor compounds, melt above 40°C without decomposition, easily emulsifiable, never interfere with the quality of fresh fruit, should have low viscosity and be economical.

Different types of edible coatings can be prepared from protein, carbohydrate, lipid materials and their derivatives. Edible coating also contain food grade plasticizers, additives, antioxidants and antimicrobial compounds that can be applied to many fruits and fruit products. Carbohydrate based edible coatings like polysaccharides, starch, modified starch, chitosan and pectin are commonly used among natural bio polymer. Edible coatings processed from cassava starch applied to strawberries were contributed efficiently in reducing respiration rate, delaying weight loss, firmness loss and retained its sensorial acceptability during storage period (Garcia *et al.*, 2011).

Protein based edible coatings have good film forming properties and good adherence to hydrophilic surfaces. Lipid based edible coatings are the hydrophobic, block the transport of

moisture and water vapor, improve surface appearance, maintain quality attributes and prolong the post harvest life of some fruits. Waxes are the most efficient lipid based edible coatings substances to decrease the water vapor pressure and was the first coating used in fruits specially in citrus fruits.

Increased shelf life at least three days relative to uncoated fruit were reported by Gallo *et al.*, (2003) for guava fruit coated in immersion solution consisting between 5 and 10% of Potato starch, sodium alginate, carragenan, pectin and glycerol between 10 and 30% at 50°C, then dried at 50°C for 30 minutes stored at 25°C and 50- 70% Relative humidity. The potato starch and pectin based coatings are highly efficient and good in the preservation of the sensorial characteristics of the fruit (size, yellow colour and aroma) with 15 days (Gallo *et al.*, 2003).

Shelf life of waxed mango fruit exceeds upto four days than non waxed mango fruit exhibited shelf stable upto 15 days that stored at room temperature ( Mandal *et al.*, 2018). According to research finding by Srinivasa *et al.* (2002) mango fruits stored in chitosan covered boxes indicated an extension of shelf life up to 18 days and without any microbial growth and off-flavour. Extended shelf life upto 13 days was recorded for Papaya fruits treated with 5% of wax emulsion while 9 days noted for untreated papaya fruit stored at room storage condition (srinu *et al.*, 2017).

Avocado fruits coated with gelatin-corn starch films have shown increased in shelf-life by reducing moisture migration, gas exchange and delaying over ripening (Aguilar-Mendez *et al.*, 2008). In addition, pine apple treated using Alginate-based edible coating containing 0.3 percent (w/v) lemongrass showed extended shelf life for 10 days by arresting physiological changes of pine apple fruits stored at 10°C and 65 percent relative humidity (Azarakhsh *et al.*, 2014). Gellan edible coating composed of gellan gum 0.56 % (w/v), glycerol 0.89 % (w/v) and sunflower oil 0.025 % (w/v) considerably lowers the respiration rate, reduces weight loss, and preserves the stiffness, colour, and sensory attributes of freshly sliced pineapple in low-temperature conditions. After 16 days of storage at 5 °C, there was no significant difference in the pH, titratable acidity, and TSS of coated and untreated samples. Cut pineapple can be kept fresh for a long time in the refrigerator with the help of the gellan edible coating solution (Azarakhsh *et al.*, 2014)

## **Calcium chloride treatments**

Post harvest application of calcium may delay senescence in fruits with no detrimental effect on consumer acceptability. Calcium chloride application on harvested fruits could have valuable role to prevent mechanical and physiological damage on fruit by making fruit cell walls hard and inhibit enzyme production of fungi and bacteria that cause fruit rotting (Mishra, 2002 and Huang et al, 2012). Shelf life upto 15.32 days was observed for papaya fruits treated with 3.0% of  $\text{CaCl}_2$  as compared to untreated fruits attained shelf life of 9 days (Srinu *et al.*, 2017). Similar research report by priyanka et al. (2012) showed that Papaya fruits treated in  $\text{CaCl}_2$  at 2 percent extended the shelf life of fruit more than 9 days of storage.

In other way, Guava fruits treated for 2 to 4 minutes in 1% and 2% of calcium chloride recorded a potential shelf life of more than 9 days under room storage condition (Manmohan *et al.*, 2019). According to Galvis *et al.*, (2003) extended shelf life upto 30 days was noted for mango fruits dipped in 10% of calcium chloride solution at ambient temperature and stored at  $12^\circ\text{C}$  and 85-90% relative humidity. Similarly, Calcium chloride ( $\text{CaCl}_2$ ) treatment significantly reduced relative weight loss, delayed the development of purple skin colour, blemishes, disease incidence and severity in avocado fruits stored for 18 days compared to untreated control (Mutui et al., 2011) .

Calcium chloride ( $\text{CaCl}_2$ ) can also change intracellular and extracellular processes that can slow down fruit ripening, maintain cell walls so that it can inhibit fruit softening, and reduce mechanical and microbiological damage (Hasmoro, 2014). Mold incidence and severity appeared on pineapple dipped in the  $\text{CaCl}_2$  solution after the 16th day of storage but it exhibited better sensory properties, particularly on texture, color, and overall acceptance After storage for 40th days (Tamalea *et al.*, 2021). Extended shelf life was reported by Youryon and Wongs (2013) for Pineapple fruits immersed in 1%, 2% and 3% of  $\text{CaCl}_2$  solution and kept under mild low temperature storage condition.

## **Hot water treatment**

Hot water treatments are post harvest thermal treatments that applied on fruits by immersion, rinsing and brushing at desired temperature ranges and duration for retaining quality and

microbial safety of fresh fruits. It is effective, particularly for fungal pathogen control because fungal spores and latent infections are either on the surface layers or under the peel of the fruit. The application of hot water treatments on different fruits have been increased in order to disinfestations of insect pests, prevent fungal rots and increase fruit resistance to chilling injury (Lurie, 1998).

The use of hot water as a disinfestations and maintaining quality of fresh fruits are affected by number of factors including maturity stage, cultivar type, temperature and duration (Djioua *et al.*, 2009). Many fruits tolerate exposure to water temperatures of 50–60°C for up to 10 min, but shorter exposure at these temperatures can control many post harvest plant pathogens (Barkai- Golan and Phillips, 1991). In contrast, hot water dips for fruit require 90 min exposure to 46°C. Several Authors reported that hot water treatment of fruits at different temperatures for several minutes to hour were applied for maintaining fruit quality during marketing and storage (Garcia *et al.*, 1995a ; kumah *et al.*, 2011; Poubou *et al.*, 2018).

Hot water treatment of mango fruits resulted reduced post harvest decay, improved color, appearance of mango fruits (Prusky, 1999). Mango fruit preliminarily treated with hot water significantly maintained firmness during 7 days of storage at ambient conditions (Pholoma *et al.*, 2020). Hot water treatment could reduce disease incidence and maintain postharvest quality during storage and prolong the shelf-life of papaya fruit ( Li *et al.*, 2013). Extended storage stability and marketability was reported for Pineapple and avocado fruits treated by hot water at different temperature and time combination (Munhuweyi *et al.*, 2020; Minh, 2021). Related research finding revealed that Guava fruits treated with hot water at 45°C for 3 minute lasts upto 31 days under room temperature of 5±2 °C and 85 to 90 per cent relative humidity (killadi *et al.*, 2021).

### **Drying**

Drying is one of the oldest, simple, cost effective, safe and wide spread methods of food preservation. It removes a large amount of water contained in products so that the growth of micro-organisms and reaction lead to deterioration are inhibited(Doymaz, 2008). It also reduces the weight and bulk of fruits which reduces transport and storage costs. Successful and appropriate fruit drying techniques produce healthy dried fruit products with good flavor, texture and color. Despite, drying does not improve food quality to obtain desired preservation and good quality of dried products sort and select the best-quality of fruits for drying.

**Solar drying** is the most effective method of drying foods using solar energy and in use in many parts of the world (Esper and Mühlbauer., 1998). It is cheaper as it has little or no equipment costs and the produce has to spread on suitable surface and allowed to dry in the sun (Mnkeni et al., 2004). Fruits dried in the solar dryer are placed on trays made of screen or wooden dowels, stainless steel, teflon coated fiberglass or plastic. Solar drying technology seems to be one of the most promising alternatives to reduce the post-harvest losses, extend shelf life with much preserved colour and texture (Mulokozi and Svanberg, 2003; Temu et al., 2008; Wiriya et al., 2009). Mangoes and pineapples dried in different solar drying and stored in low and high density polyethylene packaging containers could last up to more than six months (Mongi, 2013).

**Freeze-drying**, also known as lyophilisation, or cryodesiccation, is a water removal process typically used to preserve a perishable Agricultural commodity or make the commodities more convenient for transport and storage. Freeze-drying works by freezing the material and then reducing the surrounding pressure to allow the frozen water in the material to sublime directly from the solid phase to the gas phase. Freeze drying (FD) is the best method of dehydration as it gives a final product of the highest quality without heat compared to other methods of food drying (Irzyniec *et al.*, 1995). Different fruits dried under freeze drying method exhibited improved color and maximum retention of vitamin C, chlorophyll, flavor and taste ( Hu *et al.*, 2006; Marques *et al.*, 2006; Jangam *et al.*, 2008) .

Fruits dried under modern freeze dryer have a longer shelf life without releasing as much of their nutritional content and sensory characteristics (Valentina et al., 2016). High-quality with preserved color of Pollock avocado powder to be used in various food application was obtained using freeze drying method (Mujaffar and Dipnarine, 2020). Avocado fruit dried using freeze drying method have a shelf life of up to 9 months with exhibited slight changes in nutritional quality and sensorial properties (CastanedaSaucedo *et al.*, ; Husen *et al.*, 2014)

According to patel *et al.*, (2016) guava fruits were effectively preserved for long time with low adverse health effects when subjected to freeze drying condition. Similar research finding reported by Jangam *et al.*,(2008) revealed that freeze-dried mangoes, avocados and pineapples retained their sensory characteristics and shelf stabilities. The very low water content of freeze dried fruits helps them to have a longer shelf life (Shishegharha *et al.*,

2002). However, freeze dried fruits have a high moist air absorption tendency and additional barrier to water absorption solutes such as Arabic gum, tricalcium phosphate and maltodextrin should be added to increase its shelf stability (Ceballos *et al.*, 2012; Mosquera *et al.*, 2012).

### **Conclusion**

Application of different post harvest treatments to fruits are highly encouraged to preserve and reduce their post harvest loss. Appropriate edible coating materials, hot water treatment, dip in optimum concentration of calcium chloride solution and drying under suitable drying methods are easily applicable post harvest treatments used in the preservation of perishable tropical fruits. Those post harvest treatments have considerable role in ensuring fruits shelf stability, up to 9 months.

### **REFERENCE**

Aguilar-Mendez, M., San Martin-Martinez, E., Tomas, S., Cruz-Orea, A., and Jaime-Fonseca, M.(2008). Gelatine–starch films: Physicochemical properties and their application in extending the post-harvest shelf life of avocado (*Persea americana*). *Journal of the Science of Food and Agriculture*, 88, 185–193.

Altendorf, S. 2019. Major tropical fruits market review .2018. Rome, FAO (Food and Agriculture Organization of the United Nations).

Anjichi VA, Okong'o OM, Opile WR and Muasya RM. 2006. Effect of 'Purafil' ethylene absorber on post-harvest quality of 'Fuerte' avocados. *Proceedings of the V workshop on sustainable Horticultural production in the tropics*. Njoro, Kenya. 23rd -26th Nov 2005, pp 220-225.

Arvanitoyannis, I. and Gorris, L. G. M. 1999. Edible and Biodegradable Polymeric Materials Food Packaging or Coating. **In:** *Processing Foods: Quality Optimization and Process Assessment*, pp. 357–371. CRC Press, Boca Raton, Florida.

Azarakhsh N, Osman A, Ghazali HM, Tan CP, Adzahan NM. 2014. Effects of gellan-based edible coating on the quality of fresh-cut pineapple during cold storage. *Food and Bioprocess Technology* ; 7(7):2144-2151.

Azarakhsh N, Osman A, Ghazali HM, Tan CP, Adzahan NM.2014. Lemongrass essential oil incorporated into alginatebased edible coating for shelf-life extension and quality retention of fresh-cut pineapple. *Postharvest Biology and Technology*. 88:1-7

Barkai-Golan, R., Phillips, D.J., 1991. Post harvest heat treatment of fresh fruits and vegetables for decay control. *Plant Dis*. 75, 1085–1089

Bodini R, Sobral P, Favaro-Trindade C, 2013. Properties of gelatin-based films with added ethanol-propolis extract. *LWT-Food Sci Technol*. 51:104-10

Castaneda-Saucedo,M., Valdés,E., Tapia,E., Delgado, A., Bernardino-García, A., Rodríguez-Ramírez, R., Ramirez-Anaya, R. 2014. Effect of freeze-drying and production process on the chemical composition and fatty acids profile of avocado pulp.

Ceballos A, Giraldo G, Orrego C .2012. Effect of freezing rate on quality parameters of freeze-dried soursop fruit pulp. *J. Food Eng*. 111(2):360-365

Dhall, R.K., 2013. Advances in edible coatings for fresh fruits and vegetables: a review. *Critical reviews in food science and nutrition*, 53(5), pp.435-450.

Djioua, T., Charles, F., Lopez-Lauri, F., Filgueiras, H., Coudret, A., Freire Jr, M., Ducamp-Collin, M.N. and Sallanon, H. 2009. Improving the storage of minimally processed mangoes (*Mangifera indica* L.) by hot water treatments. *Postharvest Biology and Technology*, 52(2), pp.221-226.

Doymaz, I. 2008. Influence of blanching and slice thickness on drying characteristics of leek slices. *Chem. Eng. Process*, 47: 41-47.

Esper, A. and Mühlbauer, W., 1998. Solar drying-an effective means of food preservation. *Renewable energy*, 15(1-4), pp.95-100.

Gallo, J.Q., Amaro, M.D., Cabrera, D.M.B.G., Alvarez, M.C., Debeaufort, F. and Voilley, A., 2003. Application of edible coatings to improve shelf-life of Mexican guava. *Acta Horticulturae*, pp.589-594.

Galvis, J.A., Arjona, H. and Fischer, G., 2003. The effects of applying calcium chloride solution (CaCl<sub>2</sub>) on Van Dyke mango fruit (*Mangifera indica* L.) storage life and quality.

- Garcia, J.M., Aguilera, C., Albi, M.A., 1995a. Post harvest heat treatment on Spanish strawberry (*Fragaria X ananassa* cv Tudla). *J. Agric. Food Chem.* 43, 1489–1492.
- Garcia, L.C., Pereira, L.M., Sarantopoulos, C.I.G., and Hubinger, M.D., 2011, Effect of antimicrobial starch edible coating on shelf life of fresh strawberries. *Packag. Technol. Sci.*, 25: 413-425
- Hasmoru, H.B.S., Trisnowati, R. Rogomulyo. 2014. Effects of CaCl<sub>2</sub> on the Degradation of the Framework and Age of Buah Sawo (*Manilkara zapota* L.) van Royen). *Vegetable Journal*, 3(4): 52-62.
- Hu Q, Zhang M, Mujumdar AS, Du W & Sun J .2006. Effects of different drying methods on the quality changes of granular edamame. *Drying Technology*, 24, 1025–1032.
- Huang, S.; G. Zhu, L. Qin, X. Zhou, F. Huang, Q. Li, W. Yan, H. Huang, Z. Cen, G. Fu, and C. Hu. 2012. Enhancement of Efficacy in Controlling Postharvest Decays and Extending Shelf Life of Mangoes by Combined Pre- and Post- harvest Chemical Applications, *International Journal of Agriculture and Biology*.14 (2) : 176-182
- Husen ,R. , Andou, Y., Ismail, A., Shirai, Y. Hassan, M.A. 2014 . Enhanced polyphenol content and antioxidant capacity in the edible portion of avocado dried with superheated-steam. *International Journal of Advanced Research* 2 no. 8, 241-248.
- Irzyniec, Z. Klimezak, J. and Michalowski, S. 1995. Freeze-drying of the black currant juice. *Drying Technology* 13: 417-412.
- Jangam SV, Joshi VS, Mujumdar AS & Thorat BN .2008. Studies on dehydration of sapota (*achras zapota*). *Drying Technology*, 26, 369–377.
- Kays, S. G. 1991. Post harvest physiology of perishable plant products, AVI, Newyork, USA, 266 PP.
- Killadi, B., Guarj, P.S., Lenka, J., Chaurasia, R. and Shukla, D.K., 2021. Optimization of hot water treatment of guava fruits cv.‘Shweta’to prolong cold storage. *International Journal of Chemical Studies*, 9(1), pp.410-415.
- Kokoszka S, Lenart A. 2007. Edible coatings-formation, characteristics and use: A Review. *Pol J Food Nutr Sci.* 57:399-404. 11.

Kumah, P., Appiah, F. and Opoku-Debrah, J.K., 2011. Effect of hot water treatment on quality and shelf-life of Keitt mango.

Li, X., Zhu, X., Zhao, N., Fu, D., Li, J., Chen, W. and Chen, W., 2013. Effects of hot water treatment on anthracnose disease in papaya fruit and its possible mechanism. *Postharvest biology and technology*, 86, pp.437-446.

Lurie, S., 1998. Postharvest heat treatments. *Postharvest biology and technology*, 14(3), pp.257-269.

Mandal, D., Sailo, L., Hazarika, T.K. and Shukla, A.C., 2018. Effect of edible coating on shelf life and quality of local mango cv. Rangkuai of Mizoram. *Research on Crops*, 19(3), pp.419-424.

Manmohan, S.B, BP Bisen, Rahul Dongre and Maneesh Kumar. 2019. Effect of post-harvest treatments on shelf life and physico-chemical changes of Guava fruits. *International Journal of Chemical Studies* 2019; 7(6): 3096-3099

Marques L.G, Silveira A.M and Freire J.T .2006. Freeze-drying characteristics of tropical fruits. *Drying Technology*, 24, 457-463.

Minh, N.P. 2021. Influence of hot water treatment to quality properties of pineapple (*Ananas comosus*) fruit during storage. *Food Research*, 5(5), pp.186-194.

Mishra, M. 2002. Lead Acetate Induced Citotoxicity in Male Germinal Cell of Swiss Mice. *Swiss*. p.291-294.

Mnkeni A.P, Soundy P, Brutsch MO .2004.. Solar drying of fruit and vegetables. internet document, Department of Agriculture. WWW. solar drying of fruit and vegetables. Htm.

Mongi, R.J., 2013. *Solar drying of fruits and vegetables: dryers' thermal performance, quality and shelf life of dried mango, banana, pineapple and tomato* (Doctoral dissertation, Sokoine University of Agriculture).

Mosquera L, Moraga G, Martínez N .2012. Critical water activity and critical water content of freeze-dried strawberry powder as affected by maltodextrin and arabic gum. *Food Res. Int.* 47(2):201-06.

Mujaffar, S. and Dipnarine, T.A., 2020. The production of a dried avocado (*Persea americana*) powder. In *Proceedings of the International Conference on Emerging Trends in Engineering and Technology*, St. Augustine (pp. 44-54).

Mulokozi, G. and Svanberg, U. 2003. Effect of traditional open sun drying and solar cabinet drying on carotene content and vitamin a activity of green leafy vegetables. *Journal of Plant Foods for Human Nutrition* 58: 1 - 15.

Munhuweyi, K., Mpai, S. and Sivakumar, D. 2020. Extension of avocado fruit postharvest quality using non-chemical treatments. *Agronomy*, 10(2), p.212.

Mutui, T.M., Kimeu, E.N. and Opile, W.R. 2011. Effect of postharvest treatments on the quality and shelf life of avocado (*Persea americana* MILL.). *African Journal of Horticultural Science*, 5.

O'Sullivan, A., Shaw, N.B., Murphy, S.C., Van de Vis, J.W., Van Pelt-Heerschap, H. and Kerry, J.P. 2006. Extraction of collagen from fish skins and its use in the manufacture of biopolymer films. *Journal of aquatic food product technology*, 15(3), pp.21-32.

Patel, P., Sunkara, R., Walker, L.T. and Verghese, M., 2016. Effect of drying techniques on antioxidant capacity of guava fruit. *Food and Nutrition Sciences*, 7(07), p.544.

Pholoma, S.B., Emongor, V. and Tshwenyane, S. 2020. Physicochemical attributes in mango fruit (*Mangifera indica*) as influenced by storage temperature and hot water treatment. *Journal of Experimental Agriculture International*, 42(1), 133- 141.

Poubol, J., Techavuthiporn, C. and Kanlayanarat, S., 2018. Guava fruit treated with hot water on microbiological quality of fresh-cut 'Kimju' and 'Pan Srithong' guava. *International Food Research Journal*, 25(3), pp.903-907.

Priyanka, S., Sanjay, K., Sutanu, M., Adesh, K. and Yadav, Y.C. 2012. Effect of calcium chloride on post harvest changes in papaya fruits. *Asian Journal of Horticulture*, 7(1), pp.113-117.

Prusky D., Y. Fuchs, I. Kobilier, I. Roth, A. Weksler, Y. Shalom, E. Fallik, G. Zauberman, E. Pesis, M. Akerman, O. Ykutiely, A. Weisblum, R. Regev, And L. Artes. 1999. Effect of hot water brushing, prochloraz treatment and waxing on the incidence of black spot decay caused by *Alternaria alternata* in mango fruits. *Postharvest Biology and Technology* 15:165–174

- Sandarani, M.D.J.C., Dasanayaka, D.C.M.C.K. and Jayasinghe, C., 2018. Strategies used to prolong the shelf life of fresh commodities. *J. Agric. Sci. Food Res*, 9(1), pp.1-6.
- Shishegarha, F. Makhlof, J. Ratti, C. 2002. Freeze-drying characteristics of strawberries. *Drying Technol.* 20(1):131-45.
- Singh, R.K. and Singh, N. 2005. Quality of packaged foods. In *Innovations in food packaging* (pp. 24-44). Academic Press.
- Srinivasa, P. C., Baskaran, R., Ramesh, M. N., Prasanth, K. V. H. and Tharanathan, R. N. 2002. Storage studies of mango packed using bio-degradable chitosan film. *European Food Res. and Technol.* **215** : 504-08.
- Srinu, B., Manohar, R.A., Veena, J.K., Narender, R.S. and Harish, K.S. 2017. Effect of different post-harvest treatments on quality and shelf life of papaya. *Journal of Pharmacognosy and Phytochemistry*, 6(5), pp.1788-1792.
- Tamalea, R.R., Haryanti, P. and Wibowo, C. 2021. Application of Dipping on CaCl<sub>2</sub> as a Postharvest Treatment to Maintain the Quality of Pineapple during Storage. *Asia Pacific Journal of Sustainable Agriculture, Food and Energy*, 9(2), pp.38-45.
- Temu, A., Chove, B. and Ndabikunze, B. 2008. Development of enterprise in solar drying of fruit and vegetables for employment creation. Communication workshop for DANIDA pilot projects in SUA, 21-25 July 2009, SUA, Morogoro, Tanzania, 15pp
- Valentina, V., A.R. Pratiwi, P.Y. Hsiao, H.T. Tseng, J.F. Hsieh and C.C. Chen. 2016. Sensorial characterization of foods before and after freeze-drying. *J. Aust. Food Sci.*, 1(6):1-5.
- Wiriya, P., Paiboon, T. and Somchart, S. 2009. Effect of drying air temperature and chemical pretreatments on quality of dried chilli. *International Food Research Journal* 16: 4 - 7.
- Yahia, E.M., Ornelas-Paz, I. de J. and Ariza, F.R. 2006a. *The Mango*. Editorial Trillas, México City, Mexico, 224 pp. (in Spanish)
- Youryon, P. and Wongs-Aree, C. 2013. Postharvest application of calcium chloride affects internal browning reduction during low temperature storage of 'Sawi' pineapple. In *II Southeast Asia Symposium on Quality Management in Postharvest Systems 1088* (pp. 197-200).