

BRONCHIAL FOREIGN BODIES: TELESCOPIC APPROACH IN PORT HARCOURT

ABSTRACT

Removal of foreign bodies from the airway, especially in children require special skill and proper instrumentation. In telescopic bronchoscopy airway accessibility is made easy and hence removal becomes less challenging, compared to other methods. Aim of this study is to showcase the telescope in the process of tracheobronchial foreign body removal.

Method

The Telescopic Bronchoscope forms the major armament for this study. This is in contrast to other types of bronchoscope used in accessing the tracheobronchial tree for foreign body extraction. The German Richard Wolf Telescopic Bronchoscope was used. Other components of this instrument include, the light source and light cable for excellent illumination. The fibre optic and rigid bronchoscopes were not used in this study.

Results

24 children under this study presented in University of Port Harcourt Teaching Hospital, either referred from other centres or presented on their own. Fourteen (14) male and (10) ten female with male/female ratio of 2:1. Foreign bodies retrieval were groundnut (38%), fish bone (21%), others were plastic toy pieces, screw nuts. The right main bronchus, (61%) recorded the highest lodgement.

Conclusion

Foreign body in the airway especially in children is life threatening. The use of telescope offers excellent armament, preventing mortality.

Keywords

Telescopic Bronchoscope, foreign body aspiration, tracheobronchial tree.

Introduction

Aspiration of foreign body in the tracheobronchial airway, especially in children is life threatening. Airway compromise, many at times result in mortality¹. The use of Telescopic Bronchoscope and endotracheal intubation assures easy removal, preventing mortality which at times follows the use of other methods. Frequent de-saturation during the procedure and the delay encountered in process of de-saturation are eliminated, thereby providing speed in the process. Therefore, length of time under anaesthesia and morbidity are drastically reduced. Uninterrupted free flow of anaesthesia gas(es) is ensured as endotracheal tube is already connected to the anaesthetic machine².

The Surgeon has more time to search and remove foreign body through this process. The instrument allows for good anaesthetic approach since normal Intermittent Positive Pressure Ventilation (IPPV) is not interrupted.

The aim of this study is to highlight and showcase the use of telescopic bronchoscope in the removal of tracheobronchial foreign bodies in comparison to other methods.

Patient and Method

This is a 6 year retrospective study of children who presented with airway obstruction from foreign body requiring removal at the University of Port Harcourt Teaching Hospital. The study was between September, 2014 to September, 2020. Clinic and theatre records were utilised and information include age, gender, history of aspiration of foreign body, nature of foreign body, time of aspiration and site of lodgement.

Instrumentation

Instrumentation was the telescope bronchoscope with general anaesthesia in form of endotracheal intubation, other armaments include light sources and light carrier to provide excellent illumination and clarity of the airway.

Telescopic Instrument

The instrument is a German, Richard Wolf, optique Lumina Telescopic Bronchoscope. It has adult and paediatric sizes.

Ref: 8464,30, SN 5000229228, CE 0124, Size 16 Paediatric. Also Ref 82514,431, SN 5000192374, CE 0124 Size 35 Adult. Both have Lumina barrel forcep, where the optique telescope is inserted and passed down for viewing the airway. The Lumina barrel forcep also allows metallic or rubber suction tubing to pass for suctioning of the airway. The optique telescope has attachment for light source to properly illuminate the airway. The lumina barrel has forcep at the tip for grasping of any object within the tracheobronchial passage as you advance. These are demonstrated in the picture shown below. Good and functional suction apparatus is an essential component of these armaments.

Pictures of Instrument



The telescopic bronchoscope



The light source and light cable

Anaesthesia

The anaesthetic management in this section applied to all the patient under this study. They were assessed preoperatively to identify any associated systemic illness. Premedication with intravenous atropine 0.01mg/kg body weight, paracetamol 15mg/kg, dexamethasone 0.1mg/kg to reduce airway oedema. All prior to induction of anaesthesia.

Induction commenced with ketamine 2mg/kg body weight or propofol 2mg/kg, suxamethanium 2mg/kg. Laryngoscopy performed and endotracheal tube passed. Usually, the endotracheal tube must be less than the normal size to the patient. This is to allow the Lumina barrel of the telescope pass thorough the rima glottis. (usually size 3.0mm tube preferable). Orotracheal tube passed and secured and ventilation continued.

Results

In this study we had (24) twenty four patients that had telescopic removal of foreign body from the tracheobronchial airway. Fourteen (14) were male, and ten (10) female, with male/female ratio of 2:1.

Classical history of choking, coughing bouts before respiratory distress following meal were recorded in majority of them. Others presented because parents suspected inhalation of foreign body.

Table I: Age and gender distribution of patients

Age group	Male	Female	Total	Percentage (%)
6-12 months	1	1	2	8
1-2 years	6	4	10	42
2-3 years	3	3	6	25
3-4 years	2	1	3	13
4-5 years	1	1	2	8

5 years and above	1	0	1	4
-------------------	---	---	---	---

Table II presents time interval between aspiration and presentation. This ranged from (6) hours and (10) ten days and above.

Table II: Types of foreign body aspirated

Time Interval	Number	Percentage (%)
<24 hours	9	38
1-5 days	12	50
5-10 days	2	8
10 days and above	1	4
Total	24	100

Nineteen (19) presented through referral from other health care providers, while others presented on observing symptoms. Type of foreign bodies removal were recorded on the table III.

Table III: Types of foreign body aspirated

Object Retrieved	No of Patients	Percentage (%)
Groundnut	9	38
Fish bone	5	21
Groundnut shell	2	8
Screw nut	3	13
Eraser	1	4
Toy Plastic Piece	1	4
Cellophane Piece	1	4
Toy Bulb	1	4

No foreign body seen	1	4
----------------------	---	---

Groundnut appear as the commonest foreign body removed, 9 (38%), followed by fish bone 5(21%) and screw nut. Others are eraser, toy materials and toy bulb.

TableIV: Location where foreign body was removal

Position	Number	Percentage (%)
Right main bronchus	14	61
Left main bronchus	9	39
Total	23	100

14 (61%) foreign bodies were located in the right main bronchus, while 9 (39%) were located in left main bronchus. No foreign body was located in one patient.

Discussion

There are two main types of bronchoscopes, the fibre optic and the rigid. The rigid can be further divided into ventilating and venturi type. The method of bronchoscope used to showcases in this study is the telescopic type, which is easily applied and makes anaesthesia and ventilation easy for the procedure, Swanson et al¹.

The use of telescope minimises episodes of desaturation and discontinuity of procedure, which is usually encountered during the use of rigid bronchoscope. The reason is that there is continuous air flow from the beginning to the end of the procedure. This ensure uninterrupted ventilation during the process, with sustained adequate saturation.

Male presented more than female, with male/female ratio of 2:1. No demonstrated reason for this, but could be due to increased activity in male children. This clinical condition appear commonest between ages 1-3 years and was also observed by Fidkowski et al, when they reviewed about 12,979 cases of anaesthetic consideration of tracheobronchial foreign bodies².

The use of telescopic/endotracheal intubation assembly has advantages over other methods like: Bronchoscopic Anaesthesia-Circuit Assembly, venturi-oxygen air-entraining rigid bronchoscope and flexible bronchoscope^{3,4}. In the telescope assembly there is a wider field of view and magnification of the airway objects, better control of the ventilation with virtually absence of desaturation of patient during the procedure. Minimal trauma of airway is also observed^{5,6}. Items removed from the airway include groundnuts (38%), followed by fish bone (21%), screw nuts (14%)and pieces of plastic toys (4%) as recorded on table III.

Ezeanolueet *al* had also recorded groundnut as commonest in their (30) thirty cases series that had rigid Bronchoscopy^{7,8}. The right main bronchus had more foreign body lodged in the lumen (61%). The left main bronchus had (39%). There was no foreign body in one patient⁹.We showcased this instrument to encourage other centres to acquire it,for easy access retrieval of foreign body in the tracheobronchial tree.

More children presented within age one to two (1-2 years) compared to other age group. This could be due to poor swallowing reflexes that had not developed properly in this age group.

Table I: This was also observed by Ezeanolue *et al* in their clinical profile and technique of administering anaesthesia during rigid bronchoscope removal¹⁰. Majority of patient presented after about five days. More complication are observed due to delay in removal of foreign body.

Groundnut has acidic secretions which will cause oedematous airway with more obstruction¹¹. Table III shows the type of foreign body aspirated, groundnut presented more, followed by fish bone.

Groundnut is a common vegetable food item provided at homes and with poor swallowing reflexes, aspiration is common¹². Fourteen (14) foreign bodies were retrieved from the right main bronchus. This agree with other studies. Anyanwu C.H foreign body airway obstruction in Nigerian children^{13,14}.

Conclusion

Foreign body in the airway especially in children is life threatening. The use of telescope offers excellent armament, preventing mortality and better option to other methods of bronchoscopy.

References

1. Swanson KL, Prakesh UBS, Midthun DE, et al. *Flexible bronchoscopic management of airway foreign bodies in children*. Chest 2002;121:1695-700.
2. Fidkowski CW, Zheng H, Firth PG. *The anaesthetic consideration of tracheobronchial foreign bodies in children: a literature review of 12,979 cases*. AnaethAnalg 2010;111:1016-25 [PubMed] [Google Scholar].
3. Ramirez-figueroa J.L, Gochicoa-Rangel L.G, Ramires-san Juan D.H, et al. *foreign body removal by flexible fibre optic bronchoscopy in infants and children*. PediatrPulmono 2005;40:392-7.10.10021 ppul.20242 [pubmed] [crossRef] [Google Scholar].
4. Farrell P.T rigid bronchoscopy for foreign body removal: anaesthesia and ventilation. PaediatrAnaeth 2004;14:84-9.10.1046/j.1460-9592.2003.01194-x. [pubmed] [crossRef] [Google Scholar].
5. Adesunkamni ARK, inhaled tracheobronchial foreign bodies: A Review. Niger J. Surg. Sci. 1994;4:34-38.
6. Benjamin B; Anaesthesia for paediatric airway endoscopy. Otolaryngology Clin North Am 2000;33:29-47.

7. Banjerjee a. Subba Rao RSVR, Khanna SK Narayaman RS, Gupta BE, Selan JC, Retanan CR, Naehiappam M, Larygo-tracheo-bronchial foreign body in children J. Laryngeal et al, 1988;102:1029-1032.
8. Adegboye VO. Osinowo O. Adebo OA. Soyawo O. Management of tracheobronchial foreign bodies. Nigerian Journal of Surgical Sciences 1993;3:1-4.
9. Chatterji S and ChatterjiP. The management of foreign bodied in air passages Anaesthesia 1972;27:390-395.
10. Ezeanolue B.C, Izuora K.L.A, Ezike H.A; *tracheobronchial foreign body in Nigerian children: clinical profile and a technique of administering anaesthesia during rigid bronchosopic removal*, journal of college of medicine June 2003; 8(1):27-29.
11. Hillel A.D. Sie .F; A reliable method to maintain an airway after bronchoscopy. Laryngoscope 1998: 98:1353-1355.
12. Mark S.C, Marsh B.R, Dungeon D.L; Indications for open surgical removal of airway foreign bodies; *Ann OtolRhinol Laryngeal*; 1993:102:690-691.
13. Anyanwu CH. Foreign body airway obstruction in Nigerian children J Trap Paed 1985:30:120-172.
14. Farrel PT. Rigid bronchoscopy for foreign body removal: anaesthesia and ventilation. PaediatrAnaesth 2004; 14:84-89.