

Minireview Article

Effects of Diabetes and Hypertension on Placenta

Abstract:

Placental insufficiency is the failure of the placenta to supply nutrients to the fetus and remove toxic wastes; it is a common cause of intra uterine growth restriction (IUGR), fetal distress peri and postnatal mortality. Specifically gestational diabetes and hypertension during pregnancy alters placental architecture which contributes to insufficiency of the placenta. Placenta is the most accurate record of infant's prenatal experiences. Clinical condition such as hypertension and diabetes that complicate pregnancy leads to significant structural changes in placenta, these histological changes and morphological alterations in placenta lead to insufficiency, resulting in IUGR and fetal distress. In the present study morphology of placentas collected from diabetic and hypertensive pregnancy will be observed, the histo-pathological findings will be recorded and correlated to birth weight, APGAR scores and congenital anomalies of babies and will be compared to the findings in normal placentas also. This study will be beneficial for obstetrician and pediatrician in gathering knowledge about patho genesis of placental insufficiency; it will add to the knowledge on placental morphology and will high light the abnormalities in placental anatomy

Keywords: Placental insufficiency, congenital anomalies, toxic wastes, Hypertension

Introduction:

Placental insufficiency is the failure of the placenta to supply nutrients to the fetus and remove toxic wastes; it is a complication of pregnancy where placenta which grows during pregnancy to feed the unborn child cannot bring enough nutrients. A normal placenta at term is reddish brown in color, rounded, flattened, and discoid about 450-500 grams in weight 15-20 centimeters in diameter and 2-4 centimeters thick. Umbilical cord can arise almost at any point and fetal membranes at margins. It is derived from fetal and maternal tissue both, the maternal portion is deciduas basalis and fetal portion is called chorion fundosum. (1).

Observation

Placental insufficiency is a common cause of intra uterine growth restriction (IUGR), fetal distress peri and postnatal mortality. The reason for this in some cases is unknown but in others it is strongly associated with diabetes, hypertension, viral infections and smoking, or if it is not properly implanted at accurate site. Specifically gestational diabetes and hypertension during pregnancy which alters placental architecture contributes to insufficiency of this structure which brings oxygen and nutrients to fetus and helps remove carbon dioxide and waste.(2)(3).This is a common problem encountered by Obstetrician and Pediatrician now a days .

Placenta is the most accurate record of infant's prenatal experiences. (4). Clinical condition such as hypertension and diabetes if complicate pregnancy leads to significant structural changes in placenta. (5). These histological changes and morphological alterations in placenta lead to insufficiency, resulting in IUGR and fetal distress. Normal physiological changes occur in spiral arteries in which endovascular plugs penetrate their lumen and normally disappear at the end of 2nd trimester. (6). In some utero-placental vessel these physiological changes do not take place because of hypertensive complication leading to acute atherosclerosis. (7).

More recent studies have shown relation in hypertension and placental lesion and established relation between lesion and clinical severity but still microscopic features are to be reviewed in depth. (8)(9).

The term placenta was used for the first time in 1559, it is a mirror that reflects the intra uterine status of fetus, fetal distress, intrauterine fetal death and placental abnormalities are common in pregnancy induced hypertension and it is noticed that placental surface area is less in cases related to fetal distress. (2). Physiological changes in spiral arteries occurs in normal pregnancy but are absent in pre eclamptics. (10). Placental ischemic alterations are resulting from decrease in maternal blood supply to the placenta associate with hypertensive syndrome of pregnancy. (11).

Placenta produce hormones in addition to other function and these have an opposite effect on insulin which causes hyperglycemia in mothers ultimately contribute to fetal growth, this result in gestational diabetes which disappear often after delivery, but if mothers have type I diabetes mellitus chances of congenital birth defect increases. (12). In recent study done in 2008 by Beverley et.al. They correlated the placental weight which decreases in response to mutation in glucokinase gene of fetus. (13). Glucose, amino acid and other substrates cross placenta via different mechanisms (14), recently a significant link has been observed between measurement of umbilical cord insulin and placental weight. (15). Placental weight increases in diabetic pregnancies a condition known to be associated with fetal hyper insulinemia although this could be caused by the maternal hyperglycemic environment rather than fetal insulin levels. (16).

Scott in 2008 demonstrated that both fetal and placental weight increases in presence of maternal diabetes, their respective weight remains high and fetal weight-to placental weight ratio index (FPI) is not significantly reduced [20,21] . Lower FPI is driven primarily by higher placental weight in both control subjects and off springs of mothers with type I diabetes and is also associated with poorer maternal glycemic control in offspring of mothers with type 1 diabetes. (3).

Adiponectin concentration in umbilical cord serum is positively associated to the fetus and placental weight ratio. (17). This association reflects augmentation of insulin sensitivity and consequently increased fetal growth direct effect of fetal adiponectin on placenta are not clear which suggests that other mechanisms are also involved. (18).

Conclusion

High blood pressure (hypertension) during pregnancy can prevent the placenta (the food supply for the baby in the womb) from receiving enough blood. The lowered amount of blood to the placenta can lead to a low birth weight. Other complications can occur from high blood pressure during pregnancy Diabetes-induced placental malformation decreases placental efficiency and fetal size. Hyperglycemia disrupts rat trophoblast stem cell differentiation into invasive trophoblast. Diabetes impairs intrauterine trophoblast invasion. Hyperglycemia adversely affects placental adaptations to hypoxia.

References:

1. Baschat A A, Galan H L, Ross M G. et.al. Intrauterine growth restriction. *Obstetric-Normal and problem pregnancies .5th Ed.* New York, NY: Churchill Livingstone; 2007: chap. 29.
2. Udiana A, Bhagwat S S, Mehta C D .Relation between placental surface area, infarction and fetal distress in pregnancy induced hypertension with its clinical relevance. *J Anat Soc India* 2004; **53**(1)27-30.
3. Scott M. Nelson, Dilys J. Naveed Sattar. et.al Role of adiponectin in matching of fetal and placental weight in mothers with type I diabetes. *Diabetes Care* 2008; **31**(6).
4. Benirschke K. *The Placenta: How to examine it and what you can learn .Contemp Obst and Gynaecol* 1981; **17**:117-9.
5. Majumdar S, Dasgupta H, Bhattacharya K et.al. A study of placenta in normal & hypertensive pregnancies. *J Anat Soc India* 2005; **54**(2):1-9
6. Benirschke K, Kaufmann P. *Pathology of the human placenta.* New York: Spriger Verlag 1990; 2nd Ed.
7. Stallmach T, Hebisch G. Placental pathology: It's impact on explaining prenatal and perinatal death. *Virchows Arch* 2004; **445**:9-16.
8. Sebire N J, Goldin R D, Regan L. et.al. Term preeclampsia is associated with minimal histopathological placental features regardless of clinical severity. *J Obstet Gynecol* 2005; **25** (2):117-8.
9. Zetterstrom K, Lindeberg S N, Haglund B. et.al. Maternal complication in women with chronic hypertension: A population based cohort study. *Acta Obstet Gynecol Scand* 2005; **84**:419-24.
10. Gerretsen G, Huisjes H J, Elema J D. et.al. Morphological changes in spiral arteries the placental bed in relation to pre-eclampsia and fetal growth retardation *International journal of Obstetric and Gynecology* 2005; **88**(9): 876-81.
11. Rosana R. M. Correa, Daniel B.Gilio, Camila L. et.al. Placental morphometrical and histopathological changes in different clinical presentations of hypertensive syndromes in pregnancy. *Arch Gyneacol Obstet* 2008; **277**:201-6.
12. *Gestational diabetes .Diabetes* 2004
13. Beverley M. Shields, Gill Spyer, Annabelle S. Bridget A. Knight et.al. .Mutations in the glucokinase gene of the fetus results in reduced placental weight. *Diabetes Care* 2008; **31**(4).
14. Jansson T: Amino acid transporters in human placenta. *Pediatr Res* 2001; **49**:141-7.
15. Scott C, Dunger D: Size at birth and cord blood levels of insulin, insulin like growth factor I (IGF-I), IGF-II, IGF-binding protein –I (IGFBP-1), IGFBP-3, and the soluble IGF-II/mannose-6-phosphatereceptor in term human infants: the ALSPAC study team: Avon Longitudinal Study of Pregnancy and Childhood. *J Clin Endocrinol Metab* 2000; **85**:4266-4269.
16. Taricco E, Radaelli T, Nobile de Santis M S, Cetin. Feotal and placental weights in relation to diabetes. *Placenta* 2003; **24**:343-7.
17. Kadowaki K, Nakanishi I, Miyashita Y et.al. Adiponectin concentration in umbilical cord serum positively associated with the weight ratio of fetus to placenta. *J Clin Endocrinol Metab* 2006; **91**: 5090-4.

18. Chen j, Tan B, Karteris E, zervou S, et.al. Secretion of adiponectin by human placenta: differential modulation of adiponectin and its receptors by cytokines. *Diabetologia* 2006; **49**: 1292-1302.
19. Marina kos, Bernard Czernobilsky, Ljiljana Hlupic, Kristian Kunjko. *Croat Med J.* 2005; **46**(3):404-9.
20. Ehlers E, Talton OO, Schust DJ, Schulz LC. Placental structural abnormalities in gestational diabetes and when they develop: A scoping review. *Placenta.* 2021 Dec 1;116:58-66.
21. Sato N, Fudono A, Imai C, Takimoto H, Tarui I, Aoyama T, Yago S, Okamitsu M, Mizutani S, Miyasaka N. Placenta mediates the effect of maternal hypertension polygenic score on offspring birth weight: a study of birth cohort with fetal growth velocity data. *BMC medicine.* 2021 Dec;19(1):1-3.

UNDER PEER REVIEW