

Original Research Article

Synthesis, Characterization And Antimicrobial Screening - In Vitro And In Silico Of A Novel Hydrazone Compound Exhibiting Strong Bacterial *Escherichia Coli* Inhibitory Potentials.

ABSTRACT

The geometric increase of drug-resistant bacteria pathogens has made urgent the research, development and production of new antibacterial and antifungal compounds. As hydrazones enhanced generally with thiazoles which are 5-membered ring compound containing active nitrogen and sulphur molecules (C₃H₃NS) have been proven to exhibit strong antibacterial and antifungal activities, in this study, salicylaldehyde -4- thiazoleacetic acid hydrazone(SAFTAH) compound was synthesized. The novel compound was characterized and subjected to anti-microbial screening. The microbes employed were *staphylococcus aureus*, *Escherichia Coli*, *streptococcus* and *klebsiella aero genes*. The compound was found to be active against *Escherichia coli*. To validate this inhibition of *Escherichia coli* by the novel compound and to detect the active site of their interactions, in silico molecular docking analysis of the novel compound against aminopeptidase N from *E. coli* which is known to promote virulence to the microbe in question was carried out. Six drugs commonly used for the treatment of *E. coli* viz-a-viz, ciprofloxacin, levofloxacin, doxycycline, trimethoprim, rifamycin, rifaximin and sulfamethoxazole were subjected to same type of silico studies. The result of the docking indicated that the novel hydrazone compound was more efficient and effective than five of these *E. coli* inhibitory drugs. The success recorded from in vitro and in silico analyses stems from the efficacy of the novel compound not clear. Salicylaldehyde -4- thiazoleacetic acid hydrazone could therefore be a potential drug for the cure of *E. coli* infection.

Keywords

Escherichia coli, Antimicrobial screening, Molecular docking, Binding affinity, Salicylaldehyde -4- thiazoleacetic acid hydrazone

INTRODUCTION

Generally, acyl hydrazone are compounds that contain the -CNN- group, synthesized generally by eliminating water molecule between hydrazine and any of aldehyde or ketone (carbonyl compound) It could be by the condensation of hydrazide with carbonyl compounds [K. K. Bedia et al; 2006] Hydrazides are the acylated derivatives of hydrazine [J. Pisk, I. Dilovic et al; 2020].

A great attention is given to hydrazones because of their biological and physiological activities [Mohamma Asif and Asif Husan 2013]. They generally exhibit very strong antibacterial and antifungal activities.

A good number of researchers are interested in synthesizing variety of acyl hydrazones which are known to associate with antimalarial, antitumour, analgesic, antibacterial, antifungal, antitubercular antihelminthic, anticonvulsant, antimicrobial, antidiabetic and/or anti-inflammatory activities [M. E. Shakhofa et al, 2017, M. Asif and A. Husain; 2013]. Examples of such hydrazone include; N- (4-tert-Butylbenzoyl)-2-hydraxynthalaldehyde (BBNH) a potent inhibitor of the ribonuclease H (R Nase H) activity of human immunodeficiency virus (HIV)-1 reverse transcriptase (RT). This BBNH binds to the HIV- 1 RT R Nase H active site via coordination to the metal and to amino acid. HIV- 1 RT R Nase H can reasonably be inhibited by N- acylhydrazone and phenylhydrazones, [Y. Jin, Z. Tan et al; 2010, K. Nagano, H. Kinoshita and A. Hirakawa; 1964, R.C. Aggarwal, N.K Singh] A Research programme carried out on a series of these acyl hydrazones discovered that they are potentially tridentate ligands.

In addition, hydrazones of thiazoleacetic acid? are so enhanced that they are used as monoamine oxidase inhibitors and could be applied for curing psychotic illnesses [J.N. Nwabueze ; 1997]. [These thiazolehydrazones are of the following structure

Comment [GK1]: Can the topic be written like this? Synthesis, Characterization, In Vitro and In Silico Antimicrobial Screening of a Novel Hydrazone Compound

Comment [GK2]: Put in italics

Comment [GK3]: Put in italics

Comment [GK4]: Remove

Comment [GK5]: Change to properties

Comment [GK6]: What is this?

Comment [GK7]: Put in italics

Comment [GK8]: the

Comment [GK9]: Check all spacing and typo errors

Comment [GK10]: Put in vitro and in silico in italics

Comment [GK11]: Check sentence and remove not clear.

Comment [GK12]: Put in italics

Comment [GK13]: The introduction is too long. Edit it and leave only important details

Comment [GK14]: Check all references in the body of the text and at the reference section. Comply with Author guidelines.

Comment [GK15]: ?

Comment [GK16]: ?

Comment [GK17]: Check spacing and writing of references

Comment [GK18]: Rewrite sentence. Do not put question in the sentence and do not bold.

Comment [GK19]: The structure of the thiazolehydrazone is shown below




Fig. 1. Chemical structure of acetone[4-(5-nitro-2-furyl)-2-thiazolyl]hydrazone

Another compound known as acetone[4-(5-nitro-2-furyl)-2-thiazolyl]hydrazone is found to have anti cancer potentials[. S. Emily; 2011]

Escherichia coli (*E. coli*) belongs to a family of microbes known as Enterobacteriaceae. These are gram-negative anaerobic rod. Generally, *E. coli* are about 2.6 μm long and 1.1 -1.5 μm wide which occur as single and straight rods. This bacteria is mostly responsible for acute urinary tract infection and tract sepsia in human. It causes acute enteritis together with haemorrhagic colitis. Its presence can lead to neonatal meningitis. As this infections grows as a biofilm, it is often present in the environment.[C.A Batt(2014) in Encyclopedia of food Microbiology 2nd edition].

Amino peptidase N from *Escherichia coli* is a major metalloprotease which can be located in the small intestine and it participates in the controlled hydrolysis of peptides. That is hydrolysis of proteins, [https://www.ncbi.nlm.nih.gov] It is a M1 class amino peptidase with active region similar to the active region of thermolysin. They are zinc dependent enzymes. They are secreted by, precisely from the acinar cells of the pancreas. This enzyme is one of the known and attractive targets of *e. coli* associated disease (Sonia George et al 2014)

The *in vivo* microbial screening and *in silico* analysis of the novel compound; salicylaldehyde -4-thiazoleacetic acid hydrazone was performed in this study The docking was on Amino peptidase N from *Escherichia coli*. The Salicylaldehyde -4- thiazoleacetic acid hydrazone was tested for sensitivity using *Staphylococcus aureus*, *Escherichia coli*, *Streptococcus* and *Klebsiella aero genes* microbes, the novel compound was docked on amino peptidase N enzyme from *e. coli*. The binding energy and mechanism of action was compared with those of seven known drugs for the treatment of *e. coli* bacteria in Nigeria.

Materials and Methods

Ethyl-2-amino-4-thiazoleacetate was obtained from Sigma – Aldrich Chemical Company Ltd and used without further purification, while other reagents which include the carbonyl compound-salicylaldehyde together with the solvents used which were ethanol and methanol were from the BDH Chemical Ltd, Pools England. Known drugs for treatment of *E. coli* in Nigeria which include Ciprofloxacin, levofloxacin, doxycycline, trimethoprim, rifamycin, rifaximin and sulfamethoxazole were bought from Orchard Pharmacy Owerri and Pax Pharmacy Onitsha in Nigeria.

The Infrared spectra of the synthesized compound in Nujol were taken on FTIR-8400S Fourier Transformation Infrared Spectrophotometer and the proton NMR using dimethyl sulfoxide DMSO-d₆ and recorded on a Bruker Avance 400 NMR spectrometer were both carried out at the National Research Institute for Chemical Technology (NARICT), Zaria, Nigeria.

The antimicrobial screening of the novel compound was carried out at University of Abuja Teaching Hospital and Peak Medical Laboratory, Gwagwalada using nutrient Agar on bacteria *staphylococcus aureus*, *Escherichia Coli*, *Streptococcus* and *Klebsiella aerogenes* obtained from the Teaching hospital.

Preparation of 2-amino-4-thiazoleacetic acid hydrazide) (ATAH)

Standard method was used to prepare the hydrazides [120]?. 1mole of ethyl-2-amino-4-thiazoleacetate was reacted with 1 mole of the hydrazine hydrate to give the required hydrazide.

Comment [GK20]: What is the relationship of this compound with the synthesized compound? This must be clearly stated

Comment [GK21]: Remove bold

Comment [GK22]: Please check all references

Comment [GK23]: Spacing

Comment [GK24]: This was copied from internet. This is why it is shown as blue colour. Type it out and rephrase

Comment [GK25]: Check author guideline for reference writing

Comment [GK26]: What is MI?

Comment [GK27]: *E.coli*

Comment [GK28]: disease

Comment [GK29]: Check reference

Comment [GK30]: ?

Comment [GK31]: one word: aerogenes

Comment [GK32]: *E.coli*

Comment [GK33]: *E.coli*

Comment [GK34]: Spacing. Standard drugs.....

Comment [GK35]: Remove

Comment [GK36]: What is this?

Comment [GK37]: required

2.50ml (2.40g; 0.04moles) of hydrazine hydrate was added to 7.0g (0.040 moles) of ethyl-2-amino-4-thiazoleacetate in 50ml of absolute ethanol. Antibumping granules were added and the mixture was refluxed, on a water bath for six hours in a 100ml round-bottom flask. The mixture was poured into a beaker, left for three days to crystallize, and the crystals formed were filtered and recrystallized from ethanol. The resulting crystals were filtered and dried over silica gel in a vacuum desiccators and weighed (Yield, 4.61g; 65.86%). The crystals obtained were light brown in colour.

Comment [GK38]: Put space after figure and unit

Comment [GK39]: ?

Comment [GK40]: Give the structure of the intermediate compound

Preparation of Salicylaldehyde-2-amino-4-thiazoleacetic acid hydrazone (SAFTAH)

0.86g, (0.005 mole) of 2-amino-4-thiazoleacetic acid hydrazide (ATAH) was mixed with 0.53ml (0.601g, (0.005 moles) of salicylaldehyde in 50ml ethanol and refluxed for 4 hours in a 150ml round bottom flask on water bath. The solution was left for one day to crystallize. The crystals obtained were filtered and were recrystallized from ethanol. The yellow crystals were then dried in a desiccator over silica gel. (Yield, 0.97g; 80.78%)

Comment [GK41]: Check all figures and units

Comment [GK42]: Give the structure of the final product

Comment [GK43]: Remove the full stop

The structure and characterization of the synthesized compound was elucidated using the FTIR infrared spectra using Nujol disc and proton NMR spectrometer and some physical analyses.

Comment [GK44]: Capital I

In vitro susceptibility testing of the synthesized compound was carried out on four types of microorganisms viz *Staphylococcus aureus*, *Escherichia coli*, *Streptococcus*, *Klebsiella aerogenes*.

Comment [GK45]: Characterization and structure determination of the synthesized compound was done using the FTIR obtained using Nujol disc and proton NMR. Other physical analyses were also carried out to confirm the structure.

Comment [GK46]: In italics

Comment [GK47]: *Staphylococcus*

Comment [GK48]: *Staphylococcus aureus*

Comment [GK49]: ?

Comment [GK50]: Spacing problem and remove full stop before reference

Docking procedure

The energy minimized compounds, (synthesized and the known existing drugs) were subjected to docking analyses on aminopeptidase N from *E. coli* so as to predict their various interactions with the main binding sites on this enzyme. The Autodock Vina in Pyrx virtual screening software 20 (version 0.8) was employed for the docking analysis. [Duru et al; 2020] The grid box sizes were x centre: 19.19, y centre: 17.59, z centre: 20.88.

Biovia Discovery studio (Biovia 2020) was used to visualize the molecule-ligand interaction (enzyme-drugs) at the end of the docking process and to understudy the simulations

The binding affinities of the different compounds (drugs) docked on the protein target were obtained and the result organized on an excel spreadsheet [Chioma Nnenna Nwofor et al: 2022]

Comment [GK51]: ?

Results and Discussion

The physical analysis carried out has the following result

Comment [GK52]: Physical analysis; solubility test, conductivity, melting point, colour determination are reported in Table 1. Remove bold

Table 1. Results of physical analysis

Compound	Formular	FM WT	Colour	MPT/DEC	%Yield	Conductivity $\Omega^{-1} \text{ cm}^3 \text{ mol}^{-1}$
SAFTAH	$\text{C}_{12}\text{H}_{12}\text{N}_4\text{O}_5$	171	Yellow	210°	81	-0.289

Solubility Test

Compounds	Water	Acetone	Ethanol	Methanol
SAFTAH	IS	S	S	S

Key

IS Insoluble

S Soluble

Infrared spectra.

The infrared spectra is seen in Table 4.3. The band of interest include the $\nu(\text{OH})$, $\nu(\text{NH})$, $\nu(\text{C}=\text{N})$, $\nu(\text{C}=\text{O})$, $\nu(\text{C}=\text{N})$,

Comment [GK53]: The frequency of absorption obtained from the Infra red spectra is shown in Table 2

Compound	$\nu(\text{OH})$	$\nu(\text{N-H})$	$\nu(\text{C}=\text{O})$	$\Delta \nu(\text{C}=\text{O})$	$\nu(\text{C}=\text{N})$	$\Delta \nu(\text{C}=\text{N})$	$\nu(\text{SO}_4^{2-})$	$\nu(\text{CH}_3\text{COO})_2$	$\nu(\text{M-O})$	$\nu(\text{M-N})$
SAFTAH	3788		1750		1589					423

Comment [GK54]: Remove this if there is no peak or absorption for this functional group

H-NMR Spectrum of the novel compound - SAFTAH

Table 2. Chemical Shift and multiplicity

s/n	Chemical Shift	Multiplicity
5	11.15	Singlet
6	3.76	Singlet
8	10.06	Singlet
10	8.38	Singlet
12	8.23	Singlet
13	7.61	Multiple
14	7.26	Multiple
15	7.50	Multiple
16	6.89	Multiple
17	3.39	Singlet

Table 3. Antimicrobial test result for the compound.

Compound	<i>S.aureus</i>	<i>E.Coli-</i>	<i>S.Spp</i>	<i>K.genes</i>
SAFTAH	-	++	+	-

KEY

S.aureas= Staphylococcus aureas
E.Coli= Escherichia coli
S.spp= Streptococcus species
K.aerogenes= Klebsialla aero gene

Comment [GK55]: Put space between the alphabet and the specie and the specie should start with small letter

Comment [GK56]: All the microorganism names should be in italics

The *Ecoli* Enzyme and the various Interractions with its Key Active Regions

Comment [GK57]: ?




Fig. 2. Aminopeptidase N enzyme from *E coli*

Comment [GK58]: ?




Fig. 3. The 2D interaction of SAFTAH with the active sites of the enzyme from *E. coli*

Comment [GK59]: ?




Fig. 4 The 2D interaction of Ciprofloxacin drug with the enzyme from *E. coli*

Comment [GK60]: ?




Fig. 5. The 2D interaction of Doxycycline drug with the enzyme from E coli




Fig. 6. The 2D interaction of Rifamycin drug with the enzyme from E coli

Comment [GK61]: ?




Fig. 7. The 2D interaction of Rifaximin drug with the enzyme from E coli




Fig.8. The 2D interaction of Sulfamethoxazole drug with the enzyme from E coli




Fig. 9. The 2D interaction of Cocrystallized ligand with the enzyme from E coli

The compound obtained was active against *E. coli* used for the *in vitro* screening and *in silico*, the interaction of the novel compound with the aminopeptidase N enzyme from *E coli* is very positive. Therefore, the activity profile of the novel compound indicates the existence of a very significant correlation between the *in vitro* screening and the computational data from the *in silico* studies. Moreover, in its interaction with the *E coli* enzyme, It competes and surpasses five (5) of the commonest known drugs for the treatment of patients suffering from *E coli* in Nigeria..This is so because the novel compound binds better than these drugs at the active sites of the protein, The binding energy of the novel compound is appreciably lower than those of the known drugs as follows; Binding energy of the novel compound -6.6, while ciproxofloxacin drug is -6.0, doxycycline drug is -6.2, rifamycin drug is -5.2, rifaximin drug is -5.7 and sulfamethoxazole drug is -6.6. It follows then that the thiazole compound which has been proven to exhibit strong antibacterial and antifungal activities was highly enhanced and strengthened by the salicylaldehyde compound for the **2-amino-4-thiazoleacetic acid hydrazide** alone was subjected to same screening - *in vitro* and *in silico* and the result obtained fell below. Salicylaldehyde on its own does not have a history of drug potentiality when used alone but does exhibit the ability of adequate enhancement for compounds with drug potentials. Salicylaldehyde certainly complemented the known efficacy of **2-amino-4-thiazoleacetic acid hydrazide** making this novel compound a potential *E coli* drug. Reacting this novel hydrazide compound with some other carbonyl compounds and subjecting salicylaldehyde into reactions with different hydrazides, all for comparative studies would make a good research and offer proof of whether or not the salicylaldehyde is more efficacious than other carbonyl compounds and whether the novel hydrazide compound possesses more *e coli* treatment ability than other hydrazides.

Conclusion

The success recorded - both *in vitro* and *in silico* analyses of the novel compound synthesized, certainly stems from its efficacy. Therefore, **Salicylaldehyde-2-amino-4-thiazoleacetic acid**

Comment [GK62]: bacteria or enzyme?

Comment [GK63]: *E.coli*

Comment [GK64]: Remove bold

Comment [GK65]: ?

Comment [GK66]: Remove bold

Comment [GK67]: Put in italics

Comment [GK68]: ?

Comment [GK69]: Put as RECOMMENDATION under conclusion

Comment [GK70]: ?

Comment [GK71]: Be more specific. Efficacy in terms of what?

hydrazone (SAFTAH) emerges as an excellent potential and effective drug for the cure of *Escherichia coli* infection in Nigeria. Synthesis of the compound is not difficult and so can be very handy for the treatment of the *E coli* infection which appears to be incessant in Nigeria. The cure promises to be more stable than what obtains currently considering the low binding energy of interaction.

Comment [GK72]: Remove bold

Comment [GK73]: Italicize

Comment [GK74]: ?

Comment [GK75]: You used double full stops. Why?

REFERENCES

1. K. K. Bedia et al (2006) Synthesis and characterization of novel hydrazone-hydrazones and the study of their structure – antituberculosis activity. *European Journal of Medicinal Chemistry* 41 (11) pp 1253-1245
2. J. Pisk, I. Dilovic et al (2020) Effective methods for the synthesis of hydrazones, quinazolines and Schiff bases: Reaction monitoring using a chemometric approach. *Royal Society of Chemistry Advances*. 10. 38566-38567.
3. M. E. Shakhofa et al (2017) Homotrimetallic transition metal complexes of 2-hydroxy-N¹, N², N³-triphenylpropane-1,2,3-tricarbonylhydrazide preparation and microbe activities. *Research Journal of Pharmaceutical, Biological and Chemical Sciences*. 8 (2) 724
4. M. Asif and A. Husain (2013) Analgesic, anti-inflammatory and antiplatelet profile of Hydrazones containing synthetic molecules. *Journal of Applied Chemistry, Hindawi, India*.
5. Y. Jin, Z. Tan et al (2010) SAR and Molecular mechanism study of novel acylhydrazone compounds targeting HIV-1 C A. *Journal of Bioorganic and Medicinal Chemistry*. Vol 8 (6) pp 2135-2136
6. R.C. Aggarwal, N.K Singh and Prasad (Unpublished work)
7. J.N. Nwabueze (1997) Complexes of Copper (II) with Acetone Hydrazones derived from some cyclocarboxylic acids. *Synth. React. Inorg. met.-Org. Chem.* 27, 673
8. S. Emily (2011). Hospital Epidemiology and Infection Control in acute-care setting. *Clinical Microbiology Reviews*. 24 (1): 141–173.
9. C.A Batt(2014) in *Encyclopedia of food Microbiology* 2nd edition.
10. <https://www.ncbi.nlm.nih.gov>
11. T. Funke, (2004,2007,2010) Case. *Microbiology, An Introduction* 10th Edition. Pearson Benjamin Cummings..
12. Sonia George et al (2014) Design, docking, synthesis and anti E coli screening of novel thiazole thiourea derivatives as possible inhibitors of enoyl reductase (fabI) enzyme. *Bangladesh J. Pharmacol. India*. 9: 49-53.
13. C.E Duru, I.A Duru and A. Bilar (2020) Computational investigation of sugar fermentation inhibition by berginin at the pyruvate decarboxylase isoenzyme 1 target of *Scharyomyces cervisiae*. *J. Med. Plants Stud* 8(6) 21-25
14. C. N. Nwofor, C. E Duru and N. E Onyenwe (2022) Computational identification of small molecules in *Mitracarpus scaber* ethanolic leaf extract with fungal keratinase inhibitory potentials. *Journal of Natural Pesticide Research, Elsevier*. Vol 2 p.

Comment [GK76]: Follow author guidelines in writing the references.

UNDER PEER REVIEW

