

***In silico* screening of potential compounds from Medicinal plants by targeting
Streptococcus mutans deoxycytidylate deaminase**

Abstract

Dental caries is the chronic infectious disease caused by bacteria to form a biofilm on the tooth surface of man. Drug-resistant *Streptococcus mutans* (*S. mutans*) poses a vital public health issue. To overcome this, the development of effective drugs with novel mechanism of action is requisite. Drug repurposing is considered a viable alternative approach to overcome the above issue. In the present study, we have attempted to select unique and traditional sources used in traditional medicine. Traditionally, many cultures use chewing sticks for oral hygiene maintenance. When properly used, these chewing sticks are found to be efficient due to the combined effect of mechanical cleaning, enhanced salivation and the antimicrobial action of leached plant. Inhibition of the *S. mutans* deoxycytidylate deaminases (SmdCDs) is the most promising drug development strategy against the *S. mutans*, responsible for the biofilm formation. In the present work, out of 871 phytochemicals 211 phytochemicals were shown to be the most druggable substance with zero violation from any of the druglikeness rules. Further, the binding energy indicates the affinity of the protein structure docked with the 2 hit potential herbal compounds of which Cyclocurcumin and Androsta-1,4,6-triene-3,17-dione showed the best docking with the SmdCD.




Figure 1. outline of the study

Introduction

The oral hygiene is influenced by numerous factors, mainly diet and host immune competence, promoting the virulence and adhesion of microorganisms (Gao et al., 2018; Reibel, 2003) About 700 microbial species are identified from oral microbiome(Jr et al., 2008). Dental caries is one of the most common chronic infectious and diseases that occur at any age of humans (Gao et al., 2018). Generally, dental plaques occur when the oral microbial is low pH, thereby creating the presence of increased acid producing and acid tolerant bacteria in a structurally and functionally organized biofilm formation (Marsh, 2010). This has been attributed to the consumption of dietary free sugars (sucrose), either as additives or preservatives, and they are linked to biofilm induced tooth decay (Abdel-Aziz et al., 2020; Moynihan, 2016).The main causative agent is gram-positive bacteria *Streptococcus mutans* (*S. mutans*) associated with dental caries and dental plaque formation(Bowen & Koo, 2011).Unlike microbial biofilms display increased tolerance to the host defenses and antimicrobial agents (Algburi et al., 2017; Marsh, 2010), challenging the clinical management of dental plaque. The commonly used method for caries prevention was mechanical plaque control, such as tooth brushing and flossing. Excessive use of antimicrobials is considered as an appropriate combinatory measure for the control of dental caries, particularly

in the high risk population (Rath & Singh, 2013). Microbes within the several different approaches have been developed to prevent dental caries such as Chlorhexidine (CHX) is one of the most common antimicrobial agents. It is recognized as the principal agent for chemical plaque control (Jones, 1997). However, CHX has cytotoxic effects on a wide variety of human cells including oral mucosal cells, blood cells, keratinocytes, osteoblasts, and osteoclasts (Cabral & Fernandes, 2007; Ribeiro et al., 2004). Besides, CHX can cause taste confusions, tooth staining, and drug resistance. Thereby, alternative antibacterial agents are still needed for the control of dental plaque and caries.

The search for new drug candidates against tooth disease has led to the discovery of molecular targets, the development of drug, and explores of new bioactive substances. Despite this, there is no effective treatment available in the market. Hence in this study, we used small molecule phytochemicals against Putative deoxycytidylate deaminase(dCD) alloenzyme of *S. mutans* because several efforts have been devoted to characterizing the mechanisms of action of these phytochemicals. Numerous studies have been done on the antibacterial effects of natural medicinal plants. Several reports suggest that phytochemicals possesses remarkable inhibitory activities against bacteria.

In gram-positive bacteria and eukaryotic organisms, deoxycytidine-5'-monophosphate (dCMP) deaminases, dCDs catalyze the conversion of dCMP to deoxyuridine monophosphate (dUMP) in the pyrimidine salvage pathway. A crystal structure of dCD complexed with deoxycytidine triphosphate (dCTP) and a substrate analogue from *Streptococcus mutans* (SmdCD–dCTP) indicates an activation mechanism triggered by dCTP (Li et al., 2016). dCTP can allosterically bind to SmdCD and induce a conformational change to activate deamination. The deoxythymidine triphosphate (dTTP) bound complex adopts an inactive conformation that is consistent with its inhibitory role. To clarify the significance of the variability of the regulatory mechanism, it is necessary to compare a pair of activator-bound and inhibitor-bound SmdCD structures from the same species. As dTTP increases, dCTP can be replaced by dTTP from SmdCD and the deamination activity decreases, and vice versa. Furthermore, SmdCD reduces the efficiency of anticancer and antimicrobial drugs (Hernandez-Santiago et al., 2002)(Vellappally et al., 2017), which indicates that SmdCD inhibitors have a potential application for drug discovery.

Currently, herbal medicines have received greater attention because of their multiplicity of curing diseases, safety and being well tolerated remedies when compared with the conventional drugs. Plants are known to produce a variety of phytochemicals to protect themselves against a variety of pathogens. The use of phytochemicals present in the medicinal plants plays a crucial role in destroying the cross-links of the biofilm matrix. If we focus on the search for new drug candidates against periodontal diseases has led to the discovery of molecular targets and explores of new bioactive inhibitors. Hence in this study, we used selected medicinal plants phytochemicals derivatives against SmdCD. In the present study, we aimed to test the potency of 871 derivatives in inhibition of the SmdCD target through *in silico* methods. Therefore, drug-like properties was also predicted to ascertain the pharmacokinetics and drug ability of selected phytochemicals. The ligand based virtual screening of the library is carried out with target substrate were performed with the hit compounds.

Material and Methods

Construction of phytochemical library

Text mining analysis of plants by using Carrot2 and PubTator server which showed that selected plants photochemical had potential antimicrobial properties. Hence to find out an antibacterial activity against SmdCD enzyme, a library of 871 phytochemicals was constructed from 13 plants through searching the scientific literature and PubChem database. Further, 3D structure of each phytochemicals retrieved from PubChem (<https://pubchem.ncbi.nlm.nih.gov>) in SDF format and further converted all of them into PDB format by using Open Babel tool(O'Boyle et al., 2011).

Enzyme preparation

The 3D crystal structure of SmdCD target with PDB ID 5C2O was retrieved from the Protein Data Bank (<https://www.rcsb.org>). Here, the all water molecules, charge ions, and extra ligands were removed from the enzyme using PyMOL software(Lua & Lichtarge, 2010). After that the adding of hydrogen atoms to the enzyme was carried out by using MG Tools of AutoDock Vina software (Trott & Olson, 2009). The crystal structure of allosteric protein was then saved in PDB format for further analysis.

Ligand preparation

The 3D structure of each phytochemical was retrieved from PubChem (<https://pubchem.ncbi.nlm.nih.gov>) in SDF format and then converted into PDB files using Open Babel open source software. The crystal structure of SmdCD complexed with TTP (Compound CID: 64968) is presented at 2.35 Å resolution. The structure of reference molecule was retrieved from Protein Data Bank (<https://www.rcsb.org>).

Drug-likeness prediction

Analysis of molecular properties and drug-likeness of the screened phytochemicals is an important step in drug discovery. To be effective as a drug, a potent phytochemicals must reach its target in the body in sufficient concentration, and stay there in a bioactive form long enough for the expected biologic events to occur. Here, we used the new Swiss ADME web tool (Daina et al., 2017) that gives free access to a pool of fast yet robust predictive models for physiological property pharmacokinetics, drug-likeness and physicochemical properties. Easy efficient input and interpretation are ensured through the login-free website <http://www.swissadme.ch>. (Egan et al., 2000). Therefore, all screened phytochemicals were evaluated for their drug-like nature under different rules: Lipinski's rules of five; 'RO5 (Lipinski, 2000), Ghose filter, PAINS filter and Verber filter. The drug-likeness property of the hit molecules was checked by SwissADME web tool.

Molecular docking

Molecular Docking is the computational technique use a drug discovery. Molecular Docking is the method which predict of prefer orientation of the phytochemicals to bind at the active site of the receptor domain to form a stable complex by using AutoDock Vina software in PyRx open-source software (GUI version 0.8 of AutoDock) (Trott & Olson, 2009). Firstly docking was performed using SmdCD, reference molecule to validate docking protocol. Pdbqt format of the receptor and phytochemicals were dragged into their respective columns in form of pdbqt formate. The grid box center for docking set as X, Y, Z and with the dimensions of the grid box for SmdCD target. Now, we can run docking with Vina. Docking was performed to obtain a population of possible orientations and conformations for the phytochemical at the binding site. The finally, the binding energy table was extracted from the software. Once the analysis is completed we can check the result of best different confirmation with the lowest binding energy pose or docking score than that of the positive control was chosen after the docking search was

completed. The binding affinities of phytochemicals for SmdCD target were recorded. At the end of the docking, the best conformations were compared with the rigid docking for binding energy (kcal/mol). The best conformation of the phytochemicals which had lower binding energy as compare to reference molecule were chosen for further toxicity prediction analysis.




Figure 2. Binding pocket of SmdCD showing experimental and docked reference ligand(Blue color) binds to modulator site residues. Modulator site residues are in orange circles representation. Hydrogen bonds and that formed between protein and ligand are shown by green dotted lines and other residues are hydrophobic bond-forming residues.

Toxicity prediction

Screened phytochemicals obtained from molecular docking were preceded for toxicity prediction. The toxicity analysis of phytochemicals which have a good binding affinity with protein was carried by using OSIRIS property explorer program(Sander et al., 2009). The main aspect of US Food and drug administration toxicity risk predictor tool OSIRIS evaluated various toxicity risks properties phytochemicals such as tumorigenicity, mutagenicity, irritation, and reproductive development toxicity.

Visualization

Now, we were able to see our phytochemicals have been docked in the substrate active site. The result can be visualized and analysed different bioinformatics tool LigPlot + v.2.2.5 software. The 2 D depiction of hydrogen-bond interactions of the complex receptor-ligand structure was done by LigPlot + v.2.2.5 software (Wallace et al., 1995) to identify the interactions of amino acid between protein and ligand complex. LigPlot depicted hydrophobic bonds, hydrogen bonds, and their bond lengths in each docking pose.

Results

S.mutans is gram-positive pathogenic bacteria causing dental carie. Many treatments are already available for bacteria and various kinds of antibiotics and synthetic mouthwash are available to treat dental disease. The major problem with these chemicals is that they are toxic nature which is harmful for human health, and cause many diseases. Another problem is the drug resistance problem which arises due to the mutation in the target enzymes. To solve these entire problems, this research is focused mainly on the discovery and identification of some phytochemicals against SmdCD target . Hence our study may help identify potential therapeutic phytochemicals against targets SmdCD.

Construction of Phytochemical library

In this process takes more advantage in biomedical research and drug discovery. In this process, useful information is extracted from different data sources via the recognition and exploration of interesting patterns. However, finding the large number of medicinal plants is a laborious as well as time-consuming process. In this study, we were prepared library of 13 medicinal plants through text mining analysis Carrot2 and PubTator servers showed that these plants have potential anti-bacterial activity and phytochemicals of these plants can have anti-bacterial properties.

Hence to find out natural phytochemicals with an anti-bacterial activity against *S.mutans*, a library of 871 phytochemicals was constructed from 13 plants, namely, *Acacia nilotica*, *Acacia catechu*, *Allium sativum*, *Aloe vera*, *Azadirachta indica*, *Curcuma longa*, *Embelia ribes*,

Glycyrrhiza glabra, *Hemidesmus indicus*, *Juglans regia*, *Mangifera indica*, *Mimusops elengi* and *phyllanthus emblica* through searching the scientific literature and download from PubChem server. The 3D structures of phytochemicals were generated from PubChem Server in sdf format and then convert into pdb format by using Open Babel software.

Drug-likeness prediction

Currently, due to continuous advancement in computer science, lot of successful findings drugs from natural products using computer aided drug design methods for example the development of Dazamide, Imatinib, Dasatinib and Ponatinib etc (Ghose et al., 1999). The rationale behind these *in silico* approaches are due to relatively lower cost time factor involved compared to standard ADMET profiling (Darvas et al., 2002; DiMasi et al., 2003). In the present study we used SwissADME online software tool which is available free for the users to evaluate the ADME properties.

The results obtained from *in silico* studies clearly indicate 211 phytochemicals were showed the most druggable substance with a zero violation from any of drug likeness rules. It was interesting to note that the results from the SwissADME predictor values of Log P with the most important rules of drug likeness. eg. Lipinski, Ghose, Veber, Egan etc. Though these phytochemicals were exhibiting good hydrophilic lipophilic balance and same predicted bioavailability, the hydroxy derivative with high lipophilicity was expecting to show decent GI absorption. This hydroxy derivative with a higher value of probability of antibacterial activity and non-carcinogenic and mutagenic properties were predicted as the lead in the study.

Molecular docking

All the 211 selected phytochemicals were docked with SmdCD (PDB ID (5C2O)) using PyRx software by selecting AutoDock Vina as the docking engine to find the reasonable binding geometry and discover the protein-ligand complex, and it was found that the 3 phytochemicals have good binding affinity to the receptors as compared to reference molecule. Before performing the virtual screening, validation of the protocol was done by re-docking the reference compound (TTP) into the molulator site of SmdCD. The result showed that the docked TTP was completely superimposed with co-crystallized TTP in PDB ID(5C2O). The results showed that

all selected inhibitors were in the pocket of the target SmdCD, exhibiting a possible interaction and ranked based binding energies with SmdCD on specific binding pocket (Table 1). The binding energies of the screened phytochemicals were found to be in the range -7.9 kcal/mol to -7.5 kcal/mol and indicate good inhibition of the enzyme. 3 phytochemicals were observed to better fit strong binding in the allosteric substrate pocket. The binding energy of the screened phytochemicals was found in the following order; Cyclocurcumin(-7.9 kcal/mol) = Elatin(-7.9 kcal/mol) > Androsta-1,4,6-triene-3,17-dione(-7.6 kcal/mol) = SmdCD reference ligand(-7.5 kcal/mol). The results obtained were given in Table 1. Docking results are ranked based on binding energies. Now, after docking studies all these 3 successful natural screened phytochemicals further proceeded for toxicity prediction.

Table1: Molecular docking scores of various screened Phytochemicals with the SmdCD.

S.No	Protein-ligand complex	Compound ID	SMILES structure of compound	Binding Affinity (kcal/mol)
1	SmdCD - Reference	64968	<chem>CC1=CN(C(=O)NC1=O)C2CC(C(O2)COP(=O)(O)OP(=O)(O)OP(=O)(O)O)</chem>	-7.5
2	SmdCD - Cyclocurcumin	69879809	<chem>COc1cc(/C=C/C2=CC(=O)C[C@H](O2)c2ccc(c(c2)OC)O)ccc1O</chem>	-7.9
3	SmdCD -Androsta-1,4,6-triene-3,17-dione	104880	<chem>O=C1C=C[C@]2(C(=C1)C=C[C@@H]1[C@@H]2CC[C@]2([C@H]1CCC2=O)C)C</chem>	-7.6
4	SmdCD -Elatin	44257938	<chem>Oc1ccc(cc1)c1oc2cc(O)cc(c2c(=O)c1O)O</chem>	-7.9

Toxicity prediction

The above three phytochemicals were also further proceeded to predict their tumorigenicity, mutagenicity, irritation, and reproductive toxicity by the OSIRIS tool. The predicted toxicity of three phytochemicals is shown in Table 2. According to Table 2, the 1 phytochemical i.e. Elatin showed a high risk of toxicity, while, 2 phytochemicals i.e. Cyclocurcumin, Androsta-1,4,6-triene-3,17-dione, were non-toxic (non-mutagenic, non-tumorigenic, non-irritant, and no reproductive effect). The drug-score show ranges between 0 to 1, where as the value 1 indicates the possibility of a compound to be drug molecule, whereas, the score value 0 indicates that compounds having no possibilities of drug candidates. Therefore, these 2 phytochemicals

(Cyclocurcumin, Androsta-1,4,6-triene-3,17-dione,) can be further subjected to study their binding interaction with SmdCD.

Table 2. Toxicity profile of the reference molecule and hit phytochemicals

S. No.	Compound Name	TUM	MUT	IRR	REP	Drug-Score
1	Reference(TTP)	Non Toxic	Non Toxic	Non Toxic	Non Toxic	0.39
2	Cyclocurcumin	Non Toxic	Non Toxic	Non Toxic	Non Toxic	0.73
3	Androsta-1,4,6-triene-3,17-dione	Non-Toxic	Non-Toxic	Non-Toxic	Non-Toxic	0.75
4	Elatin	Toxic	Non-Toxic	Non-Toxic	Non-Toxic	0.46

Visualization

LigPlot+ v.1.4.5 was used to visualize the protein-ligand interactions. The docked poses of these two compounds with SmdCD is shown in Figure 3. SmdCD -reference the docked TTP showed interaction with the amino acid residues by hydrogen and hydrophobic bonds as found in the experimental structure shown in Figure 2. It forms seven hydrogen-bonds with five residues Thr42, Tyr44, Asn53, Thr23 and Asn75 of SmdCD. It also formed the five hydrophobic bonds with residues Ala49, Gly47, Gly46, Gly43 and Gln79 of SmdCD as shown in Figure 3. According to Figure. 2(A), SmdCD - Cyclocurcumin formed one hydrogen bonds with Thr42 which have the bond distance 3.08 Å and It also formed nine hydrophobic bonds with Ala49, Gln79, Ala41, Gly47, Gly46, Asn75, Tyr44, Val48, and Lys82. SmdCD - Androsta-1,4,6-triene-3,17-dione interacted with Tyr44 that make one hydrogen bonds having the bond distance 3.14 Å and seven hydrophobic bonds with Asn53, Gly47, Asn75, Gly43, Gln79, Ala49 and Asp50, residues were found to participate in SmdCD - Androsta-1,4,6-triene-3,17-dione complex showed in table 3.

Table 3. 2D details Interactions between the SmdCD and top hits after the molecular docking. The bold residues represent the common interacted residues between SmdCD-reference complex and SmdCD -screened phytochemical complex.

S. No	Compounds Name	Number of H-bonds	Interacted residues with SmdCD	Common active site residues
1	Reference(TTP)	7	Ala49, Gly46, Gly47, Gln79, Gly43, Thr23, Tyr44, Asn53, Asn75, Thr42	Ala49, Gly47, Gln79, Tyr44, Asn75
2	SmdCD - Cyclocurcumin	1	Tyr44, Gly46, Gly47, Asn75, Ala41, Thr42, Gln79, Ala49, Val48, Lys82	
3	SmdCD - Androsta-1,4,6-triene-3,17-dione	1	Ala49, Gly47, Gln79, Asn75, Tyr44, Asn53, Gly43, Asp50	




Figure 3. 2D interaction of protein-ligand complexes with Hydrogen and Hydrophobic-bonds between hit phytochemicals (reference ligand, Cyclocurcumin, and Androsta-1,4,6-triene-3,17-dione) and dotted green lines denote hydrogen bonds, red half arcs indicate hydrophobic interactions.

Discussion and Conclusion

Despite the improvement in pharmacology and conventional chemistry in developing rapidly new synthetic antibiotics, altering existing antibiotics or finding suitable protein targets against which inhibitors can be developed. Present global drug development programs may not be able to afford new effective antibiotics for the next decade (Abdallah EM., 2011). In our study, we used different compounds from selected medicinal plant against allosteric substrate of SmdCD. These are all of the plants used in Ayurveda and the ancient medicinal system with antibacterial, anti-inflammatory, antiviral, antioxidant, anticancer, and antidiabetic activities. *Glycyrrhiza glabra* is one of the extensively used herbs from the ancient medicinal history of Ayurveda. The antimicrobial activity of *G. glabra* has been both researched and exploited medicinally for many years. Natural Plant products (NPPs) of *G. glabra* is considered as antitussive, mucolytic, expectorant, antimicrobial, immunostimulant as well as a flavoring agent (Vandeputte et al., 2011). Saponins have also been reported to possess antimicrobial activity due to their detergent like nature they can cause leakage in the membrane by interacting with proteins and certain enzymes from the bacterial cell (S.A. Tamil et al., 2011). Rationale based selection of *G. glabra* against *P. aeruginosa* by employing bioprospection, in silico and in vitro study is reported previously (Chakotiya, Chawla, et al., 2016; Chakotiya, Tanwar, et al., 2016).. Curcumin is a active component *C. longa* is having many properties such as antioxidant, anti-inflammatory, anti-viral, antibacterial, antifungal and anticancer activities and also works against various malignant diseases such as diabetes (Spinetti et al., 2013), arthritis, Alzheimer's (Mishra & Palanivelu, 2008) and other chronic diseases has been reported. Cyclocurcumin, a curcumin derivative of *C. longa*, exhibits immune-modulating ability and is a potential phytochemical for the treatment of rheumatoid arthritis. TNF- α is a key factor in a variety of inflammatory diseases. The role of cyclocurcumin in overcoming p38 α -induced production of TNF- α and hence can be used as a therapeutic agent to target rheumatoid arthritis (Fu et al., 2017). Cyclocurcumin can be used as a herbal drug or proved to be a good lead compound for oral and cervical cancers

(Nomin et al., 2005; ULLMAN et al., 1996). The information is quite significant in drug development for oral and cervical cancers. Here in this study, we employed *in silico* techniques to investigate the natural compound Cyclocurcumin and Androsta-1,4,6-triene-3,17-dione as possible to control biofilm formation.

Therefore, to find out potential phytochemicals, we prepared a library of phytochemicals of 10 selected medicinal plant and buildup a library of 871 phytochemical. Afterthat, It is used to filtered a library of 871 phytochemical through Swiss ADME web tool. Further filtered 211 phytochemical were subjected to molecular docking against SmdCD. Based on Virtual screening of Elatin phytochemical of selected pants we found the top 3 phytochemicals against SmdCD, namely Cyclocurcumin, Androsta-1,4,6-triene-3,17-dione, Elatin, and all these compounds showed good binding energy with SmdCD as compared to reference compounds. After virtual screening we were checked toxicity prediction of screened phytochemicals through OSIRIS softwere. Now, we get the result out of 3 phytochemicals, 2 phytochemical were non toxic in nature. Through these results, we can suggest these phytochemicals can be used against SmdCD target. Our drug repurposing study, both phytochemicals were found to inhibit the SmdCD and these phytochemicals may be used against the *S.mutans* microbial infection. Finally, we suggest based on a future perspective on the *in vitro* and *in vivo* research, that these phytochemicals namly Cyclocurcumin, and Androsta-1,4,6-triene-3,17-dione may become development of organic mouthwash and potent anti biofilm drugs to oral care.

Reference

- Abdallah EM. (2011). Plants: an alternative source for antimicrobials. . *J App Pharm Sci.*, 1, 16–20.
- Abdel-Aziz, M. M., Emam, T., & Raafat, M. M. (2020). Hindering of Cariogenic Streptococcus mutans Biofilm by Fatty Acid Array Derived from an Endophytic *Arthrographis kalrae* Strain. *Biomolecules*, 10(5), 811. <https://doi.org/10.3390/biom10050811>
- Algburi, A., Comito, N., Kashtanov, D., Dicks, L. M. T., & Chikindas, M. L. (2017). Control of Biofilm Formation: Antibiotics and Beyond. *Applied and Environmental Microbiology*,

83(3). <https://doi.org/10.1128/AEM.02508-16>

Bowen, W. H., & Koo, H. (2011). Biology of <i>Streptococcus mutans</i>-Derived Glucosyltransferases: Role in Extracellular Matrix Formation of Cariogenic Biofilms.

Caries Research, 45(1), 69–86. <https://doi.org/10.1159/000324598>

Cabral, C. T., & Fernandes, M. H. (2007). In vitro comparison of chlorhexidine and povidone-iodine on the long-term proliferation and functional activity of human alveolar bone cells.

Clinical Oral Investigations, 11(2), 155–164. <https://doi.org/10.1007/s00784-006-0094-8>

Chakotiya, A. S., Chawla, R., Thakur, P., Tanwar, A., Narula, A., Grover, S. S., Goel, R., Arora, R., & Sharma, R. K. (2016). In vitro bactericidal activity of promising nutraceuticals for targeting multidrug resistant *Pseudomonas aeruginosa*. *Nutrition*, 32(7–8), 890–897.

<https://doi.org/10.1016/j.nut.2016.01.024>

Chakotiya, A. S., Tanwar, A., Narula, A., & Sharma, R. K. (2016). Alternative to antibiotics against *Pseudomonas aeruginosa*: Effects of *Glycyrrhiza glabra* on membrane permeability and inhibition of efflux activity and biofilm formation in *Pseudomonas aeruginosa* and its in vitro time-kill activity. *Microbial Pathogenesis*, 98, 98–105.

<https://doi.org/10.1016/j.micpath.2016.07.001>

Daina, A., Michielin, O., & Zoete, V. (2017). SwissADME: a free web tool to evaluate pharmacokinetics, drug-likeness and medicinal chemistry friendliness of small molecules.

Scientific Reports, 7(1), 42717. <https://doi.org/10.1038/srep42717>

Darvas, F., Keseru, G., Papp, A., Dorman, G., Urge, L., & Krajcsi, P. (2002). In Silico and Ex Silico ADME Approaches for Drug Discovery. *Current Topics in Medicinal Chemistry*,

2(12), 1287–1304. <https://doi.org/10.2174/1568026023392841>

DiMasi, J. A., Hansen, R. W., & Grabowski, H. G. (2003). The price of innovation: new estimates of drug development costs. *Journal of Health Economics*, 22(2), 151–185.

[https://doi.org/10.1016/S0167-6296\(02\)00126-1](https://doi.org/10.1016/S0167-6296(02)00126-1)

Egan, W. J., Merz, , Kenneth M., & Baldwin, J. J. (2000). Prediction of Drug Absorption Using Multivariate Statistics. *Journal of Medicinal Chemistry*, 43(21), 3867–3877.

<https://doi.org/10.1021/jm000292e>

Fu, M., Chen, L., Zhang, L., Yu, X., & Yang, Q. (2017). Cyclocurcumin, a curcumin derivative, exhibits immune-modulating ability and is a potential compound for the treatment of rheumatoid arthritis as predicted by the MM-PBSA method. *International Journal of Molecular Medicine*, 39(5), 1164–1172. <https://doi.org/10.3892/ijmm.2017.2926>

Gao, L., Xu, T., Huang, G., Jiang, S., Gu, Y., & Chen, F. (2018). Oral microbiomes: more and more importance in oral cavity and whole body. *Protein & Cell*, 9(5), 488–500. <https://doi.org/10.1007/s13238-018-0548-1>

Ghose, A. K., Viswanadhan, V. N., & Wendoloski, J. J. (1999). A Knowledge-Based Approach in Designing Combinatorial or Medicinal Chemistry Libraries for Drug Discovery. 1. A Qualitative and Quantitative Characterization of Known Drug Databases. *Journal of Combinatorial Chemistry*, 1(1), 55–68. <https://doi.org/10.1021/cc9800071>

Hernandez-Santiago, B., Placidi, L., Cretton-Scott, E., Faraj, A., Bridges, E. G., Bryant, M. L., Rodriguez-Orengo, J., Imbach, J. L., Gosselin, G., Pierra, C., Dukhan, D., & Sommadossi, J. P. (2002). Pharmacology of β -Thymidine and β -2'-Deoxycytidine in HepG2 Cells and Primary Human Hepatocytes: Relevance to Chemotherapeutic Efficacy against Hepatitis B Virus. *Antimicrobial Agents and Chemotherapy*, 46(6), 1728–1733. <https://doi.org/10.1128/AAC.46.6.1728-1733.2002>

Jones, C. G. (1997). Chlorhexidine: is it still the gold standard? *Periodontology 2000*, 15(1), 55–62. <https://doi.org/10.1111/j.1600-0757.1997.tb00105.x>

Jr, R. P., Chalmers, N., Rickard, A., & Kolenbrander, P. (2008). Community Development in Bacterial Biofilms of the Oral Cavity. *Microscopy and Microanalysis*, 14(S2), 1554–1555. <https://doi.org/10.1017/S1431927608088831>

Li, Y., Guo, Z., Jin, L., Wang, D., Gao, Z., Su, X., Hou, H., & Dong, Y. (2016). Mechanism of the allosteric regulation of *Streptococcus mutans* 2'-deoxycytidylate deaminase. *Acta Crystallographica Section D Structural Biology*, 72(7), 883–891. <https://doi.org/10.1107/S2059798316009153>

- Lipinski, C. A. (2000). Drug-like properties and the causes of poor solubility and poor permeability. *Journal of Pharmacological and Toxicological Methods*, 44(1), 235–249. [https://doi.org/10.1016/S1056-8719\(00\)00107-6](https://doi.org/10.1016/S1056-8719(00)00107-6)
- Lua, R. C., & Lichtarge, O. (2010). PyETV: a PyMOL evolutionary trace viewer to analyze functional site predictions in protein complexes. *Bioinformatics*, 26(23), 2981–2982. <https://doi.org/10.1093/bioinformatics/btq566>
- Marsh, P. D. (2010). Microbiology of Dental Plaque Biofilms and Their Role in Oral Health and Caries. *Dental Clinics of North America*, 54(3), 441–454. <https://doi.org/10.1016/j.cden.2010.03.002>
- Mishra, S., & Palanivelu, K. (2008). The effect of curcumin (turmeric) on *Alzheimer's disease* : An overview. *Annals of Indian Academy of Neurology*, 11(1), 13. <https://doi.org/10.4103/0972-2327.40220>
- Moynihan, P. (2016). Sugars and Dental Caries: Evidence for Setting a Recommended Threshold for Intake. *Advances in Nutrition*, 7(1), 149–156. <https://doi.org/10.3945/an.115.009365>
- Nomin[◆], Y., Charbonnier, S., Miguet, L., Potier, N., Dorsselaer, A. Van, Atkinson, R. A., Trav[◆], G., & Kieffer, B. (2005). 1H and 15N resonance assignment, secondary structure and dynamic behaviour of the C-terminal domain of human papillomavirus oncoprotein E6. *Journal of Biomolecular NMR*, 31(2), 129–141. <https://doi.org/10.1007/s10858-004-7802-y>
- O'Boyle, N. M., Banck, M., James, C. A., Morley, C., Vandermeersch, T., & Hutchison, G. R. (2011). Open Babel: An open chemical toolbox. *Journal of Cheminformatics*, 3(1), 33. <https://doi.org/10.1186/1758-2946-3-33>
- Rath, S. K., & Singh, M. (2013). Comparative clinical and microbiological efficacy of mouthwashes containing 0.2% and 0.12% chlorhexidine. *Dental Research Journal*, 10(3), 364–369.
- Reibel, J. (2003). Tobacco and Oral Diseases. *Medical Principles and Practice*, 12(Suppl. 1), 22–32. <https://doi.org/10.1159/000069845>
- Ribeiro, D. A., Bazo, A. P., da Silva Franchi, C. A., Marques, M. E. A., & Salvadori, D. M. F.

- (2004). Chlorhexidine induces DNA damage in rat peripheral leukocytes and oral mucosal cells. *Journal of Periodontal Research*, 39(5), 358–361. <https://doi.org/10.1111/j.1600-0765.2004.00759.x>
- S.A. Tamil, M.G. Dinesh, R.S. Satyan, B. Chandrasekaran, & C. Rose. (2011). Leaf and seed extracts of *Bixa orellana* L. exerts anti-microbial activity against bacterial pathogens. *J. Appl. Pharm. Sci.* , 1, 116–120.
- Sander, T., Freyss, J., von Korff, M., Reich, J. R., & Rufener, C. (2009). OSIRIS, an Entirely in-House Developed Drug Discovery Informatics System. *Journal of Chemical Information and Modeling*, 49(2), 232–246. <https://doi.org/10.1021/ci800305f>
- Spinetti, G., Fortunato, O., Caporali, A., Shantikumar, S., Marchetti, M., Meloni, M., Descamps, B., Floris, I., Sangalli, E., Vono, R., Faglia, E., Specchia, C., Pintus, G., Madeddu, P., & Emanuelli, C. (2013). MicroRNA-15a and microRNA-16 impair human circulating proangiogenic cell functions and are increased in the proangiogenic cells and serum of patients with critical limb ischemia. *Circulation Research*, 112(2), 335–346. <https://doi.org/10.1161/CIRCRESAHA.111.300418>
- Trott, O., & Olson, A. J. (2009). AutoDock Vina: Improving the speed and accuracy of docking with a new scoring function, efficient optimization, and multithreading. *Journal of Computational Chemistry*, NA-NA. <https://doi.org/10.1002/jcc.21334>
- ULLMAN, C. G., HARIS, P. I., GALLOWAY, D. A., EMERY, V. C., & PERKINS, S. J. (1996). Predicted α -helix/ β -sheet secondary structures for the zinc-binding motifs of human papillomavirus E7 and E6 proteins by consensus prediction averaging and spectroscopic studies of E7. *Biochemical Journal*, 319(1), 229–239. <https://doi.org/10.1042/bj3190229>
- Vandeputte, O. M., Kiendrebeogo, M., Rasamiravaka, T., Stévigny, C., Duez, P., Rajaonson, S., Diallo, B., Mol, A., Baucher, M., & El Jaziri, M. (2011). The flavanone naringenin reduces the production of quorum sensing-controlled virulence factors in *Pseudomonas aeruginosa* PAO1. *Microbiology*, 157(7), 2120–2132. <https://doi.org/10.1099/mic.0.049338-0>
- Vellappally, S., Divakar, D. D., Al Kheraif, A. A., Ramakrishnaiah, R., Alqahtani, A., Dalati, M. H. N., Anil, S., Khan, A. A., & Harikrishna Varma, P. R. (2017). Occurrence of

vancomycin-resistant *Staphylococcus aureus* in the oral cavity of patients with dental caries. *Acta Microbiologica et Immunologica Hungarica*, 64(3), 343–351.
<https://doi.org/10.1556/030.64.2017.033>

Wallace, A. C., Laskowski, R. A., & Thornton, J. M. (1995). LIGPLOT: a program to generate schematic diagrams of protein-ligand interactions. *Protein Engineering, Design and Selection*, 8(2), 127–134. <https://doi.org/10.1093/protein/8.2.127>

UNDER PEER REVIEW