

Short Research Article

ASSESSMENT OF THE KNOWLEDGE, ATTITUDE AND PRACTICE OF HEALTHCARE WORKERS ABOUT HAND HYGIENE

ABSTRACT

Aims: The purpose of this study was to assess the knowledge, attitude and practice of healthcare workers about hand hygiene.

Study design: A cross-sectional study was used.

Place and Duration of Study: Memon Medical Institute Hospital, a tertiary care private hospital Karachi, Pakistan from November 2021 to December 2021.

Methodology: Non-random convenience sampling method was used in this study.

Results: Most participants have an acceptable level of knowledge of hand hygiene. The highest scores were in the infection control domain and the lowest scores defined the hand hygiene domain. Multivariate analysis showed that work experience and previous education were the most important predictors of participants' knowledge of hand hygiene.

Conclusion: Hand hygiene is the simplest yet effective way to stop the spread of germs/microbes non-invasive independent predictors for screening esophageal varices may decrease medical as well as financial burden, hence improving the management of cirrhotic patients. These predictors, however, need further work to validate reliability.

Comment [OF1]: reduce

Keywords: Hand hygiene, nurses, healthcare workers, infection-prevention, knowledge, attitude, practice.

1. INTRODUCTION

Hand hygiene is the compliance of cleaning hands with cleaning soap and water or with antiseptic hand rub to remove temporary microorganisms from hands and preserve the circumstance of the skin⁽¹⁾. It is one of the most significant issues in the world and the chain of infection could be broken by hand hygiene compliance and to decrease health related infections⁽²⁾. There would be lots of microorganisms in the hand during the dealing with patients unless there is hand hygiene compliance following recommended guidelines^(3, 4). Morbidity, length of stay, and treatment cost increases with Healthcare associated infections (HAI), which can be reduced by following infection control guidelines⁽⁵⁾.

Comment [OF2]: broken

Comment [OF3]:

The mode of transmission mainly begins with healthcare provider's hands⁽⁶⁾. Infection control guideline should be followed to reduce hand related microorganisms spreading. The healthcare workers wash their hands properly before and after touching the patient to prevent HAI infections⁽⁶⁾. With the decrease in infection rates reported after better compliance with handwashing, it states to the professional practice of hand hygiene⁽⁷⁾.

Comment [OF4]: ???????

Transmission of the pathogen among the patients can be potentially caused by the hands of healthcare workers (HCW). Manifestations of the hand hygiene programs reached a high standard in care for the patients and decrease healthcare-associated infections (HAIs) by about one infection per 1000 patients^(7, 8). 9% is the prevalence of HAI in Southeast Asia, between 2.5% and 14.8% in Africa^(9, 10). Whereas, 7.6% prevalence in diverged patient population in high-wages countries⁽¹¹⁾.

Comment [OF5]: through

Comment [OF6]: Don't start a sentence with a figure.
Suggestion:
Prevalence of HAI in Southeast Asia is 9%, and between 2.5% and 14.8% in Africa whereas it is 7.6% in diverged patient populations in high-waged (high-income) countries

Patients admitted in Intensive Care Units (ICUs) have a higher rate of incidence of HAI. Though, in certain contexts, the frequency of patient contact may be too great to ensure 100% hand hygiene

Comment [OF7]: Be mindful about the use of these terms; prevalence and incidence, please!

compliance among HCWs^(12, 13). Invasive devices commonly utilized on ICU patients act as entrance points for virulent microorganisms, resulting in a higher risk of HAIs⁽¹⁴⁾.

Comment [OF8]: Tense!

The proper compliance with hand hygiene can decrease the spread of HAIs, length of stay in hospital, HAIs related mortality and morbidity, healthcare expenses, and encourages the patient's health and safety which have been proved by the various studies^(15, 16). The practice of hand hygiene for HCWs has ~~vebeen~~ recommended by the international public health agencies including the World Health Organization (WHO). However, compliance remains low presently^(17, 18), hence the need for continuous surveillance.

2. METHODOLOGY

A cross sectional method was conducted among healthcare providers at Memon Medical Institute Hospital in 2021. The samples consisted of -was 209 healthcare workers. Sample size was calculated by Open Epi software. They HCWs were invited to participate in the study through a non-random convenient sampling method. Inclusion criteria included Healthcare workers who have minimum 1-year clinical experience, Aged 20 to 50 years, Doctors, nurses, midwives having valid PMDC or PNC License. Exclusion criteria consisted of non-clinical staff, not willing to be part of study.

The research tool was WHO questionnaire named as "HAND HYGIENE KNOWLEDGE QUESTIONNAIRE". The questionnaire consisted of two parts, first part covered demographic data and second part was based on 9 questions for ~~healthcare worker~~ HCWs regarding ~~to~~ the knowledge of hand hygiene practice/procedure. In addition to questionnaire responses, demographic detail was collected including participant's educational institution, age, gender, and previous experience of working in the healthcare industry.

Data analyzed using SPSS version 21. Data collection was conducted in November 2021 after the ethical approval by IRB of Memon Medical Institute Hospital.

3. RESULTS

Table 1: shows ed the demographic data of the study participants. e Out of 209 study participants, majority were female 135 (64.6%) and 74 (35.4%) were male participants. Out of 209 participants ~~majority were~~ 39.23% were in the age group of 20-25 years. Majority of participants (74.16%), had more than one-year (work) experience. More than half of the participants were- had educational qualification with Diploma in Nursing:- 11% had BSN, 21.53% had MBBS, 9.09% had CMW, 4.3 % had Nursing Assistant, and 2.87% had DPT. Most of the participants were Nurses (134, (64.11%) while 23 (20/58%) were doctors, 14(6.69%) were midwives, 10 (4.78%) were Nursing Assistants, 2 (0.95%) were technicians, and 6 (2.87%) therapists.

Comment [OF9]: majority were females or majority was female


Comment [OF10]: Do you mean 20.58%?

Table 1 Socio Demographic Characteristics of Study Participants

Characteristics	n	%
Age (years)		
20-25	82	39.23
26-30	77	36.84
31-35	43	20.57
36-40	6	2.87
41-45	1	0.47
Gender		
Male	74	35.4
Female	135	64.6


Marital Status

Married	112	53.58
Unmarried	92	44.01
Divorced	5	2.39
Working Experience		
1-year	32	15.31
less than 1 year	22	10.52
more than 1 year	155	74.16
Education		
BSN	23	11
MBBS	45	21.53
CMW	19	9.09
Nursing Assistant	9	4.3
Diploma in Nursing	107	51.19
DPT	6	2.87
Designation		
Nurse	134	64.11
Healthcare assistant	10	4.78
Midwife	14	6.69
Doctor	43	20.574
Technician	2	0.95
Therapist	6	2.87




Graph Fig. 1. Main route of cross transmission of potentially harmful germs between patients in health care facility.

Formatted: Right



Graph Fig. 2. Shows the most frequent source of germs responsible for health care-associated infections



Graph Fig. 3. Shows the minimal time needed for alcohol-based hand rub to kill most germs on hands

Graph Fig. 1 shows the Main route of cross-transmission of potentially harmful germs between patients in health care facilities. Most of the participants (74, 35%) said health-care-worker HCWs' hands were not clean. While air circulation in hospital 44 (21.05%) and Patients exposure to colonized surfaces 55 (26.32%) and sharing noninvasive objectives are 36 (17%)

Comment [OF11]: Tense

Graph Fig. 2 show depicts the most frequent sources of germs responsible for health care associated infections. Most of the participants (78, 37.32%) answered that the hospital environment is the most frequent source of germs responsible for health care associated infections.

Graph Fig.3. shows the most minimal time needed for alcohol-based hand rub to kill most germs on hands. Most of the participants (128, 60.29%) said-revealed that the minimal time needed for alcohol-based hand rub to kill most germs on hands is 20 seconds.

Table 2 show presents the Practice of health care provider about HH. The study group categorized on the base of formal training received or not more than half the participants 110 (52.32%) received formal training for HH in the last three years. Most of the participants (179, 85%) said they use alcohol-based hand rub for hand hygiene while 44 (21.05%) do not use alcohol-based hand rub for hand hygiene.

Table 2. HAND HYGIENE PRACTICE QUESTIONNAIRE

Characteristics	N	%
Did you receive formal training in hand hygiene practice in the last three years?		
Yes	99	47.37
No	110	52.63
Do you routinely use an alcohol-based hand rub for hand hygiene?		
Yes	179	85.65
No	30	14.35

Table 3 shows the study participants knowledge about HH. Study group was categorized based on whether they believed that the use of hand hygiene before touching the patients prevents transmission of germs, 204 (97.61%) said yes and 5(2.39%) no. When participants were questioned about whether Hand hygiene immediately before a clean/aseptic procedure prevents transmission of germs, 203 (96.7%) said yes and 6 (2.87%) said no. When participants were asked about whether Alcohol-based hand rub and handwashing with soap and water is true immediately after a risk of body fluid exposure 99% participants said yes and only 1% said no. When participants were asked about whether wearing jewelry should be avoided, as associated with increased likelihood of colonization of hands with harmful germs 201 (96.17%) said yes and 6 (3.87%) said no. When participants were asked about whether Damaged skin should be avoided, as associated with increased likelihood of colonization of hands with harmful germs, most of the participants (197, 97.26%) said yes and 12 (5.74%) said no. When participants were asked about Whether Artificial fingernails should be avoided, as associated with increased likelihood of colonization of hands with harmful germs, most of the participants (180, 86.12%) said yes and 29 (13.88%) said no. When participants were asked about Whether Regular use of a hand cream should be avoided, as associated with increased likelihood of colonization of hands with harmful germs more than half of the participants 110 (52.63%) said yes and 99 (47.33%) said no.

Table 3. HAND HYGIENE KNOWLEDGE QUESTIONNAIRE

Characteristics	n	%
Hand hygiene before touching the patient prevents transmission of germs		
Yes	204	97.61
No	5	2.39
Hand hygiene Immediately after a risk of body fluid exposure prevents transmission of germs		
Yes	203	97.13
No	6	2.87
Hand hygiene Immediately before a clean/aseptic procedure prevents transmission of germs		
Yes	201	96.17
No	8	3.83
Hand hygiene after exposure to the immediate surroundings of a patient prevents transmission of germs		
Yes	209	100.00
No	0	0.00
Alcohol-based hand rub and handwashing with soap and water are true after touching a patient		

Yes	204	97.61
No	5	2.39
Alcohol-based hand rub and handwashing with soap and water are true immediately after a risk of body fluid exposure		
Yes	209	100
No	0	0
Alcohol-based hand rub and handwashing with soap and water are true after exposure to the immediate surroundings of a patient		
Yes	207	99.00
No	2	1.00
Hand rubbing is more rapid for hand cleansing than handwashing		
Yes	192	91.90
No	17	8.10
Hand rubbing causes skin dryness more than handwashing		
Yes	202	96.70
No	7	3.30
Wearing jewelry should be avoided, as associated with increased likelihood of colonization of hands with harmful germs?		
Yes	201	96.17
No	8	3.83
Damaged skin should be avoided, as associated with increased likelihood of colonization of hands with harmful germs?		
Yes	197	94.26
No	12	5.74
Artificial fingernails should be avoided, as associated with increased likelihood of colonization of hands with harmful germs?		
Yes	180	86.12
No	29	13.88
Regular use of a hand cream should be avoided, as associated with increased likelihood of colonization of hands with harmful germs?		
Yes	110	52.63
No	99	47.37

Formatted: Space After: 0 pt, Don't keep with next, Tab stops: 0.99", Left + 1.9", Left

[No evidence that statistical analysis was carried out as stated earlier \(SPSS, version 21\). No statistical values were provided and no comparative analysis of the data gathered.](#)

DISCUSSION

This study showed that ~~the~~ most of the participants had good knowledge about hand hygiene. ---% participants were aware of knowledge about hand hygiene through which the rate of nosocomial infection is also low in MMIH. Comparatively, India have reported HH compliance ranging from 20–85.5%.¹⁹ In Qassim, Saudi Arabia, 58% of healthcare workers have moderate knowledge of suggested hand hygiene.²⁰ Other similar studies have shown the influence of multiple variables on nurses' knowledge of hand hygiene.²¹

Comment [OF12]: ????????

Work experience and occupational type were significantly associated with nurses' knowledge of hand hygiene. Perhaps as nurses gain more practical experience in clinical settings and determine their employment patterns, they will be more motivated to learn more and comply with the latest guidelines.

Jang et al.²² discussed this issue in a different way, noting that as long as ~~health care worker~~ HCWs practice hand hygiene for self-protection, education on this has little impact. ~~And found~~ They also reported that educating caregivers on good communication patterns, teamwork, and how to adhere to hand hygiene guidelines are important factors despite heavy workloads.²²

Variations in knowledge among the hospitals surveyed, particularly with regard to hand hygiene definitions, were attributed to differences in hospital policies, ~~differences in~~ teaching methods, and ~~differences in~~ the conditions of selected neonatal wards (number of new-borns, needing a ventilator, etc.). Number of new-borns, or prior care, number of stations, equipment availability, and management style).²²

Comment [OF13]: RE-structure, please

4. CONCLUSION

Continue to implement structured, regular and ongoing training programs in a variety of effective ways to maintain and strengthen caregiver knowledge of hand hygiene and reduce and eliminate knowledge gaps; we need to increase accuracy and supervision. There is also a need to identify factors that promote nurses' positive perceptions and attitudes towards hand hygiene.

Comment [OF14]: These are recommendations which should be based on the conclusion reached. So no conclusion stated

ETHICAL APPROVAL

This research has been conducted after ethically approved by IRB committee of Memon Medical Institute Hospital.

REFERENCES

1. Pittet D. Improving adherence to hand hygiene practice: a multidisciplinary approach. *Emerging infectious diseases*. 2001 Mar;7(2):234.
2. Pittet D. Improving adherence to hand hygiene practice: a multidisciplinary approach. *Emerging infectious diseases*. 2001 Mar;7(2):234.
3. Mathur P. Hand hygiene: back to the basics of infection control. *The Indian journal of medical research*. 2011 Nov;134(5):611.
4. Pittet D, Allegranzi B, Boyce J, World Health Organization World Alliance for Patient Safety First Global Patient Safety Challenge Core Group of Experts. The World Health Organization guidelines on hand hygiene in health care and their consensus recommendations. *Infection Control & Hospital Epidemiology*. 2009 Jul;30(7):611-22.
5. Nejad SB, Allegranzi B, Syed SB, Ellis B, Pittet D. Health-care-associated infection in Africa: a systematic review. *Bulletin of the World Health Organization*. 2011;89:757-65.
6. Martín-Madrado C, Salinero-Fort MA, Abanades-Herranz JC, Arnal-Selfa R, García-Ferradal I, Espejo-Matorral F, Carrillo-de Santa-Pau E, Soto-Díaz S. Effectiveness of a training programme to improve hand hygiene compliance in primary healthcare. *BMC Public Health*. 2009 Dec;9(1):1-8.
7. Chen YC, Sheng WH, Wang JT, Chang SC, Lin HC, Tien KL, Hsu LY, Tsai KS. Effectiveness and limitations of hand hygiene promotion on decreasing healthcare-associated infections. *PLoS one*. 2011 Nov 16;6(11):e27163.
8. Labrague LJ, McEnroe-Petitte DM, Van de Mortel T, Nasirudeen AM. A systematic review on hand hygiene knowledge and compliance in student nurses. *International nursing review*. 2018 Sep;65(3):336-48.
9. Sands M, Aunger R. Determinants of hand hygiene compliance among nurses in US hospitals: A formative research study. *PLoS one*. 2020 Apr 7;15(4):e0230573.
10. Gould DJ, Moralejo D, Drey N, Chudleigh JH, Taljaard M. Interventions to improve hand hygiene compliance in patient care. *Cochrane database of systematic reviews*. 2017(9).
11. Ceylan B, Gunes U, Baran L, Ozturk H, Sahbudak G. Examining the hand hygiene beliefs and practices of nursing students and the effectiveness of their handwashing behaviour. *Journal of Clinical Nursing*. 2020 Nov;29(21-22):4057-65.
12. Ng WK, Shaban RZ, van de Mortel T. The effect of a hand hygiene program featuring tailored religion-relevant interventions on healthcare workers' hand rubbing compliance and beliefs in

- the United Arab Emirates: A cohort study. *Infection, disease & health*. 2019 Aug 1;24(3):115-23.
13. Suchitra JB, Devi NL. Impact of education on knowledge, attitudes and practices among various categories of health care workers on nosocomial infections. *Indian journal of medical microbiology*. 2007 Jul 1;25(3):181-7.
 14. Randle J, Clarke M, Storr J. Hand hygiene compliance in healthcare workers. *Journal of Hospital Infection*. 2006 Nov 1;64(3):205-9.
 15. Al-Hussami M, Darawad M, Almhairat II. Predictors of compliance handwashing practice among healthcare professionals. *Healthcare Infection*. 2011 Jun 23;16(2):79-84.
 16. Biswal M, Angrup A, Rajpoot S, Kaur R, Kaur K, Kaur H, Dhaliwal N, Arora P, Gupta AK. Hand hygiene compliance of patients' family members in India: importance of educating the unofficial 'fourth category' of healthcare personnel. *Journal of Hospital Infection*. 2020 Apr 1;104(4):425-9.
 17. Biswal M, Angrup A, Rajpoot S, Kaur R, Kaur K, Kaur H, Dhaliwal N, Arora P, Gupta AK. Hand hygiene compliance of patients' family members in India: importance of educating the unofficial 'fourth category' of healthcare personnel. *Journal of Hospital Infection*. 2020 Apr 1;104(4):425-9.
 18. Randle J, Clarke M, Storr J. Hand hygiene compliance in healthcare workers. *Journal of Hospital Infection*. 2006 Nov 1;64(3):205-9.
 19. Bharara T, Gur R, Duggal S, Chugh V. Evaluation of hand hygiene compliance over the years, in an intensive care unit of a north Delhi hospital preparing for accreditation: A 3-year study. *Journal of Family Medicine and Primary Care*. 2020 Apr;9(4):1939.
 20. Abalkhail A, Mahmud I, Alhumaydhi FA, Alslamah T, Alwashmi AS, Vinnakota D, Kabir R. Hand hygiene knowledge and perception among the healthcare workers during the COVID-19 pandemic in Qassim, Saudi Arabia: a cross-sectional survey. *InHealthcare* 2021 Nov 24 (Vol. 9, No. 12, p. 1627). MDPI.
 21. Parmeggiani C, Abbate R, Marinelli P, Angelillo IF. Healthcare workers and health care-associated infections: knowledge, attitudes, and behavior in emergency departments in Italy. *BMC infectious diseases* . 2010; 10 (1): 35
 22. Jang JH, Wu S, Kirzner D, Moore C, Youssef G, Tong A. et al. Focus group study of hand hygiene practice among healthcare workers in a teaching hospital in Toronto, Canada. *Infect Control Hosp Epidemiol* . 2010; 31 (2):144–50

[Kindly follow the format of the Journal for listing of References. I think tht there is no need to state the month of publication once the volume and number of Journal have been stated, please.](#)

[Listing publications that appeared in 2022 would definitely enhance the value of the article.](#)

APPENDIX

QUESTIONNAIRE

- Tick **only one answer** to each question.
- Please read the questions carefully before answering. Your answers will be kept confidential.

PART 1 DEMOGRAPHICAL INFORMATION

1. **Name:** _____ (optional)

2. **Age**

- | | | |
|----------|----------|----------|
| a) 20-25 | c) 31-35 | e) 41-45 |
| b) 26-30 | d) 36-40 | f) 45-50 |

3. **Gender**

- | | |
|---------|-----------|
| a. Male | b. Female |
|---------|-----------|

4. **Marital status**

- | | | |
|------------|---------------|-------------|
| a. Married | b. Un married | c. Divorced |
|------------|---------------|-------------|

5. **Working experience**

- | | | |
|-----------|---------------------|---------------------|
| a. 1-year | b. Less than 1-year | c. More than 1-year |
|-----------|---------------------|---------------------|

6. **Education**

- | | |
|--------------|----------------------|
| a. BSN | d. Nursing assistant |
| b. MBBS | e. RN diploma |
| c. Midwifery | |

7. **Profession**

- | | |
|-------------------------|-------------------|
| a. Nurse | d. Medical doctor |
| b. Healthcare assistant | e. Technician |
| c. Midwife | f. Therapist |

8. **Department**

- Emergency
- Gymea/ LR
- OPD
- IPD
- Private/ Semi Private
- ICU
- Paediatric Ward

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47

PART 2 HAND HYGIENE KNOWLEDGE QUESTIONNAIRE

1. Did you receive formal training in hand hygiene in the last three years?

Yes No

2. Do you routinely use an alcohol-based hand rub for hand hygiene?

Yes No

3. Which of the following is the main route of cross-transmission of potentially harmful germs between patients in a health-care facility? (tick one answer only)

- a. Health-care workers' hands when not clean
- b. Air circulating in the hospital
- c. Patients' exposure to colonised surfaces (i.e., beds, chairs, tables, floors)
- d. Sharing non-invasive objects (i.e., stethoscopes, pressure cuffs, etc.) between patients

4. What is the most frequent source of germs responsible for health care-associated infections? (tick one answer only)

- a. The hospital's water system
- b. The hospital air
- c. Germs already present on or within the patient
- d. The hospital environment (surfaces)

Which of the following hand hygiene actions prevents transmission of germs to the patient?

- e. Before touching a patient Yes No
- f. Immediately after a risk of body fluid exposure Yes No
- g. Immediately before a clean/aseptic procedure Yes No
- h. After exposure to the immediate surroundings of a patient Yes No

5. Which of the following statements on alcohol-based hand rub and handwashing with soap and water are true?

- i. After touching a patient Yes No
- j. Immediately after a risk of body fluid exposure Yes No
- k. Immediately before a clean/aseptic procedure Yes No
- l. After exposure to the immediate surroundings of a patient Yes No

6. Which of the following statements on alcohol-based hand rub and handwashing with soap and water are true?

- m. Hand rubbing is more rapid for hand cleansing than handwashing True False
- n. Hand rubbing causes skin dryness more than handwashing True False
- o. Hand rubbing is more effective against germs than handwashing True False

48 p. Handwashing and hand rubbing are recommended to be performed in sequence
49 True False

50
51 7. What is the minimal time needed for alcohol-based handrub to kill most germs on
52 your hands? (tick one answer only)

53 q. 20 seconds

54 r. 3 seconds

55 s. 30 seconds

56 t. 10 seconds

57
58 8. Which type of hand hygiene method is required in the following situations?

59 u. Before palpation of the abdomen Rubbing Washing None

60 v. Before giving an injection Rubbing Washing None

61 w. After emptying a bedpan Rubbing Washing None

62 x. After removing examination gloves Rubbing Washing None

63 y. After making a patient's bed Rubbing Washing None

64 z. After visible exposure to blood Rubbing Washing None

65
66
67
68 9. Which of the following should be avoided, as associated with increased likelihood
69 of colonisation of hands with harmful germs?

70 aa. Wearing jewellery Yes No

71 bb. Damaged skin Yes No

72 cc. Artificial fingernails Yes No

73 dd. Regular use of a hand cream Yes No

74

75

76
77
78
79
80 **Thank you very much for your time!**

81

82

Appendix B

83

INFORMED CONSENT FORM

84

85

86

This is Mehwish Aqeel student at Memon College of Nursing conducting the study for the requirement of Bachelor of Science of Nursing Degree. You are invited to participate in this study

87

88

89

By signing in in this form you have voluntarily agreed to participate in a research study entitled: "Assess the Knowledge, Attitude and Practice of Healthcare Workers about Hand Hygiene"

90

91

92

To be under supervision of Principal Investigator.

93

94

The purpose of the research study on topic "Assess the Knowledge, Attitude and Practice of Healthcare Workers about Hand Hygiene" is to determine the hand hygiene compliance of healthcare team.

95

96

You will be requested to fill the questionnaire consist of demographic information and hand hygiene knowledge

97

98

There will be no risk.

99

100

The study research records will be kept confidential and you will not be identified in any verbal or written reports the research records related to this study will be kept in a secure password protected computer file.

101

102

You will not be charged for procedures performed that are purely related to your participation in this study.

103

104

Your only 1hour will be needed in this participation and for any information you can call the supervisor of this study.

105

106

Your participation in this study is voluntary. You may be a participant in it only for your wish and you can withdraw from this study at any time.

107

108

109

Name of Participant: _____ **Signature** _____

110

111

112

Researcher Name: XXXXXXXXXX **Signature** _____

113

Email id: XXXXXXXX@gmail.com