

Original Research Article

ISOLATION AND CHARACTERIZATION OF ANTIBIOTIC PRODUCING FUNGI FROM SOIL

ABSTRACT

This study was aimed to isolate and characterize antibiotic producing fungi from the soil environment within Ahmadu Bello University main campus Samaru, Zaria. Soil samples were collected from five different locations within Ahmadu Bello University main campus for the Isolation of fungi. Spread plate method involving serial dilution technique was used for the Isolation using Saboraud Dextrose Agar (SDA). Six species of fungi were isolated from the soil samples and then characterized microscopically and macroscopically. The fungi were *Aspergillus niger*, *Aspergillus fumigatus*, *Aspergillus* sp, *Penicillium* sp, *Fusarium* sp (P14) and *Fusarium* sp (P15). Sensitivity test using *Staphylococcus aureus*, *Escherichia coli* and *Klebsiella pneumoniae* as test pathogens was employed to determine the ability of the fungal isolates to produce antimicrobials. All the fungal isolates were found to inhibit the growth of at least one of the test pathogens except *Fusarium* sp (P14). *Aspergillus niger* produces zones of inhibition of 9mm, 5mm and 6mm against *Staphylococcus aureus*, *Escherichia coli*, and *Klebsiella pneumoniae*, *Aspergillus fumigatus* produces zones of inhibition of 5mm against *Escherichia coli* and *Klebsiella pneumoniae*. *Aspergillus* sp produced zones of 6mm against *Staphylococcus aureus* and *Escherichia coli*, *Penicillium* sp which produces a zone of 10mm, 7mm and 6mm against *Staphylococcus aureus*, *Escherichia coli* and *Klebsiella pneumoniae*. The results of this investigation demonstrate that strains of antibiotic-producing fungi are present in the soil of Ahmadu Bello University main campus Samaru, Zaria and these could be harnessed by the pharmaceutical industries for the production of antibiotics from local sources.

Comment [H1]: Kindly add a mini-introduction before aim. A one sentence introduction.

Comment [H2]: Please put all genus and species in italics

Comment [H3]: Same here

1.0 INTRODUCTION

Comment [H4]: Please kindly write according to the journal standard.

1.1 Background of Study

Soil is a reservoir for a large number of microorganisms such as bacteria, viruses, fungi and protozoa (Raja *et al.*, 2017). Microbes of soil increase the soil fertility, maintain ecosystem sustainability, antibiotic and enzyme production and biodegradation are the distinct beneficial effects (Varnam and Evans, 2000).

Fungi are geographically widely distributed and have been observed in a broad range of habitats principally in soil (Magnuson and Lasure, 200).

The term 'antibiotic' literally means 'against life'. According to Talaro and Talaro (2002) antibiotics are substances produced by natural metabolic processes of some microorganisms that can inhibit or destroy other microorganism. Duta (2005) defined antibiotics as the miracle drugs of modern times that act as magic bullet shooting down infective organisms that have invaded the human body and caused infections.

In 1928, the first clinical trial of penicillin was carried out on humans by an English Bacteriologist Late Sir Alexander Fleming. This antibiotic was obtained from a blue green mould of the soil called *Penicillium notatum* (Dutta, 2005).

Subsequent to the identification of penicillin production by *Penicillium notatum*, screening experiments revealed that *Penicillium chrysogenum* was a superior producer of penicillin. A typical fermentation yields three types of penicillin namely, penicillin F, penicillin G and penicillin V (Denyer *et al.*, 2004; Schlegel, 2003; Talaro *et al.*, 2002).

As more antibiotics were discovered, designed and studied; scientists found that they had different properties. Some of these properties include their source, range of activity and their kinds. These were used to classify them (Sommer *et al.*, 2006). The ability to produce antibiotics

has been found mainly in fungi of the group Aspergillales, and in a few other bacteria (Schlegel, 2003).

Perhaps one of the few most important discoveries regarding the beneficial use of fungi for human was the identification in 1928 by sir Alexander Fleming, that an isolate of *Penicillium notatum* produced a substance capable of killing gram positive bacteria (Dutta, 2005; Walsh, 2003; Krishna, 2006). This compound was the first member of the E- lactam class of antibiotics to be discovered. These compounds function by inhibiting peptidoglycan synthesis in bacteria and their use has reduced the importance of the gram positive as cause of disease (Berg et al., 2003; Nester et al., 2004).

Comment [H5]: Put all et al in italics. Check other pages for same correction

A number of antibiotic drugs have been discovered from soil- inhabiting microorganisms which include Fungi (20% of isolated antibiotics), Actinomycetes (70%) and Eubacteria (10%) (Bredy, 1974; Lechevalier, 1975).

A number of Aspergillus-related patents have been issued for medicinal compounds. Lovastatin produced by *Aspergillus terreus* was one of the first commercially cholesterol-lowering drugs lam, 1983. A number of antibiotics, antitumor and antifungal agents have been derived from Aspergillus metabolites. The experimental anti-Candida drug, Cilofungin, is a semi-synthetic drug, derived from a chemical modification of echinocandin B which is produced by *Emericella nidulans/ rogulosa* (Huang et al., 1990).

Aspergilli are known for their ability to secrete a variety of biologically chemical compounds including antibiotics, mycotoxins, immune-suppressants, and cholesterol- lowering agents (Goldman and Osmani, 2008).

1.2 Statement of Research Problem

Treatment of infectious diseases caused by pathogenic bacteria was one of the most traditional problems in the clinical field (Mehrgan *et al.*, 2008). In addition, antimicrobial resistance presents as major challenge in the effective treatment of infectious disease, resulting in increased patient morbidity and mortality (O'Neill, 2014).

1.3 Justification of the Study

As more and more bacteria continue to develop resistance to currently produced antibiotics, research and development of new antibiotics continue to be important. The discovery of penicillin in 1928 by Alexander Fleming, showed its efficacy in laboratory cultures against many diseases producing bacteria. This discovery marked the beginning of the development of antibacterial compounds produced by living organism (Schlegel, 2003; Sommer *et al.*, 2006)

This research will help to reveal if strains of antibiotic-producing fungi is present in the soil that could be harnessed by pharmaceutical industry for modification to produce new high-yielding strains for the production of new antibiotics.

1.3 Aim of Study

The aim was to isolate and characterize antibiotic-producing fungi from the soil within Ahmadu Bello University main campus Samaru, Zaria.

1.4 Objectives

1. to determine the physico-chemical properties of the soil samples.
2. to isolate and Identification of fungi from soil samples.
3. To assess the antibacterial activity of the fungal isolate metabolites against *Escherichia coli*, *Klebsiella pneumoniae* and *Staphylococcus aureus*.

Comment [H6]: Please write according to the journal's standard. Go to the author's guideline or journal archives and download a published work, then follow the format.

2.0 LITERATURE REVIEW

2.1 General Characteristics of Fungi

Fungi are eukaryote with a nuclear membrane and cell walls. As heterotrophs, they depend on living or dead organic materials for both their carbon and their energy. Fungi are aerobic organisms, although, some can tolerate the rather low oxygen concentrations and high levels of carbon dioxide found in wet or compacted soils. Strictly speaking, fungi are not entirely microscopic, since some of these organisms such as mushrooms, form macroscopic structures that can easily be seen without magnification (Brady and weil, 1999).

The fungi constitute a group of living organisms devoid of chlorophyll. They resemble green plants in that, with few exceptions, they have definite cell walls, they are usually non-motile, although they may have motile reproductive cells, and they reproduce by means of spores. They do not possess stems, roots, or leaves, nor have they developed a vascular system, as have the more advanced types of plants. Fungi are usually filamentous and multi-cellular; their nuclei can be demonstrated with relative ease (Alexopoulos and Mims, 1979).

The kingdom Fungi is enormous, the identified species and those not yet classified are up to over 300,000 species. The majority of these species are Microscopic fungi (yeast and molds) often used for the production of antibiotics. A relatively small number of species have reproductive systems known as "mush rooms" that can be easily observed in the field, fungi of various shapes and colours can be observed. Examples of common shapes and appearances are the saddle or honey comb shapes, corals shape appearance, umbrella shaped (with a cap and a stipe). Similar variation exists in the colour, taste and smell of fungi. However, such macroscopic observations are not sufficient to achieve a proper scientific identification and classification of fungi and consequently, microscopic studies are also necessary (Sarad *et al.*, 2016).

Fungi belong to their own kingdom (Kingdom fungi). They obtain their nutrition through a range of ways including degradation of organic material and symbiosis (as lichen) among others (Carris *et al.*, 2012). As such, they are categorized as heterotrophic because they are unable to synthesize their own food (they lack chlorophyll). They can reproduce sexually or asexually with a majority of fungi being spore producers. Examples include Moulds, yeast, mushrooms to larger multicellular capable of forming hyphae threads of false roots. For this reason, fungi are classified according to their morphologies. Fungi are very important microorganisms because of their variety of uses which include mycorrhizal relationship between some of the fungi and plants that benefits various plants in many ways. For instance, whereas the fungi obtain nutrients from the plants, the fungi may make various materials such as phosphates readily available for the plant. Some others help in degradation of organic materials (Smith, 2008).

2.1.1 Occurrence of Soil Fungi

Soil fungi comprise of an extremely diverse group of micro-organisms. Tens of thousands of species have been identified in soils, representing some 170 genera. Although their numbers are usually somewhat smaller than bacterial, their relatively large sizes result in their dominating the biomass and metabolic activities in many soils. Their biomass is commonly ranges from 1,000 to 15,000kg/ha in the upper 15cm. (Brady and Weil, 1999).

Surprisingly, fungi may be the most abundant eukaryotic members of some deep-sea sediment (Edgcomb *et al.*, 2011). A key question is the extent to which these taxa are active to these hostile environments, as opposed to surviving in highly resistant, dominant spore stages following introduction by wind or by other vectors (Bridge and Spooner, 2012).

The most commonly measured characteristics of soil fungi are their biomass, elemental stoichiometry, growth rates and efficiencies and activities of their extracellular enzymes.

(Sinsabaugh and Follstadshah, 2011; Sinsabaugh *et al.*, 2013; Wallander *et al.*, 2013). These parameters define the size of fungal C and nutrient pools and rates of fungal substrates by enzymatically degrading complex organic matter outside the cell. However, control of the of the organisms. Hence, a substantial fraction of the enzymatic potential of soils is a legacy of enzymes that are spatially and temporally displaced from their origins (Burns *et al.*, 2013). In some cases, particularly for oxidative enzymes, these enzymes catalyze degradation and condensation reactions secondary to their original functions (Sinsabaugh, 2010).

Soil-inhabiting fungi are an important and wide spread component of most terrestrial ecosystems and play a key role in micro-ecosystem (Jackson *et al.*, 2000). Fungi have profound influences on biogeochemical cycles through their growth habits, which include external digestion of food resources using a powerful arsenal of degradative enzymes and secondary metabolism. (Floudas *et al.*, 2012). Fungi are prominent and present in all soils. At broad phylogenetic scale, the aphorism “everything is everywhere” does seem to apply to fungi; beyond soils, they are also found in nearly every other habitat on earth, including deep sea hydrothermal vents and sediments, sub glacial sediments, ancient permafrost, sea ice, hot and cold deserts, and soils of dry valleys of antarctica (Cantrell *et al.*, 2011).

Fungal biomass varies widely within and across biomes in relation to litter composition, root density, and nutrient availability. Fungi may comprise up to 20% of the mass of decomposing plant litter. Although fungal abundance and ratios of fungal to bacterial biomass tends to increase as soil pH decreases (Hogberg *et al.*, 2007; Jorgensen and Wichem, 2008). Other studies suggest that fungal distributions are more influenced by nitrogen and phosphorous than pH (Fierer *et al.*, 2009; Lauber *et al.*, 2009).

2.1.2 Structure of Fungi

The body or vegetative structure of a fungus is called a thallus (pl., thalli). It varies in complexity and size, ranging from the single-cell microscopic yeasts to multi-cellular molds, macroscopic puff balls, and mushrooms. The fungal cell usually is encased in a cell wall of chitin. Chitin is a strong but nitrogen-containing polysaccharide consisting of N-acetyl glucosamine residues (Talaro and Talaro, 2002).

2.1.3 Division of Soil Fungi

Fungi may be divided into three groups which are yeast, mushrooms and moulds. Yeasts which are single-celled organisms live principally in water logged, anaerobic soils. Molds and mushroom are both considered to be filamentous fungi, because they are characterized by long, thread-like, branching chains of cells. Individual group of filaments called hypha are often twisted together to form mycelia that appear somewhat like woven ropes. Fungal mycelia are often visible as thin, white or colored strands running through decaying plant litter. The filamentous fungi reproduce by means of spores, often formed on fruiting bodies which may be microscopic (i.e. molds) or macroscopic (i.e. mushrooms) Brady and Weil (1999).

The molds are distinctively filamentous, microscopic or semi-macroscopic fungi that play a much more important role in soil organic matter breakdown than the mushroom fungi. Molds develop vigorously in acid, neutral, or alkaline soils. Some are favored rather than harmed, by lower pH. Consequently, they may dominate the micro-flora in acid surface soils, where bacteria and actinomycetes offer only mild competition. The ability of molds to tolerate low pH is especially important in decomposing organic residues in acid forest. Many genera of molds are found in soils. Four of the most common are *Penicillium*, *Mucor*, *Fusarium*, and *Aspergillus*.

Species from these genera occur in most soils. Soil conditions determine which species are dominant (Brady and Weil, 1999).

2.1.4 Activities of Fungi

As decomposers of organic materials in soil, fungi are the most versatile and persistent of any group. They usually dominate in the upper horizons of forested soils, as well as in very acid or sandy soils. They carry out the largest share of the decomposition in many cultivated soils as well. Fungi become more efficient than bacteria that they assimilate into their tissues a larger proportion of the organic materials they metabolize. Soil fungi can synthesize a wide range of complex organic compounds in addition to those associated with soil humus. It was from a soil fungus, *Penicillium* species, that the first modern antibiotic drug, penicillin, was obtained. By killing bacteria, such compounds probably help fungi to out compete rival microorganisms in the soil (Brady and Weil, 1999).

2.1.5 Importance of Fungi to Man

The human use of fungi for food preparation or preservation and other purposes is extensive and has a long history. The study of the historical uses and sociological impact of fungi is known as ethno mycology. Because of the capacity of this group to produce enormous range of natural products with antimicrobial or other biological activities, many species have long been used or are being developed for industrial production of antibiotics, vitamins, anti-cancer and cholesterol-lowering drugs. More recently, methods have been developed for the genetic engineering of fungi, enabling metabolic engineering of fungal species. For example, genetic modification of yeast species (Holkins and smolke, 2008) which are easy to grow at fast rates in large fermentation vessels has opened up ways of pharmaceutical production that are potentially more efficient than production by the original source organisms (Huang *et al.*, 2008).

2.2 The Genus *Penicillium*

2.2.1 Occurrence

The *Penicillium* species are as common and cosmopolitan as the *Aspergilli*. They are the so-called green molds and blue molds which are so frequently found on citrus and other fruits, and on other food stuffs that have become contaminated with their spores. The conidia of *Penicillium*, like those of *Aspergillus*, are everywhere in the air and in the soil (Alexopoulos and Mims, 1979).

2.2.2 Economic Importance

Many species of *Penicillium* are capable of producing organic acids, such as citric, fumaric, oxalic, gluconic, and gallic. Industrially, the penicillia are important in making cheese and antibiotics. *Penicillium notatum* and *Penicillium chrysogenum* have come to prominence as sources of the now famous antibiotic, Penicillin. Although other species of *Penicillium* also produce this antibiotic, certain strains of the latter species have proved to be most efficient in its manufacture and are being used exclusively in the commercial preparation of this drug. Of great interest in this connection is the ability of mycologists to produce new high-yielding strains of *Penicillium chrysogenum* by subjecting conidia to ultra-violet irradiation and testing the colonies originating from the conidia for Penicillin production. Yields of penicillin have been increased enormously in this way, and this has been the primary factor in the price reduction which has made this first "wonder drug" available to all (Alexopoulos and Mims, 1979).

2.2.3 Morphology

The life history of a typical *Penicillium* is very much like that of *Aspergillus*, but the morphology of the structures differs considerably. The mycelium produces simple, long erect conidiophores which branch about two thirds of the way to the tip, in symmetrical and asymmetrical, broom-

like fashion. The conidiophores commonly referred to as the brush, is technically known as the penicillus (pl. *Penicilli*; I; *Penicillium* = small brush). The conidia are globule to ovoid, and under the microscope resemble glass beads. They are formed in the same manner as in *Aspergillus*. The enormous quantities of greenish, bluish, or yellow conidia which are produced are chiefly responsible for the characteristic colony colour of various species of *Penicillium*. The conidia germinate easily into germ tubes from which mycelia develop (Alexopoulos and Mims, 1979).

2.3 The Genus *Aspergillus*

2.3.1 Occurrence

Aspergillus is a genus of anamorphic fungi reproducing by production of phialospores (conidia borne on phialides). It is a large genus with over 180 recognized species (Pitt *et al.*, 2000).

Thom and Raper (1945) recognized 78 species of *Aspergillus*. The black aspergilli or *Aspergillus niger* group are the fungi which we commonly call black mold. The genus is widely distributed from the arctic region to the tropics. The air everywhere seems to contain the conidia of these organisms. The soil, too, contains the spores of the aspergilli. The aspergilli are capable of utilizing an enormous variety of substances for food because of the large numbers of enzymes they produce. Indeed, it is difficult to find a substance containing some organic matter and a little moisture on which the aspergilli cannot grow (Alexopoulos and Mims, 1979).

2.3.2 Economic Importance

Because of their great enzymatic activities, aspergilli are employed in several industrial processes. Citric and gluconic acid are manufactured commercially by the use of *Aspergillus niger*. Many other acids and various other chemicals are produced in large or small quantities by members of this genus. Enzyme preparations are made commercially through the use of these

fungi, and a number of antibiotics have been isolated from aspergillus cultures, though none has proved equal to Penicillin or to the actinomycete product in therapeutic properties. In Japan, *Aspergillus oryzae* is used to make sake, an alcoholic beverage concocted from rice, and to manufacture various fermented foods. In Java, *aspergillus wentii* is employed in processing soybeans because of its ability to loosen the hard tissue of the bean (Alexopoulos and Mims, 1979).

The greatest positive economic impact of the *Aspergilli* has been in the exploitation of the enzymes and acids produced a number of species. Two of the most important industrial products produced by *Aspergilli* are amylase and citric acid.' The Koji molds' (*Aspergillus oryzae*, *Aspergillus sojae* and *Aspergillus awamoni*) have been used for more than a thousand years to produce a number of a mold strain grown on steamed grain. The mold produce amylases which break down the starches and contribute to the flavour and colour of the product (Hara *et al.*, 1992). Citric acid production by *Aspergillus niger* was discovered in 1916. By the mid-1920's over three quarters of the citric acid, used worldwide was produced by fungal fermentation. It is still used today to produce more than 500,000 tons per year (Roehr *et al.*, 1992)

Unfortunately, the positive economic impact of *Aspergilli* and their metabolites are more than balanced by the negative aspects. *Aspergilli* are a major cause of degradation of agricultural products both before and after harvest. Some species are human and animal pathogens, and a number of species are allergenic. They also produce variety of mycotoxins which are secondary metabolites that are harmful to humans and animals (Maren, 2002). Mycotoxins are products of secondary metabolism indicating that their production is not necessary for the survival of the fungus. Unlike antibiotics which are toxic only to microorganisms, mycotoxins are harmful to humans and or animals. Aflatoxin is the most economically important mycotoxin in the world.

The most toxic form of aflatoxin is aflatoxin B. Aflatoxin is formed in oil seed crops (cotton seed, corn and peanuts) and tree nuts under drought conditions in the field. It can form in virtually any stored grain that can support fungal growth (Diener *et al.*, 1987).

2.3.3 Somatic Structures of *Aspergilli*

The mycellium of *Aspergilli* resembles that of other fungi. The hyphae are well developed, profusely branched, septate, and hyaline; their cells are, as a rule, multinucleate (Alexopoulos and Mims, 1979).

Aspergillus is characterized by its distinctive conidiospore. The base of the conidiospore usually forms 'T' or 'L' shape where it connects with the vegetative hyphae. This is commonly called the 'foot cell' even though it is not a separate cell. The stipe extends from the foot cell and may be quite short (50 micro meter or less) to several millimeters in length. The Apex of the stipe expands into a vesicle. The vesicles may have various characteristic shapes. In some species, the conidia-bearing phialides arise directly from the vesicle. This form is called uniseriate. In other species, there is a second layer of cells between the vesicle and the phialides. These cells are called 'metulae' and *Aspergilli* with metulae are referred to as biseriate species. An important character that distinguishes *Aspergillus* from several closely related genera is that the phialides/metulae arise simultaneously on the vesicle (Maren, 2002).

2.4.0 Test Organisms

2.4.1 *Escherichia coli*

Escherichia coli is a gram negative, rod shaped bacteria and a typical coliform of the intestinal tract. It is important;

- (a) As a natural commensal (inhabit the intestine of humans and animals, thus belonging to the group of microorganisms known as *Enterobacteria*).

- (b) As a pathogen, particularly of the urinary tract, and also in diarrhea, appendicitis, peritonitis, neonatal meningitis and pyelitis.
- (c) In proving evidence of faecal contamination of water supplies.

It grows well in most media and it is typically indole-positive, produce gas from glucose and haemolysis is seen on blood agar. It has a complex antigenic structure (Brooks, 1998; Cheesbrough, 2006).

2.4.2 *Klebsiella pneumoniae*

Klebsiella pneumoniae is a gram negative, non-motile, encapsulated rod. it is lactose fermenting, producing mucoid pink colonies on macConkey agar, yellow mucoid colonies on CLED medium and grey white, usually mucoid colonies on blood agar. It is also urease positive, citrate positive and utilizes malonate.

It is a major pathogen of humans, present in the respiratory tract and faeces of about 5% of normal individuals. It causes a small proportion of (about 1%) of bacterial pneumonia. *Klebsiella pneumoniae* causes chest infections and occasionally severe bronchopneumonia with lung abscesses. It occasionally produces urinary tract infections and bacteremia. Infections of *Klebsiella pneumoniae* are often opportunistic occurring in patients with existing diseases or in debilitated patients (Brooks, 1998; Cheesbrough, 2006).

2.4.3 *Staphylococcus aureus*

Staphylococcus aureus are gram positive, non-sporing, non-motile cocci which forms a regular grape-like clusters. They grow rapidly and abundantly under aerobic condition. They can't operate high concentration of salt and are resistant to heat. They are catalase positive, oxidase negative and have the ability to produce coagulase (Harish *et al.*, 2002).

They cause different diseases, such as septicaemia, boils, styes, toxic food-poisoning, Staphylococcus aureus can survive on dry surfaces for months and infect its host from the skin, mouth and nose (Morgenstern *et al.*, 2016)

Comment [H7]: Please remove. Is not needed. Follow the guideline

UNDER PEER REVIEW

METHODOLOGY

Comment [H8]: remove

3.0 MATERIALS AND METHODS

3.1 Study Area

This research was carried out within Ahmadu Bello University main campus Samaru Zaria. The soil samples were gotten from the following Area/sites: Dangote hostel, Sassakawa hostel, Akenzua hostel, Danfodio hostel and Division of Agricultural College (DAC).

3.2 Collection of Samples

Soil samples was collected aseptically from five different locations within Ahmadu Bello University. Spatula cleansed with 70% alcohol was used to collect the soil samples into a transparent sterile polythene bag. The samples were then immediately taken to the Microbiology Laboratory, where they were analyzed. The five locations where the samples were collected includes: Dangote hostel, Sassakawa hostel, Akenzua hostel, Division of Agricultural College (DAC) and Danfodio hostel.

Comment [H9]: were

3.3 Determination of Soil Temperature

The temperature of the soil at the five different sites was determined by the use of thermometer. The thermometer was inserted into the soil up to depth of 5cm and allowed to stay for 10minutes, after which the temperature readings were then obtained. The average of three consecutive readings was recorded for each site (Dix and Webster, 1995).

3.4 Determination of Soil pH

The soil pH values were determined in Institute of Agricultural Research (IAR), soil analysis laboratory by pH meter using standard methods of Watson and Brown (1998). Using this method, 3g of soil sample was weighed into a beaker containing 3ml of distilled water, which was then stirred for five seconds and allowed to stand for 10 minutes. The electrode of the pH

meter was then inserted into the slurry and swirled carefully. The reading was taken thereof and the average of the consecutive readings was recorded for each site.

3.5 Soil Types

Comment [H10]: Put in bold

The soil types were determined in Institute of Agricultural Research (IAR), soil analysis laboratory (Ahmadu Bello University) according to the method of (Whitbread *et al.*, 1996; Pettijohn, 2000).

3.6 Isolation and Identification of Fungal Isolates

Comment [H11]: Put in bold

The soil fungi were isolated by Soil Dilution Techniques using the spread plate method. The Media used for the isolation was Sabouraud Dextrose Agar (SDA) which was prepared according to the manufacturer's instruction. 1g of soil sample was measured using a weighing balance and was then immersed into 9ml of distilled water in a bijou bottle. Serial dilution was then carried out in ten (10) folds in bijou bottles each containing 9 ml of distilled water to obtain 1/10 dilution (stock solution). 1ml of the stock solution was transferred into the first bijou bottle which was shaken vigorously to have a uniform distribution. Another 1ml was taken from the first diluents to the second bottle and 1ml was also taken from the second diluents to the third bottle respectively (Waksman and Fred, 1922). 0.1 Aliquot each of 10^{-3} dilution was then taken by the use of a syringe which was then dispensed into a Petri dish and the same was done for all the remaining samples. The Plates were incubated for three (3) to seven (7) days at room temperature. Pure cultures of the fungal isolates were identified using both macroscopic (cultural) and microscopic (morphological) features with reference to Atlas of Mycology.

3.7 Determination of Percentage Occurrence Frequency of Fungal Isolates

The percentage frequency of occurrence for each species of fungus isolated was determined by the methods of Sampo *et al.* (1997). This was computed by the formula: $A/B \times 100$; where A =

Number of plates in which the species appeared, and B = Total number of plates incubated for each sites.

3.8.0 Test Organisms

The test bacteria that were used for this study were *Escherichia coli*, *Klebsiella pneumoniae* and *Staphylococcus aureus*. The bacterial isolates were obtained from Ahmadu Bello University medical center, Samaru, Zaria, where it was isolated from patients. The Isolates of these organisms were taken to the laboratory of the Microbiology Department Faculty of Life Sciences, Ahmadu Bello University Zaria. The organisms were sub-cultured on Nutrient Agar for reconfirmation.

Comment [H12]: The isolates were taken to microbiology lab. at

3.8.1 Reconfirmation of *Staphylococcus aureus* Isolates

3.8.1.1 Catalase Test

A drop of hydrogen peroxide was added to the surface of a clean, grease-free glass slide and colonies of the suspected *Staphylococcus aureus* was added to it with a sterile straight wire loop and emulsified. Bubbles formation indicates positive test while absence of bubbles indicates negative result. *Staphylococcus aureus* is catalase positive (Cheesbrough, 2006).

3.8.1.2 Coagulase Test

A drop of plasma was added to already emulsify distilled water with colonies of the suspected *Staphylococcus aureus* on clean grease-free slide, and then the slide was rocked for 10 seconds. Positive test was indicated by clumping, while negative test shows no clumping. *Staphylococcus aureus* was coagulase positive (Cheesbrough, 2006).

3.8.2 Reconfirmation of *Escherichia coli* and *Klebsiella pneumoniae* Isolates

3.8.2.1 Indole Test

The indole test was carried out by adding three drops of Kovac's indole reagent to 5ml of young broth culture of the isolated bacteria in the test tube and shaken gently for a minute and positive result was indicated by a red layer at the top of the broth culture while a negative result was indicated by lack of colour change at the top of the broth culture. *Escherichia coli* and *Klebsiella pneumoniae* are indole positive (Cheesbrough, 2006)

3.8.2.2 Methyl-Red Test

Tubes containing Methyl Red-Voges Proskauer (MR-VP) broth was inoculated with the suspected isolates and incubated for 24 hours at 37°C. Three drops of Methyl-Red solution was added to each tube. After incubation, a distinct red colour indicated a positive test for *Escherichia coli* while absence of colour change indicated negative test for *Klebsiella pneumoniae* (Cheesbrough, 2006).

3.8.2.3 Voges-Proskauer Test

This test uses alpha naphthol and potassium hydroxide to test for the presence of acetyl methyl carbinol (acetoin). Tubes of MR-VP broth was inoculated with the suspected isolates and incubated for 24 hours at 37°C. 1ml of the broth was transferred into another tube, 0.6ml alpha-naphthol solution and 0.2ml 40% KOH was added and shaken, it was then allowed to stand for 30-60 minutes. A yellowish color is an indication of a negative result while absence of color change indicated a positive result. *Escherichia coli* is VP negative while *Klebsiella pneumoniae* is VP positive (Cheesbrough, 2006).

3.8.2.4 Citrate Utilization Test

A simmon citrate agar was inoculated with the suspected isolates and incubated at 37c for 24 hours after which the medium was observed for color change. Citrate positive organisms, growth will be visible on the slant surface and the medium will be an intense prussian blue. The alkaline carbonates produced as by-product of citrate catabolism raises the pH of the medium to above 7.6, causing a colour change from the original green color to blue. *Klebsiella pneumoniae* is citrate utilization positive while *Escherichiacoli* are citrate utilization negative (Chesbrough, 2006).

Comment [H13]: Write properly

Comment [H14]: Space out

3.9 Determination of Antibiotic Production by the Fungal Isolates

Methods used by Williams and Cross (1971) was employed to determine the antibiotic production potential by fungal isolates *in vitro* against common human pathogens. The human pathogens used for this study were: *Escherichia coli*, *Klebsiella pneumonia* and *Staphylococcus aureus*. Each fungal isolate was streaked on Nutrient Agar as a straight line and incubated at 30°C. After two days of incubation, the human pathogenic test organism *Escherichia coli*, *Klebsiella pneumonia* and *Staphylococcus aureus* were streaked perpendicular to the streaked line of the growing fungus. This was then incubated at 30°C for 24 hours, after which the Zone of Inhibition of each test organisms from the streaked line of the growing fungus was measured.

RESULTS AND FINDINGS

Comment [H15]: Delete

4.0 Results

An investigation was carried out in this study to determine the presence of antibiotic-producing fungi in the soil samples collected from five different sites within Ahmadu Bello University Main Campus Samaru, Zaria. The experimental result obtained in this study were presented in a tabular form. The list of tables is shown below:

Table 1: Show results of the physico-chemical Properties of soil samples gotten from the soil environment of Danfodio, Division of Agricultural College (DAC), Dangote, Akenzua and Sassakawa. The soil pH varies from 5.92 to 8.03 and the temperature range is between 26°C to 32°C respectively.

Table 2: Shows the macroscopic and microscopic characteristics of the fungal isolates. A total of six species of fungi were isolated. The macroscopic characteristics were based on colony colour, texture, elevation and reverse and also microscopic characteristics was based on their conidia, stipes, and vesicles.

Comment [H16]: Restructure the grammar to add whole meaning to the sentence.

Table 3: shows the occurrence of the fungal isolates from each soil samples. *Aspergillus niger*, *Aspergillus fumigatus* and *Fusarium* sp had occurrence of 80% each *Penicillium* sp and *Aspergillus* sp had 40%

Table 4: Show the results of the antimicrobial activities of the fungal isolates metabolites against *Staphylococcus aureus*, *Escherichia coli*, and *Klebsiella pneumonia*. The zone of inhibition produced against the test pathogens ranges from 3mm to 10mm respectively.

Comment [H17]: Justify to be on a straight line

Table 1: Physio-chemical properties of the soil samples from the different locations within Ahmadu Bello University Campus Samaru, Zaria

| Locations | Soil Types | Soil pH | Temperature (°C) |
|------------------|-------------------|----------------|-------------------------|
| DAC | Loamy | 7.33 | 30 |
| Dangote | Clay | 7.44 | 27 |
| Danfodio | Loamy | 5.92 | 26 |
| Akenzua | Sandy | 6.35 | 32 |
| Sassakawa | Clay | 8.03 | 28 |

Table 2: Macroscopic and Microscopic characteristics of the fungal Isolates

| Sample code | Macroscopic Characteristics | Microscopic Characteristics | Inference |
|-------------|---|---|------------------------------|
| P10 | <p>Colour: White at first then turns black to dark brown</p> <p>Texture: powdery</p> <p>Elevation: Flat</p> <p>Reverse: White to Yellow</p> | <p>Conidia: Globose and irregularly rough.</p> <p>Stipes: Thick, smooth, slightly brown especially near the apices</p> <p>Vesicles: Nearly spherical.</p> | <i>Aspergillus niger</i> |
| P11 | <p>Colour: Dull green colour</p> <p>Texture: Floccose</p> <p>Elevation: plane</p> <p>Reverse: Dull Yellow</p> | <p>Conidia: Globose to broadly ellipsoidal, smooth</p> <p>Stipes: Uncoloured or grayish near the apices, smooth-walled</p> <p>Vesicles: Pyriform.</p> | <i>Aspergillus fumigatus</i> |
| P12 | <p>Colour: Green Yellow</p> <p>Texture: slightly granular, wrinkled</p> <p>Elevation: Plane</p> <p>Reverse: Yellow</p> | <p>Conidia: Globose, smooth walled</p> <p>Stipes: Un coloured to pale brown, walls smooth to slightly roughed</p> <p>Vesicles: Subglobose to pyriform.</p> | <i>Aspergillus sp</i> |
| P13 | <p>Colour: Blue green</p> <p>Texture: powdery</p> <p>Elevation: plane</p> <p>Reverse: White to Yellowish</p> | <p>Mycelium produces simple long erect conidiophores which branch about two-third of the way with phialides in brush- like clusters.</p> | <i>Penicillium sp</i> |
| P14 | <p>Colour: White</p> <p>Texture: Cottony</p> | <p>Hyphae slender and tapering with no chlamydospore</p> | <i>Fusarium sp</i> |

Comment [H18]: Don't put sp in italics.

Elevation: Raised

Reverse: Yellowish brown

P15

Colour: Red

Texture: Cottony

Elevation: Raised

Reverse: Red brown

Hyphae slender, longer and tapering (septate) with swollen, thick-walled chlamydospore

Fusarium
sp

UNDER PEER REVIEW

Table 3 Occurrence of fungal isolates gotten from each soil samples

| | DAC | DANG | DANF | AKEN | SASS | OCCURRENCE(%) |
|------------------------------|-----|------|------|------|------|---------------|
| <i>Aspergillus niger</i> | + | + | - | + | + | 80 |
| <i>Aspergillus fumigatus</i> | - | + | + | + | + | 80 |
| <i>Aspergillus</i> sp | + | - | - | + | - | 40 |
| <i>Penicillium</i> sp | + | - | + | - | - | 40 |
| <i>Fusarium</i> sp | + | + | + | - | + | 80 |
| <i>Fusarium</i> sp | + | - | - | - | - | 20 |

KEY

- = Absent

+ = Present

DAC= Division of Agricultural College, DANG= Dangote, DANF= Danfodio, AKEN= Akenzua, SASS= Sassakawa

Table 4 Antibacterial activity of the fungal isolates against the test bacteria

| Sample code | Isolate Identity | Zones of Inhibition (mm) | | |
|-------------|------------------------------|--------------------------|----------------|----------------------|
| | | <i>S. aureus</i> | <i>E. coli</i> | <i>K. pneumoniae</i> |
| P10 | <i>Aspergillus niger</i> | 9 | 5 | 6 |
| P11 | <i>Aspergillus fumigates</i> | NA | 5 | 5 |
| P12 | <i>Aspergillus sp</i> | 6 | 6 | 4 |
| P13 | <i>Penicillium sp</i> | 10 | 7 | 6 |
| P14 | <i>Fusarium sp</i> | NA | NA | NA |
| P15 | <i>Fusarium sp</i> | NA | 3 | NA |

KEY

NA= No Activity

5.0 Discussion, Conclusion and Recommendation

Comment [H19]: remove

5.1 Discussion

Comment [H20]: Please kindly discuss your findings in line with other scholar's findings.

The soil pH, is therefore, described as the "master soil variables" that influences myriads of soil biological, chemical and physical properties and processes that affect plant growth and biomass yield (Minasny *et al.*, 2016).

In this study, the pH values of the soil samples ranges from 5.92 to 8.03. Danfodio had a pH of 5.92 and Sassakawa had pH of 8.03.

The variation in soil pH might be due to the activities carried out by microorganisms in the soil. This is in agreement with recent studies done by (Neina, 2020).

The temperature of the soil as at the time of collection (rainy season) revealed that the soil environment had a temperature range of 26°C to 32°C.

Aspergillus niger, *Aspergillus fumigatus* and *Fusarium* sp (P14) had the highest occurrence of 80% each. *Fusarium* sp (P15) had the least occurrence of 20%.

The high occurrence might be because the soil condition was favourable for the fungal growth and also because hyphae forming microorganisms (fungi) are best adapted to benefit from increased nutrient concentration in soil. The hyphae of fungi are able to colonize discrete pieces of organic matter very rapidly and are able to form bridges between nutrient rich areas (Varnam *et al.*, 2000).

Penicillium sp and *Aspergillus niger* produce the highest zone of Inhibition of 10mm and 9mm and *Fusarium* sp (P15) produced the least zone of Inhibition of 3mm.

The zone of inhibition produced might be as a result of some bio-active compounds or metabolites produced by the fungal isolates due to limiting nutrients. The antibiotic production is of importance in enhancing the ability to compete for nutrient (Varnam *et al.*, 2000).

Comment [H21]: Justify too

5.2 Conclusion

The soil sample from each sites was found to be vary from slightly acidic to slightly alkaline with pH ranging from 5.92 to 8.03. The soil temperature ranges from 26°C to 32°C.

Comment [H22]: were

A total of six fungi were isolated from the soil and they were identified based on their macroscopic and microscopic characteristics. They were *Aspergillus niger*, *Aspergillus fumigatus*, *Aspergillus* sp, *Penicillium* sp, *Fusarium* sp (P14) and *Fusarium* sp (P15).

Aspergillus niger, *Aspergillus fumigatus* and *Fusarium* sp (P14) had the highest occurrence of 80% and *Fusarium* sp (P15) had the least occurrence of 20%.

Comment [H23]: Please justify the write up so as to be on a straight line. Highlight and press control J

All the fungal isolates metabolites inhibited at least one of the test organisms except *Fusarium* sp (P14). *Penicillium* sp and *Aspergillus niger* metabolites produced the highest zone of inhibition of 10mm and 9mm against *Staphylococcus aureus*.

This investigation reveals that some Fungal species isolated from the soil environment do produce some form of antimicrobials.

5.3 Recommendation

1. The soil harbors numerous kinds of microorganisms such as fungi and so activities involving the use of soil should be done appropriately or carefully.
2. Fungi are said to be both beneficial and harmful and so care should be taken while culturing them in the laboratory as some are said to be sporogenous (spore formers) which could be inhaled.
3. In the determination of antibiotic producing fungi, further methods should be used such as aqueous extract of the fungal isolates.

4. This investigation is a primary study; further investigation needs to be embarked upon to determine the type of antimicrobial substance(s) produced or the type of effect they cause on the pathogens whether static or cidal.

UNDER PEER REVIEW

REFERENCES

- Alexopoulos, C.J. and Mims, C.W. (1979). *Introductory Mycology*. 3rd edition. John Wiley and sons, Inc., New York, London, Sydney. 271-281.
- Baumgardner, D.J. (2012). Soil-related bacterial and fungal infections. *Journal of the American Board of Family Medicine Practice*, **25**(5), 734-744.
- Berg, J.M., Tymolko, J.L. and Stryer, L. (2002). *Biochemistry*, 5th edition. W.H Freeman and Company, New York.
- Brady, N.C. and Weil, R.R. (1999). *The nature and properties of soils*. 12th edition. Prentice Hall international Upper Saddle River, New Jersey 07458, 425-428. ISBN 0-13-852444-0.
- Bredy, J. (1974). Recent developments of antibiotic research classification of antibiotics according to chemical structures. *Advanced Applied Microbiology* **18**: 309-406.
- Bridge, P., Spooner, B. (2012). Non-Lichenized Antarctic Fungi: transient visitors or members of a cryptic ecosystem? *Fungal Ecology*, **5**: 381- 394.
- Brooks, G.F., Butel, J.s., Morse, S.A., Jawetz, Melnick and Adelberg. (2004). *Medical Microbiology*, 23rd edition. McGraw Hill Companies, Singapore.
- Burns, R., Deforest, J., Marxsen, J., Sinsabaugh, R., Stromberger, M., Wallenstein, M., Weintraub, M., Zoppini, A. (2013). Soil enzyme research: current knowledge and future directions. *Soil Biology and Biochemistry*, **58**: 216 - 234.
- Cantrell, S.A., Dianese, J.C., Fell, J., Gunde-Cimerman, N., Zalar, P. (2011). Unusual fungal niches. *Mycologist*, **103**: 1161-1174.
- Carris, L.M., Little, C.R. and Stiles, C.M. (2012). *Introduction to Fungi*. The Plant Health Instructor. DOI: 10.1094/PHI-1-2012-0426011.
- Cheesbrough, M. (2006). *District Laboratory Practice in Tropical Countries*, 5th edition. Cambridge University Press, Cambridge.
- Denyer, S.P., Hodges, N.A. and German, S.P. (2004). *Hugo and Russell's Pharmaceutical*, 7th edition. *Blackwell Science*, India.
- Dix, N.J. and Webster, J.W. (1998). *Fungal Ecology*. Chapman and Hall, London.
- Dobashi, K., Matsuda, N., Hamada, M., Naganawa, H., Takita, T. and Takeuchi, T., Novel (1998). Antifungal antibiotics octacosamicins A and B: Taxonomy, fermentation and isolation, physicochemical properties and biological activities. *Journal of Antibiotics*. **41**: 1525-1532.
- Domsch, K.H., Gams, W. and Anderson, T. (1980). *Compendium of Soil Fungi*. Academic Press, London.
- Dutta, A.C. (2005). *Botany for Degree Students* 18th edition. Oxford University Press, New York.
- Edgcomb, V.P., Beaudoin, D., Fast, R., Biddle, J.F., Teske, A. (2011). Marine Subsurface Eukaryotes: the fungal majority. *Environmental Microbiology*, **13**: 172-183

Comment [H24]: Kindly add citations that are not here. Also, remove references that were not cited.

- Fierer, N., Strickland, M.S., Liptzin, D., Bradford, M.A and Cleveland, C.C. (2009). Global patterns in below ground communities. *Ecological Letter*, **12**: 1238-1249.
- Floudas, D., Blinder, M., Riley, R., Barry, K., Blanchetle, R.A., Henrissat, B., Martinez, A.T., Otilar R., Spatafora J.W and Yadav J.S. (2012). The Paleozoic origin of enzymatic lignin decomposition reconstructed from 31 fungal genomes. *Science*, **336**: 1715-1719.
- Goldman, G.H. and Osmani, S.A. (2008). The aspergilli. CRC press.
- Harris, I.G, Foster, S.J, and Richard, R.G. (2002). An introduction to *Staphylococcus aureus* and Techniques for identifying and Quantifying *Staphylococcus aureus* Adhesion in Relation to Adhesion to Biomaterials. *Journal of European cells and Materials* **4**: 39-60. ISSN: 14732262.
- Hawkins, K.M and Smolke, C.D. (2008). Production of benzyloquinoline alkaloids in *Saccharomyces cerevisiae*. *Nature Chemical Biology*, **4**(9): 564-73
- Hawksworth, D.L. and Licking, R. (2017). Fungal Diversity Revisited 2.2 to 3.8 Million Species. *Microbiology spectrum* **5**: 79-95.
- Hogberg, M.N., Hogberg, P. and Myrold, D.D. (2007). Microbial community composition in Boreal forest soils determined by pH, C-to-N ratio, the trees, or all three? *Ecology*, **150**: 590-601.
- Huang, B., Guo, J., To, B., Yu, X., Sun, L. and Chen, W. (2008). Heterologous production of secondary metabolites as pharmaceuticals in *Saccharomyces cerevisiae*. *Biotechnology Letter*, **30**(7): 1121-1137.
- Jorgensen, R.G. and Wichern, F. (2008). Quantitative assessment of the fungal contribution to microbial tissue in soil. *Soil Biology and Biochemistry*, **40**: 2977-2991.
- Krishna, V.S. (2006). Biopharmaceuticals II, 1st edition. New Age International publishers, New York.
- Lauber, C.L., Hamady, M., Knight, R. and Fierer, N. (2009). Pyrosequencing-based assessment of soil pH as a predictor of soil bacterial community structure at the continental scale. *Applications*, **75**: 5111.
- Lechevalier, H. (1975). Production of some antibiotics by members of different genera of microorganisms. *Advanced Applied Microbiology*, **19**: 25-45.
- Magnuson, J.k. and Lasure, L.L. (2002). Fungal diversity in soils as assessed by direct culture & molecular techniques, in 102nd *General Meeting of the American Society for Microbiology*, Salk Lake City, 19-23.
- Maren, A.K. (2002). Identification of common *Aspergillus* species, 1-2.
- Morgenstern, M., Erichsen, C., Von R., and Stephen, H. (2016). *Staphylococcal* orthopedic device- related infections in older patients. *Injury*, **47**(7): 10.
- Neina, D. (1999). The Role of Soil pH in Plant Nutrition and Soil Remediation. *Applied and Environmental Soil Science*, 2019: 1-9.
- Nester, E.W., Anderson, D.G., Roberts, C.E., Pearsall N.N. and Nester M.T. (2004). *Microbiology. A Human perspective*, 4th edition. McGraw Hill, New York.

- O'Neill, J. (2014). Antimicrobial resistance: tackling a crisis for health and wealth of nations. *Review Antimicrobials*. Resistance 16
- Raja, M., praveena, G., and Williams, S. (2017). Isolation and identification of fungi from soil in loyola college campus, chennai, India. *Internation Journal of Current Microbiology and Applied sciences*, volume **6**(2):1789-1795.
- Sampo, S., Begero, R., Buffa, G. and Lumpimosa, A.M. (1997). *Soil Fungi*. Academic Press, London.
- Schlegel, H.G. (2003) *General Microbiology*, 7th edition. Cambridge University Press, Cambridge.
- Sinsabaugh, R.L. (2010). Phenol oxidase, peroxidase and organic matter Dynamics of soil. *Soil Biology and Biochemistry*, **49**: 391- 404.
- Sinsabaugh, R.L. and Follstadshah J.J. (2011). Eco-enzymatic stoichiometry of recalcitrant organic matter decomposition: the growth rate hypothesis in reverse. *Biogeochemistry*, **102**: 31- 43
- Sinsabaugh, R.L., Manzoni, S., Moorhead, D.L. and Richter, A. (2013). Carbon use efficiency of microbial communities. *Stoichiometry, methodology and modelling*, **16**: 930- 939.
- Smith, D. (2008). Mycorrhizal symbiosis. The symbionts forming arbuscular mycorrhizas, 13-41.
- Sommer, C.V, Shapp M.G, Gerald F.C, Feder B, and Martin L.A. (2006). *The New Book of Knowledge*, 306-312.
- Talaro, K. and Talaro (2002) A., *Foundations in Microbiology*, 4th edition. McGraw Hill, New York.
- Varnam, A.H and Evans M.G. (2000). *Environmental Microbiology*. University of North London, U.K. ISBN 1-55581-218-X.
- Waksman, S.A., Fred, E.D. (1992). A tentative outline of the plate method for determining the number of microorganisms in the soil. *Journal of Soil Science*. **14**(1):27-28.
- Wallander, H., Ekblad, A., Godbold, D., Johnson, D., Bahr, A., Baldrian, P., Bjork, R., Kieliszewska-Rokicka, B., Kjoller, R. and Kraigher, H. (2013). Evaluation of methodsto estimate production, biomass and turnover of ectomycorrhizal mycellium in forest soils. *Soil Biology and Biochemistry*, **57**: 1034- 1047.
- Watson, M.E. and Brown, J.R. (1998). pH and Lime requirements. *Soil Science Society of American Journal*. **221**:13-16.
- Whitbread, A.M., Lefroy, R.D. and Blair, G.J. (1996). Changes in physical properties and soil organic fractions with a cropping on a red brown earth soil. *Australian Journal of Agro*, **12**: 23-29.
- Williams, S.T. and Cross, T. *Actinomycetes*. In: J.R. Norris, D. W Robbins (1971). *Methods in Microbiology*. Academic Press, New York, 29334.