

# CHARACTERIZATION AND IDENTIFICATION OF BACTERIA PRESENT IN THE BATH TOWELS OF FEMALE STUDENTS IN RIVERS STATE UNIVERSITY HOSTELS

## Abstract

Bath towels are an important part of our everyday life, but they often amass a large number of micro-organisms which may sometimes be harmful to us. For the study a total of 10 samples plus a control were collected from bath towels of female students in NDDC hostel, post-graduate hostel and control from Mile 3 market respectively between November 2021 to February 2022. Standard methods were employed for the sampling and determination of microbiological characteristics. Identification of bacteria was carried out using colonial, morphological and biochemical characteristics. Statistical analysis were performed using the T-test method and sensitivity was carried out on the isolates to detect pathogenicity. Range of microbial counts of NDDC hostel were: Total heterotrophic bacteria  $2.0 \times 10^4$  cfu/ml to  $3.17 \times 10^4$  cfu/ml, Total coliform count  $0.48 \times 10^4$  cfu/ml to  $2.99 \times 10^4$  cfu/ml in the first sampling. Post graduate hostel ranged from: Total heterotrophic bacteria  $1.48 \times 10^4$  cfu/ml to  $0.18 \times 10^4$  cfu/ml in the first sampling. For the second sampling, NDDC hostel microbial count ranged from  $0.73 \times 10^4$  cfu/ml to  $1.15 \times 10^4$  cfu/ml in the total heterotrophic bacteria, and  $0.31 \times 10^4$  cfu/ml to  $0.13 \times 10^4$  cfu/ml for total coliform count. Post graduate hostel ranged from  $0.27 \times 10^4$  cfu/ml to  $0.93 \times 10^4$  cfu/ml for total heterotrophic bacteria and  $0.22 \times 10^4$  cfu/ml to  $0.18 \times 10^4$  cfu/ml for total coliform count. The control microbial count for total heterotrophic bacteria were  $0.12 \times 10^4$  and  $0.16 \times 10^4$  cfu/ml for the first and second sampling respectively and for the total coliform count,  $0.18 \times 10^4$  cfu/ml and  $0.17 \times 10^4$  cfu/ml for the first and second sampling respectively. Statistical analysis using student's T-test was carried out. The mean test values for total heterotrophic bacteria in the first and second samplings were 2.2750E2 and 57.4000 in NDDC hostel and 1.5060E2 and 66.8000 in post graduate hostel. The mean values for total coliform count for the first and second samplings were 1.6220E2 and 24.5000 in NDDC hostel and 99.8000 and 26.9000 in Post graduate hostel. The test revealed that there was no significant difference in the bacterial load of NDDC hostel and Post graduate hostel from the two samplings. In the antimicrobial susceptibility test carried out, results showed that Ciproflox, Reflacine and Tarrivid were most effective against the isolates, whileth the isolates mostly showed resistance against Amplicox, Amoxil and Azithromycin. The bacteria species isolated were *Staphylococcus aureus*, *Escherichia coli*, *Staphylococcus spp*, *Bacillus cereus*, *Klebsiella spp*, *Bacillus spp*. The study demonstrated that significant numbers of *E.coli*, and staphylococcus as well as other microbes occur in bath towels.

Key Words: Towels, Bathroom, Hostel, Bacteria, Coliforms, Pathogenicity.

## INTRODUCTION

Bath towels are clothes we use to wipe or clean our body after bath. They collect and accumulate microorganisms from the body and surroundings where they are kept creating an avenue for changes in incidence, pathogens and outcome (Martin, 2012).

Commensals as well as mutualistic microorganisms habit the skin preventing pathogens from taking over the skin ( Hadaway, 2003). At times these normal skin flora cause diseases especially in immune compromised persons (Fedricks, 2007).

As we know the environment is laden with consortium of microorganisms in dust suspensions, bath water bath sponge e.t.c. sometimes bacteria and viruses that habit the intestinal tract of humans can get into bathing water when it gets contaminated with faeces and hence when we clean our body with towels, these organisms are lodged on the bath towels. Some of these organisms include *Escherichia coli*, *Salmonella*, *Rotavirus*, *Enterovirus*, *Pseudomonas aeruginosa*, *Aeromonas hydrophilia*, *Staphylococcus epidermidis*, and *Staphylococcus aureus* (Corazza *et al.*, 2002, Botone & Perez, 1993; Madigan *et al.*, 2000; Chapelle, 2002).

According to Roth and James, (1988) some of these microorganisms are opportunistic especially when there are injuries to the skin or immune compromised persons, they cause severe infections.

Justification of sample size and implication of the study is due to the way female students in those hostels mentioned use and take care of their towels.

### **Aim**

To characterise and identify difference bacteria in bath towels of female students in Rivers State University.

### **MATERIALS AND METHODS**

## **Study Area:**

### **Rivers State University Female hotels (NDDC and POST GRADUATE)**

**Station I:** NDDC Female Hostel

**Station II:** Post-graduate Female Hostel

**Station III:** Control Towel Purchased from Mile 3 Market

## **Sample Collection**

Female bath towels were swabbed from the hostels using sterile swab sticks while wearing sterile gloves. The swab sticks were put in sterile zip lock bags and transported to the laboratory.

## **Preparation of Media**

All the media used in this work including Nutrient Agar, Eosine Methylene blue and MacConkey was prepared according to manufacturers description

## **Antibiogram (Agar disk diffusion method)**

Antibiogram or Antibiotic sensitivity testing is the measurement of the susceptibility of bacteria to antibiotics. A sterile swab stick was dipped into a tube containing, the bacteria suspension and its turbidity is equivalent to 0.5m Mcfarland turbidity standard and the swab stick was pressed against the tube above the fluid level to remove evenly which contained already prepared Mueller hinton agar in three dimension rotating the plate about 60°c each time. The agar plate was allowed to dry for 5 minutes then the antimicrobial disk was impregnated into the agar using a sterile forcep or the surface of the inoculated plate 1.5ml

away from the edge of the plate. Using the head of the sterile forcep, the disk is slightly preserved down to ensure good contact with the agar. After applying the disk, the plates were incubated in an inverted position at 35°C for 16 to 18 hrs. After incubation, the test plates were examined to ensure confluence growth or near confluence. The diameter of each zone of inhibition was measured in millimetres using a ruler on the underside of the plate and recorded for reference purposes (CLSI, 2017).

### **Biochemical Tests**

Biochemical tests are one of the traditional methods for the identification of micro-organisms, usually performed with phenotypic identification.

The ability of micro-organism to utilize certain biomolecules resulting in useful organic compound for themselves forms the basis of various biochemical tests. Biochemical tests are different types, where the identification or distinction between different microorganisms is made on various bases.

#### **Motility Test**

The test is used to differentiate between motile and non-motile micro-organisms. Double strength nutrient agar was dispensed into test tubes, autoclaved to sterilize and allowed to solidify using sterilized wire loop, each isolate was inoculated by stabbing to half the depth of media and incubated at room temperature for about 48 hours. Growth away from the inoculation was recorded as evidence of motility.

#### **Catalase Test (Slide Method)**

This test is used to identify organisms that produce the enzyme catalase. This enzyme detoxifies hydrogen peroxide by breaking it down into water and oxygen. A sterile wire loop is used to transfer a loopful of the test organism to a grease-free slide emulsified with a drop of distilled water. One drop of hydrogen peroxide (6%) was added and observed for effervescence within 3 seconds. The production of air bubbles indicates a positive result.

### **Oxidase Test (Filter Paper Method)**

This test is used to identify micro-organisms containing the enzyme, cytochrome oxidase (important in the electron transport chain). A small portion of the isolate was smeared on part of the filter paper impregnated with freshly prepared oxidase reagent. The reaction was observed within 10 seconds to see if there was any colour change. Deep purple colourations appearing within 5-10 seconds indicate a positive reaction, a weak positive reaction appeared within 70-60 seconds and a negative reaction was indicated by non colour change.

### **Indole Test**

This test is used to ascertain the ability of some microbes to hydrolyse the amino acid tryptophan to produce indole. Tryptophan is made available by trypton in the medium of 10ml of peptone water and dispensed test tubes and sterilized by autoclaving. It was allowed to cool before inoculating isolates into the sterile broth. The broth culture was incubated at 37°C for 48hrs after which 0.5ml of Kovac's reagent was added into each of the culture test tube. The test tubes were shaken and allowed to stand for 5 minutes. Positive results show a red colour at the surface of the medium and negative result showed no red colour at the surface of the medium.

### **Methyl Red Test**

This test is used to identify bacteria producing stable acid by mechanism of mixed acid fermentation of glucose. 17g of methyl red vogues-proskauer (MRVP) broth was suspended in 100ml distilled water. 5ml MRVP broth were distributed into each test tube and autoclaved at 121°C for 15 minutes. A loopful of the test organism was inoculated into the broth and incubated for 48hrs. After incubation 2-5 drops of methyl red indicator was added to the culture. Red colour change indicated a positive result and yellow colour change indicated a negative result.

### **Vogues-Proskauer Test**

This test is used to detect acetone (an important physiological metabolite excreted by many micro organisms) in a bacteria broth culture. A loopful of the test organisms is inoculated into MRVP broth and incubated for 24hours. After incubation, 0.6ml (9 drops) of  $\alpha$ -naphthal and 0.2 (3days) of potassium hydroxide were dropped into the broth culture and was shaken and allowed to stand for 15minutes. A pink or red colour at the surface of the medium indicates a positive result and a copper colour at the surface of the medium indicates a negatives result.

### **Sugar Fermentation**

This test is used to share the utilization of sugar by bacteria as a sole source of carbon with the production of either acid or gas, or both and colour change. 2.08g of peptone is dissolved in 200ml of distilled water (Peptone water broth). An indicator to indicate the reactivity by colour change (methyl red some) was added to top tone water. Equal quantity of the already prepared peptone was is dispensed into different sterile conical flasks the available sugars (1.0g) are respectively dissolved into the peptone water and then dispensed

in 10ml into test tubes containing inverted Durham tubes and autoclaved. The respective isolates were inoculated aseptically and incubated for 24-48hrs at 37°C. Positive results have colour changes from red to yellow or from yellow to colourless indicating acid production. Gas production was noted by presence of air space in the Durham tubes within the test tubers. Retention of initial colour indicates negative result.

## RESULTS

The results for the total heterotrophic count of bacteria for the first and second sampling of bath towels were presented in table 4.1 and 4.2. The results showed that NDDC hostel had the highest count for the first sampling at  $3.17 \times 10^4$  cfu/ml and post graduate hostel had the highest count for the sample sampling at  $1.17 \times 10^4$  cfu/ml.

In table 4.3 and 4.4, results for total coliform are presented and they showed that NDDC hostel had the highest count for first sampling at  $2.99 \times 10^4$  cfu/ml and post graduate hostel had the highest count for the second sampling at  $0.41 \times 10^4$  cfu/ml.

In table 4.5 and 4.6, results for occurrences of isolates on the samples were presented. In the first sampling the isolates *Staphylococcus aureus*, *Bacillus cereus*, *E.coli* and *Klebsiella* sp had the highest occurrence at 9.30% and the lowest occurrence at 2.33% for *Staphylococcus* sp & *Bacillus* sp. In the second sampling, the highest occurrence happened in *E.coli* at 21.7%, while the lowest was *Staphylococcus aureus* at 8.7%.

In table 4.7 and 4.8, results for morphological and Biochemical test were presented. The results showed that about 59% of the isolates were gram positive and 41% gram negative.

**Table 4.1: Total Heterotrophic count (First Sampling)**

Location	Sample	THB1	THB2	Mean	CFU/ml
NDDC Hostel	A	190	210	200	2.0 x10 <sup>4</sup>
	B	113	92	103	1.03 x10 <sup>4</sup>
	C	308	294	301	3.01 x10 <sup>4</sup>
	D	220	215	218	2.18 x10 <sup>4</sup>
	E	315	318	317	3.17 x10 <sup>4</sup>
PG Hostel	F	150	145	148	1.48 x10 <sup>4</sup>
	G	68	42	55	0.55 x10 <sup>4</sup>
	H	207	195	201	2.01 x10 <sup>4</sup>
	I	40	37	39	0.39 x10 <sup>4</sup>
	J	321	301	311	3.11 x10 <sup>4</sup>
	Control	8	15	12	0.12 x10 <sup>4</sup>

**Table 4.2: Total Heterotrophic count (Second Sampling)**

Location	Sample	THB1	THB2	Mean	CFU/ml
NDDC Hostel	A	76	69	73	$0.73 \times 10^4$
	B	40	33	37	$0.37 \times 10^4$
	C	46	54	50	$0.5 \times 10^4$
	D	17	10	14	$0.14 \times 10^4$
	E	117	112	115	$1.15 \times 10^4$
PG Hostel	F	33	20	27	$0.27 \times 10^4$
	G	50	28	39	$0.39 \times 10^4$
	H	66	52	59	$0.59 \times 10^4$
	I	112	122	117	$1.17 \times 10^4$
	J	87	98	93	$0.93 \times 10^4$
Mile 3 Market	Control	10	22	16	$0.16 \times 10^4$

**Table 4.3: Total Coliform count (First Sampling)**

<b>Location</b>	<b>Sample</b>	<b>THB1 (10<sup>-1</sup>)</b>	<b>THB2 (10<sup>-1</sup>)</b>	<b>Mean</b>	<b>CFU/ml</b>
NDDC Hostel	A	50	46	48	0.48 x10 <sup>4</sup>
	B	47	36	42	0.42 x10 <sup>4</sup>
	C	140	210	175	1.75 x10 <sup>4</sup>
	D	281	215	248	2.48 x10 <sup>4</sup>
	E	292	305	299	2.99 x10 <sup>4</sup>
PG Hostel	F	288	292	290	2.9 x10 <sup>4</sup>
	G	15	21	18	0.18 x10 <sup>4</sup>
	H	140	165	153	1.53 x10 <sup>4</sup>
	I	26	16	21	0.21 x10 <sup>4</sup>
	J	18	17	18	0.18 x10 <sup>4</sup>
	Control	17	19	18	0.18 x10 <sup>4</sup>

**Table 4.4: Total Coliform count (Second Sampling)**


<b>Location</b>	<b>Sample</b>	<b>THB1 (10<sup>-1</sup>)</b>	<b>THB2 (10<sup>-1</sup>)</b>	<b>Mean</b>	<b>CFU/ml</b>
NDDC Hostel	A	33	29	31	0.31 x10 <sup>4</sup>
	B	31	25	28	0.28 x10 <sup>4</sup>
	C	36	24	30	0.30 x10 <sup>4</sup>
	D	24	17	21	0.21 x10 <sup>4</sup>
	E	16	10	13	0.13 x10 <sup>4</sup>
PG Hostel	F	16	28	22	0.22 x10 <sup>4</sup>
	G	21	32	27	0.27 x10 <sup>4</sup>
	H	32	23	28	0.28 x10 <sup>4</sup>
	I	39	42	41	0.41 x10 <sup>4</sup>
	J	15	21	18	0.18 x10 <sup>4</sup>
Mile 3 Market	Control	13	20	17	0.17 x10 <sup>4</sup>

**Table 4.5: Frequency Table (First Sampling)**

Isolates	A	B	C	D	E	F	G	H	I	J	Control	Total	Percentage (%)
HO <sub>1</sub>	+			+					+	+		4	9.30%
HO <sub>2</sub>			+			+				+		3	6.97%
HO <sub>3</sub>									+	+	+	3	6.97%
HO <sub>4</sub>	+								+			2	4.65%
HO <sub>5</sub>									+			1	2.33%
HO <sub>6</sub>			+	+					+			4	9.30%
HO <sub>7</sub>						+			+			2	4.65%
HO <sub>8</sub>			+			+			+			3	6.97%
HO <sub>9</sub>	+			+	+						+	4	9.30%
HO <sub>A</sub>					+				+			2	4.7%
HO <sub>B</sub>									+		+	2	4.7%
HO <sub>C</sub>									+			1	2.33%
HO <sub>D</sub>			+									1	2.33%
HO <sub>E</sub>						+						1	2.33%
HO <sub>F</sub>	+					+						2	4.65%
HO <sub>E1</sub>			+			+				+	+	4	9.30%
HO <sub>E2</sub>		+			+					+	+	4	9.30%
<b>Total</b>												<b>43</b>	<b>100%</b>

**Table 4.5: Frequency Table (Second Sampling)**

Isolates	1	2	3	4	5	6	7	8	9	10	Control	Total	Percentage (%)
1	+			+						+		3	13.0%
2			+					+				2	8.7%
3			+			+		+				3	13.0%
4	+								+	+		3	13.0%
5	+				+		+					3	13.0%
6		+		+	+						+	4	17.4%
7	+				+				+	+	+	5	21.7%
<b>Total</b>												<b>23</b>	<b>100%</b>



**Figure 4.1: % Occurrence of Bacterial Isolates in the Towel Sample**

**Table 4.7**

Colonial Morphology							Biochemical Reaction							Sugar Fermentation			Probable Organism		
Isolates	Margin	Colour	Texture	Shape	Elevation	Surface	Gram	Indole	Citrate	Methyl Red	Voges-Proskauer	Motility	Oxidase	Catalase	Lactose	Sucrose	Glucose	Mannitol	
HO <sub>1</sub>	Entire	Yellow Gold	Moist	Cocci	Raised	Smooth	+	-	-	+	-	-	-	+	A	A	A	-	<i>Staphylococcus aureus</i>
HO <sub>2</sub>	Entire	Creamy	Moist	Rod	Raised	Smooth	-	+	+	-	+	+	+	+	-	A	AG	AG	<i>Escherichia coli</i>
HO <sub>3</sub>	Entire	Creamy	Dry	Rod	Flat	Smooth	-	-	-	+	+	+	+	-	-	A	AG	AG	<i>Klebsiella sp</i>
HO <sub>4</sub>	Entire	Creamy	Dry	Rod	Flat	Smooth	+	-	+	+	-	-	-	+	-	-	A	-	<i>Bacillus sp</i>
HO <sub>5</sub>	Entire	Yellow Gold	Moist	Cocci	Raised	Smooth	+	+	+	+	+	+	-	+	-	-	A	-	<i>Staphylococcus sp</i>
HO <sub>6</sub>	Entire	Milky	Dry	Rod	Flat	Smooth	+	-	+	+	-	+	+	+	-	-	A	-	<i>Bacillus cereus</i>
HO <sub>7</sub>	Serrated	Creamy	Dry	Cocci	Flat	Smooth	+	-	+	-	+	-	+	+	-	A	AG	AG	<i>Staphylococcus aureus</i>
HO <sub>8</sub>	Entire	Yellow Gold	Moist	Cocci	Raised	Smooth	+	+	+	-	-	-	+	+	A	A	A	A	<i>Staphylococcus sp</i>
HO <sub>9</sub>	Entire	Creamy	Moist	Rod	Raised	Smooth	-	+	+	+	-	+	+	+	-	AG	A	AG	<i>Escherichia coli</i>
HO <sub>A</sub>	Serrated	Creamy	Moist	Rod	Raised	Smooth	-	+	+	-	+	-	+	+	A	AG	A	A	<i>Escherichia coli</i>

<b>HO<sub>C</sub></b>	Entire	Creamy	Dry	Rod	Raised	Smooth	+	+	+	+	-	+	+	+	A	A	A	-	<i>Bacillus sp</i>
<b>HO<sub>D</sub></b>	Entire	Creamy	Moist	Rod	Raised	Smooth	+	-	+	-	+	+	+	+	-	-	-	A	<i>Bacillus sp</i>
<b>HO<sub>E</sub></b>	Serrated	Creamy	Moist	Rod	Flat	Smooth	-	+	+	-	+	+	+	+	A	AG	AG	AG	<i>Escherichia coli</i>
<b>HO<sub>F</sub></b>	Entire	Creamy	Moist	Cocci	Flat	Rough	+	+	+	+	-	+	+	+	A	A	A	A	<i>Staphylococcus sp</i>
<b>HO<sub>E1</sub></b>	Entire	Creamy	Moist	Rod	Raised	Smooth	-	-	+	-	+	-	+	+	-	AG	A	AG	<i>Klebsiella sp</i>
<b>HO<sub>E2</sub></b>	Entire	<b>HO<sub>B</sub></b>	Entire	Creamy	Dry	Rod	-	-	+	-	+	+	-	+	+	+	A	A	A

**Table 4.8**

Colonial Morphology							Biochemical Reaction						Sugar Fermentation				Probable Organism		
Isolates	Margin	Colour	Texture	Shape	Elevation	Surface	Gram reaction	Indole	Citrate	Methyl Red	Vogus-Proskauer	Motility	Oxidase	Catalase	Lactose	Sucrose	Glucose	Mannitol	
1	Entire	Creamy	Moist	Rod	Raised	Smooth	-	+	+	-	+	+	+	+	-	AG	A	AG	<i>Escherichia coli</i>
2	Entire	Yellow	Moist	Cocci	Raised	Smooth	+	-	-	+	-	-	-	+	A	A	A	-	<i>Staphylococcus aureus</i>
3	Entire	Yellow Gold	Moist	Cocci	Raised	Smooth	+	-	-	+	-	-	-	+	A	A	A	-	<i>Staphylococcus aureus</i>
4	Entire	Creamy	Moist	Rod	Flat	Smooth	-	+	+	-	+	-	+	+	-	A	AG	AG	<i>Klebsiella sp</i>
5	Serrated	Creamy	Dry	Cocci	Flat	Smooth	+	+	+	+	-	-	+	+	A	A	A	-	<i>Staphylococcus sp</i>
6	Entire	Milky	Dry	Rod	Flat	Smooth	+	-	+	+	-	-	-	+	-	A	-	-	<i>Bacillus sp</i>
7	Entire	Creamy	Moist	Rod	Flat	Smooth	-	+	+	-	+	+	+	+	A	AG	AG	AG	<i>Esherichia coli</i>

**Table 4.9: Antibiotic susceptibility pattern of Bacterial isolates from Bath Towels**

Isolates	Gram Positive									
	CPX	S	SXT	E	PEF	CN	APX	Z	AM	R
Bacillus sp.	S	I	R	I	S	S	I	S	S	S
Staphylococcus sp.	S	S	S	I	S	S	R	R	R	I
Bacillus cereus	S	R	R	R	S	R	R	R	R	R
Bacillus sp.	S	R	R	S	S	S	R	R	R	R
Staphylococcus sp.	S	I	R	I	S	R	R	R	R	I
Staphylococcus aureus	S	I	R	I	S	R	R	R	R	I
Bacillus sp.	S	R	R	S	S	I	R	R	R	R
Bacillus sp.	S	I	S	S	S	R	R	R	R	I
Staphylococcus sp.	S	S	R	R	S	R	R	R	R	I
Staphylococcus sp.	S	S	S	I	S	S	R	R	R	I

**Table 4.10: Antibiotic susceptibility pattern of Bacterial isolates from Bath Towels**

Isolates	Gram Negative									
	OFX	S	SXT	CH	SP	CPX	AM	AU	CN	PEF
E. coli	S	S	S	S	S	S	R	R	R	S
Klebsiella sp.	S	R	S	R	S	S	R	R	R	S
E. coli	S	I	S	I	S	S	R	I	R	I
E. coli	R	R	R	R	R	R	R	R	R	R
E. coli	S	S	I	R	S	S	R	R	I	S
E. coli	S	R	R	R	S	S	R	R	R	S
Klebsiella	S	R	S	R	S	S	R	R	R	S

**Key:**

CPX	-	Ciproflox,	PEF	-	Reflacine	S	-	Sensitive
S	-	Streptomycin	CN	-	Gentamycin	I	-	Intermediate
E	-	Erythromycin	AM	-	Amoxil	R	-	Resistant
APX	-	Ampiclox	SXT	-	Seprin			
Z	-	Azithromycin	R	-	Rifampicin			
AU	-	Augmentin	SP	-	Spectinomycin			
CH	-	Chloramphenicol	OFX	-	Tarrivid			

## DISCUSSION

The analyses in this study were carried out to know the characteristics and identified bacteria in towels of female students in Rivers State University hostels. Counts of total heterotrophic bacteria ranged  $2.0 \times 10^4$  cfu/ml to  $3.17 \times 10^4$  cfu/ml in the NDDC hostel and  $1.48 \times 10^4$  cfu/ml to  $3.11 \times 10^4$  cfu/ml in post graduate hostel for the first sampling with the control at  $0.12 \times 10^4$  cfu/ml. Total heterotrophic count for the second sampling ranged from  $0.73 \times 10^4$  cfu/ml to  $1.15 \times 10^4$  cfu/ml in NDDC hostel,  $0.12 \times 10^4$  cfu/ml to  $0.93 \times 10^4$  cfu/ml in the post graduate hostel with the control having  $0.16 \times 10^4$  cfu/ml. Total coliform count for first sampling ranged from  $0.48 \times 10^4$  cfu/ml to  $2.99 \times 10^4$  cfu/ml in NDDC hostel and  $2.9 \times 10^4$  cfu/ml to  $0.18 \times 10^4$  cfu/ml in the post graduate hostel with the control having  $0.18 \times 10^4$  cfu/ml. Total coliform count for the second sampling ranged from  $0.31 \times 10^4$  to  $0.13 \times 10^4$  cfu/ml in NDDC hostel,  $0.22 \times 10^4$  to  $0.18 \times 10^4$  cfu/ml in the post graduate hostel and the control having  $0.17 \times 10^4$  cfu/ml.

The bacterial isolate identified were *staphylococcus aureus*, *Escherichia coli*, *staphylococcus sp.*, *Bacillus cereus*, *klebsiella spp*, *Bacillus spp*. This agrees with the work of (Hannah et al., 2020) in which *E.coli* and *Staphylococcus aureus* were isolated from towel samples.

The NDDC hostel recorded the highest heterotrophic bacteria count at  $3.17 \times 10^4$  for the first sampling while post graduate hostel had the highest heterotrophic bacteria count for the second sampling at  $1.17 \times 10^4$ . For the total coliform count, NDDC hostel recorded the highest count at  $2.99 \times 10^4$  cfu/ml for the first sampling and post graduate hostel had the highest count for the second sampling at  $0.41 \times 10^4$  cfu/ml. The controls for both total heterotrophic bacteria

and total coliform counts for the first and second sampling were significantly low compared to these of NDDC hostel and post graduate hostel. This may be attributed to it being a new towel that had not been used for cleaning purposes.

The significant differences in bacterial counts between the two hostels may be due to variations in hostel conditions, water supply/quality and general handling by females in the different hostels. This is in agreement with the work of (Sturt, 2015, Bradford, 2018). The differences also in counts between the first and second sampling may be attributed to the week interval between the samplings. The students were told to wash the towels and use for a few days prior to the second sampling.

Based on this study, it was observed that *E.coli* occurred in about 57% of the samples. This can be attributed to the fact that bathrooms of all hostels sampled for this study were close to toilets. Toilets are very likely sources of *E.coli* contamination (Hannah et al., 2020). *Staphylococcus aureus* also occurred in about 36% of the samples. This is attributed to *S. aureus* being a normal flora of the skin. This agrees with the work of (Oller and Mitchel, 2008) that *S. aureus* occurs in cotton towels. It also agrees with the work of (Neely and Maley, 2000) that observed that *Staphylococcus* could survive for 19-21 days on cotton fabrics, which can be attributed to the significant occurrence of *Staphylococcus* species in the second sampling even after they were washed (Oller et al., 2008).

The results were analyzed using student's T test to compare bacterial loads in the samples from the two hostels and the mean values were: Total heterotropic bacteria 2.2750E2 for NDDC hostel and 1.5060E2 for PG hostel in the first sampling, 57.4000 for NDDC hostel and 66.8000 for post graduate hostel in the second sampling. Total coliform counts were 1.6220E2

for NDDC hostel and 99.8000 for post graduate hotel in the first sampling and 24.5000 for NDDC hostel and 26.9000 for post graduate hostel in the second sampling. It showed that there was no significant difference in the bacterial loads of NDDC hostel and post graduate hostel in the first and second sampling in our study, we have tested isolated bacterial species for their sensitivity pattern against the commonly prescribed antibiotics according to the CLSI guideline Antibiotic Suceptibilityy of Staphylococcus sp was tested against 10 commonly prescribes and available antibiotics (Ciproflox, Reflacine, Streptomycin, Gentamycin, Septrin, Erythromycin, Ampiclox, Amoxil, Rifampicin and Azithromycin), using agar diffusion method.

The results show that the susceptibility of Staphylococcus sp was higher Ciproflox, and Reflacine (100%) followed by Septrin (60%) and Gentamycin (40%). Staphylococcus sp showed higher resistance to Ampiclox and Amoxil with 100% for each of them.

For Bacillus species isolates, Ciproflox and Reflaxine also had the highest susceptibility rates at 100% each, followed by Erythromycin and Gentamycin with 40% sensitivity each Septrin had the highest resistance rate with 100% resistance followed by Ampiclox with 80% resistance.

In the case of E.coli isolates, the results showed that the susceptibility was higher for Tarrivid (100%), followed by Ciproflox with 80% sensitivity. The highest percentage of resistance was recorded against Ampiclox (100%) and Augmentin and Gentamycin at 80% sensitivity each.

For the Klebsiella species isolates, the results showed the susceptibility was 100% for Tarrivid, Septrin, Ciproflox and Reflacine while it recorded 100% resistance against Septrin, chloramphenicol, Augmentin, Amoxil and Gentamycin. In general, the results show that

Ciproflox and Reflacine were most effective against the isolates. All the isolates showed high level of resistance against Ampiclox, Amoxil and Azithromycin. E. coli and Klebsiella sp. were also highly sensitive to Tarrivid.

## **Conclusion**

The study on the characterization and identification of bath towels of female students in Rivers State University hostels revealed that micro organisms are present in baths towels in various varieties and degrees. The morphological and biochemical identification of bacterial isolates revealed the presence of *Escherichia coli*, *Klebsiella spp*, *Staphylococcus spp* and *Bacillus sp*. Some of the genera (*Staphylococcus spp*, *Klebsiella spp*, *Bacillus spp* etc) were observed to be capable of being opportunistic pathogens under the right conditions.

Towels are necessary and useful items, but can also be a breeding ground for bacteria.

This however can be greatly reduce if these items and kept clean at all times.

## **References**

- Botone, E. S., & Perez, A.A. (1993). *Pseudomonas aeruginosa* folliculitis acquired through use of a contaminated loofah sponge: an unrecognized potential public health problem. *J. Clin Microbiol.* 1993 Mar; 31 (3) 480-483.
- Bradford, A. (2018). How often should you wash bath towels? It depend: [Online Available from [https://www-inet-com.cdn.ampproject.org/v/s/www.cnet-com/google-amp/news/how-often-should-you-wash-bath-towels-heres-the-answers/?amp\\_js\\_v=a2&usqp=mp331AQareferrer=https%3A%2F%2Fwww.google.com&tf=From%20%251%24s&sampshare](https://www-inet-com.cdn.ampproject.org/v/s/www.cnet-com/google-amp/news/how-often-should-you-wash-bath-towels-heres-the-answers/?amp_js_v=a2&usqp=mp331AQareferrer=https%3A%2F%2Fwww.google.com&tf=From%20%251%24s&sampshare)] Accessed: 20<sup>th</sup> June, 2019.]
- Chapelle, F.H. (2000). Ground water Microbiology and Geochemistry. New York: John Wiley & Sons, 2000. 97-107.

- Clinical and Laboratory Standard Institute (2017). Performance Standards for Antimicrobial Susceptibility Testing twenty-five information supplement. CLSI document M100-S21 (ISBN 1-56238-742-1) Clinical and Laboratory Standard Institute, 940 West Valley road, Suite 1400, wayne, Pennsylvania 19087 USA, 30(1), 68-70.
- Corazza, M., Carla, E., Rossi, M.R., Pedna, M.F., & Olgili, A. (2002). Face and body sponges : Beauty aids or Potential Microbiological Reservoir? *Eur J Dermatol.* 2002 Nov-Dec; 13(6): 511-3 PMID: 14721777.
- Fredricks, D. N. (2001). Microbial ecology of human skin in health and disease. *J. investing Dermatol symp proc.* 2001:6:167-167.
- Hadaway, L. C. (2002). Skin flora and infection. *J. Infus Nurs.* 4(2): 3-9.
- Hannah, T., Asumang, Z. A., Imoro & S. T. Kporde (2020). Toothbrush and Towel Handling and their Microbial Quality. The case of students of University for Development Studies, Nyankpala Campus, Ghana. *Afr, J. Infect Dis* 15(1): 41-46.
- Madigan, M. M., J. Martinko, & Parker, J. (2000). Brock Biology of Microorganisms. 8<sup>th</sup> ed Upper Saddle River, NJ: Prentice Hall 2000.
- Martin, G. S. (2012). “Sepsis, severe sepsis and septic shock: changes in incidence, pathogens and outcomes.” *Expert Review of Anti-Infective Therapy.* 10(6): 701-6
- Neely, A. N., & Maley, M. P. (2000). Survival of Enterococci and Staphylococci on hospital fabrics and plastics. *J Clin Microbiol* 38:724-726.
- Oller, A. R. & Mitchell, A. ( 2008). Staphylococcus aureus recovery from cotton towels. *J infect Developing countries.* 3(3): 224-228.
- Roth, R. R. & James, W. D. (1988). “Microbial ecology of the skin.” *Anna Rev Microbiol.* 42 (1): 4441-64.
- Sturt, K. (2015). How often you should actually be washing your towels and streets [Online] Available from: <https://m.huffnostcom/us/entr/us-55cb927e4boflcbfle72> [Accessed : 20th June, 2019].

