

Antimicrobial Activity and Phytochemical Screening of Methanolic Leaf Extract of *Vernonia amygdalina*

ABSTRACT

Aims: Crude methanolic leaf extract of *Vernonia amygdalina* was evaluated to determine its bioactive constituents, the antimicrobial properties, measure the minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) against some selected bacterial organisms.

Place and Duration of Study: The investigation was carried out at University of Maiduguri in Borno State, Nigeria. The herb obtained from Lake Chad Research Institute were identified and validated by the department of Forestry and wild life, Mohammed Lawan College of Agriculture (MOLCA) in Maiduguri, Borno State.

Methodology: Crude methanolic leaf extraction of plant, qualitative phytochemical screening, antimicrobial sensitivity against some disease-causing organisms, minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) were carried out.

Results: Phytochemical screening of *V. amygdalina* leaves extract revealed the presence of tannins, saponins, terpenoids, flavonoid, carbohydrates and cardiac glycosides. The antimicrobial sensitivity shows *P. aeruginosa*, had the highest sensitivity with effect at all concentrations (26mm at 1000mg/ml), the lowest against *Salmonella typhi* were 9mm at 1000mg/ml and 7mm at 200mg/ml, while the highest were 17mm at 400mg/ml and 14mm at 200mg/ml. When tested at all concentrations (200-1000 mg/ml), *S. aureus*, *S. pyogenes*, *B. subtilis*, *Corynebacteria species*, and *K. pneumonia* showed greater sensitivity than *Salmonella typhi* but less sensitivity to *P. aeruginosa*. Based on the outcomes of the minimum inhibition concentration (MIC) and minimum Bactericidal (MBC) results, respectively shows *S. aureus* was found to be sensitive at only 100-200mg/ml, however *P. aeruginosa* was found to have the highest sensitivity with effect at all concentrations (25-200mg/ml) extract for the lowest (12.5).

Conclusion:

The study highlighted the antimicrobial effects of *V. amygdalina* leave extracts on some pathogens thereby verifying the traditional healer's claim. Also, it was concluded that the **extract of *V. amygdalina* contained** pharmacologically active phytochemicals which could be responsible for the numerous medicinal properties exhibited by the plant leaf extract.

Keywords: *Vernonia amygdalina*; methanol; antimicrobial; phytochemicals; minimum inhibitory concentration; minimum bactericidal concentration.

INTRODUCTION

Plants have been utilized to cure and manage a variety of ailments due to the presence of phytochemicals. Worldwide, there is a steady rise in the usage of medicinal herbs. The advent of resistant diseases and the advancement of scientific understanding of herbal remedies as significant treatment alternatives have led to an increase in the quest for therapeutic molecules originating from plant species^[1]. Despite the significant advances in modern medicine, phytotherapy is still widely utilized today since medicinal plants have been used to prevent and treat a variety of health issues for many years^[2-4]. Depending on a number of variables like the weather, ecology, the time and age of

collection, and other considerations, the various phytochemicals generated by plants can vary both qualitatively and quantitatively ^[4-5].

Vernonia amygdalina is well known for its wide range of uses in traditional medicine ^[4]. A member of the family Asteraceae, *Vernonia amygdalina* is also known as bitter leaf in English, ewuro in Yoruba, Shiwaka in Hausa, and olubo in Igbo ^[6].

Bitter leaf is a 2–5 m tall shrub plant that thrives in a variety of ecological zones throughout Africa. It produces enormous amounts of forage and is drought tolerant. The leaf is green and has a distinct odour and harsh taste ^[7]. Non-nutritional elements such as alkaloids, saponins, tannin, terphenoids, glycosides, and steroids are to blame for the bitter test ^[6,8].

Vernonia amygdalina is a perennial shrub. It is a vegetable that is frequently utilized in tropical African cuisine and folk medicine. Traditional users claim that this plant has palpable health advantages. Numerous vitamins, minerals, and phytochemicals are present in it, including iodine, alkaloids, anthraquinones, edotides, sesquiterpene lactones, and steroid glycosides. The health advantages of the plant are linked to these bioactive substances. Traditional medicinal uses not yet supported by scientific evidence have been indicated ^[9].

Vernonia amygdalina favourable effects on health are invariably attributed to its nutritional and phytochemical richness. Many of its mechanisms of action have not been fully understood in humans, despite the presence of compounds with anticancer effects, antioxidant properties, antimalarial properties, anti-inflammatory properties, antibacterial characteristics, and hypolipidemic benefits ^[9].

Alkaloids, steroids, flavonoids, phenols, saponins, terpenes, cyanogenic glucoside, tannin, anthraquinone, phytate, oxalate, and lignans have been identified in *V. amygdalina* by phytochemical analysis ^[10,11].

These phytochemicals may be the cause of its physiological effects, which include those that are hypolipidemic, anti-diabetic, anti-microbial, anti-allergic, antibacterial, antimalarial, anti-fungal, anti-cancer, anti-leukemia, anti-inflammatory, analgesic, antipyretic, anti-oxidative, anti-helminthic, and antimicrobial ^[11-13].

V. amygdalina extracts in aqueous, ethanol, and methanol suppress bacteria and fungi linked to food deterioration and pathogenicity. Extracts of *V. amygdalina* leaves have been reported to be effective against a variety of microorganisms, including *Candida albicans*, *Pseudomonas aeruginosa*, *Bacillus subtilis*, *Staphylococcus aureus*, and *Escherichia coli*. Periodontal bacteria are prevented from proliferating when chewing on sticks made of *Vernonia amygdalina* wood. There is evidence that leaves can inhibit a number of dangerous bacteria and viruses ^[14-16]. Alkaloids, flavonoids, and saponins are what give *V. amygdalina* its antibacterial characteristics. Vernodalol, vernodalin, and vernolide are some of the specific chemicals linked to this plant's antibacterial properties (sesquiterpene lactones) ^[16]. Sesquiterpene lactones have been linked to antibacterial, anti-protozoal, and anti-tumor effects. ^[17-19]

The leaves of *Vernonia amygdalina* are primarily consumed by humans. Washing is frequently required to lessen the harsh taste that water-soluble saponins generate. Sesquiterpene lactones (such as vernodalin, vernolepin, and vernomygdin) and steroid glycosides are what give *V. amygdalina* its bitter flavor (vernioniosides). Usually, a fresh leaf is washed before being added to food while it is being prepared. African breadfruit, yam porridge, and other foods including soups (such as Ogbono, Egusi, Okro, and the well-known bitter leaf soup) can all be made using *V. amygdalina*. The leaves are consumed as a vegetable and aid with digestion. The leaves have been used in place of hops in the beer-brewing sector ^[20]. The stems are eaten by domestic animals. They are also used by humans as a chewing stick. The leaves may be used as a hop substitute and an antioxidant ^[21].

Traditional medicine has utilized *Vernonia amygdalina* extensively to manage and treat a variety of illnesses. It is employed in some situations to control and treat conditions like diabetes, malaria, and gastrointestinal problems. Additionally, a local infusion of the leaves or roots is used to treat microbiological infections, schistosomiasis, cough, hepatitis, and other disorders that are spread through sexual activity ^[22,23]. The roots and leaves are used in traditional medicine to cure fever, hiccups, and renal issues while, ^[24], the wood, from the root, can be used to clean the tooth ^[25]. According to reports ^[26], *V. amygdalina*'s fresh leaves work wonders as a purgative. In some parts of Africa, it is used to make cough medicine ^[27].

One of *Vernonia amygdalina's* most potent substances, saponins are complex molecules whose glycans are triterpenoids or steroidal ^[29,28]. Saponins are starch glycosides that have three hydroxyl groups connected to complicated sugar components ^[29]. It is an amorphous white or light brown powder that initially tastes pleasant before turning bitter and even acidic. The gastrointestinal tract is thought to be too irritated by saponins for them to be employed internally ^[29,30]. In experiments, *Vernonia amygdalina's* steroidal saponin levels appear to be higher than those of other plants ^[29,31]. The leaves of *Vernonia amygdalina* also contain flavones (flavonoids), which are antioxidants with therapeutic benefits against carcinogens and aging ^[29,32].

Clinical research is needed to support the benefits of this plant product on health. More animal models' experiments are required to prove the potency of this plant product. Grey areas where to encourage the efficient use of this plant product, study is indicated Based on conclusions from compositional analysis and animal bioassays, this plant product is believed to have health advantages. There is a need to establish these claims in humans. There is only very little information on the food uses of *V. amygdalina* which has been related to potential health benefits. There are a lot of claims on this plant-based on ethnomedicine and traditional uses ^[9].

Numerous studies indicate that its use is largely dose-dependent for successful results ^[33]. When taken as an antimalarial or to increase fertility ^[34], it may occasionally exhibit adverse consequences ^[35]. Therefore, the goal of this study is to evaluate the Phytochemicals and antimicrobial activities of *Vernonia amygdalina* (Better Leaf) leaf extracts.

MATERIALS AND METHODS

Study Area

The investigation was carried out at University of Maiduguri in Borno State, Nigeria. Between latitudes 10.2° N and 13.4° N and longitudes 9.8° E and 14.4° E, Maiduguri is at a latitude of 354 m above sea level. The dry (October–May) and wet (June–September) seasons are the most common. The yearly rainfall is between 9 and 198 mm, the temperature goes from 13 °C to 41 °C, and the relative humidity fluctuates between 19% and 78% but stays around 45% during the wet season. Seven to nine hours a day are spent in the sun ^[36].

Plant Components

The Lake Chad Research Institute provided the *V. amygdalina* leaves, which the Forestry Department of the Mohammed Lawan College of Agriculture (MOLCA) in Borno State identified and validated the plant. The leaves are cleaned with fresh water before being dried under a shed at room temperature. The dried leaves are then ground into a fine powder using a wooden pestle and mortar and stored in a clean plastic container for further processing.

Methodology

Extraction of the Plant Material

1.5 liters of 95% methanol were added to the two-liter round bottom flask in which the powdered material had been previously placed. The combination was refluxed for approximately two hours, after which the solution was withdrawn for filter paper-based debris removal. After evaporation a (green) substance was obtained, the entire solution was concentrated on hot air ovum at 40 °C – 50 °C. Take the weight and transfer to an air tight container for analysis.

Phytochemical Screening of the Extract

Using the approach outlined by ^[37] ^[38], the crude aqueous extract of *V. amygdalina* was subjected to qualitative chemical screening for the identification of the various classes of chemical constituent.

i. Test for alkaloids

On a steam bath, precisely 0.5g of the extract were mixed with 5.0 ml of 2 M aqueous hydrochloric acid. 1.0 ml of the filtrate will be treated separately with a few drops of Mayer's reagent; the appearance of buff-colored will be an indication for the presence of alkaloid; the appearance of

orange red precipitate from Drangendoff's reagent; and the appearance of dark-brown precipitate from Wangner's reagent; will be an indication for the presence of alkaloid ^[39].

Test for Tannins

10ml of distilled water and precisely 0.5g of plant extracts was combined. A few drops of a 1% solution of ferric chloride will be added to the mixture after filtering, and 2.0 ml of the filtrate will be used for the test. The presence of tannin will be indicated by the presence of blue-black, green, or blue-green precipitate colors. The filtrate will be mixed with an equal volume of 10% lead ethanoate. The occurrence of tannins will be revealed by the production of white precipitate. Three drops of 10% HCL and one drop of methanol will be added to the filtrate of the extract when it is heated. The presence of tannins will be determined by a crimson precipitate ^[40-41].

i. Test for Phlobatannins

The extracts were heated in a little amount of distilled water before being filtered. With 1% aqueous HCl, the filtrate will be further heated. Phlobatannins' appearance ^[41].

ii. Test for Glycosides

a. Liebermann-Bur Chard's Test. (Test for Steroid Nucleus).

2ml of acetic anhydride was added to the extract (0.5g). The mixture will cool in ice before being placed in a cone. Carefully added tetraoxosulphate (vi) acid will be used. If the color changes from violet to blue or bluish-green, a steroidal ring is present ^[42].

a. Salkowski's Test. (Test for steroidal Nucleus)

Tetraoxosulphate (VI) acid was carefully added by the side of the test tube to generate a lower layer after the addition of 2ml of chloroform to extract (0.5g). The presence of a steroidal ring is indicated by the appearance of a reddish-brown color or yellow in the interphase ^[42].

Test for Saponins

To ascertain the presence of saponins, the froth test and emulsion test as reported by ^[43] will be employed. In a 100 ml beaker, distilled water (20 ml) was mixed with a little amount of each plant extract, boiled, and the filtrate utilized for the test:

(a) The froth test was performed using 5 ml of filtrate diluted with 20 ml of water, forcefully shaken, and left to stand for 30 minutes. The outcome was noted.

(b) A tiny amount of filtrate was mixed with 2.5 ml of fehling's solutions A and B in an equal volume. A brick-red precipitate's appearance was a sign that glycosides were present ^[44].

Test for free anthraquinones (Borntrager's test)

10ml of aqueous tetraoxosulphate (vi) acid was added to the extract (0.5g), agitated, and then transferred while it is still hot. Five milliliters of benzene will be mixed with the filtrate. After being separated, the benzene layer will be given half of its own volume of 10% ammonia solution. Indicators of mixed anthraquinones include pink, red, and violet coloring in the ammoniac (lower) phase ^[41].

Test for flavonoids

A. Test for Ferric Chloride

A tiny amount of the extract will be heated with water before being filtered. the filtrate, to 2ml.

B. The Shinoda Test

The extract (0.5g) were warmed and filtered after being resolved in ethanol. Then, a few drops of com will be poured to the filtrate, followed by the pieces of magnesium chips. HCl Flavonoids will be present if the coloration is pink, orange, or red to purple ^[45].

C. Lead Acetate Test

5ml of distilled water were used to dissolve up to 0.5g of the extract. 3.0 ml of 10% lead acetate solution was added. The presence of flavonoids was revealed by the formation of buff-colored precipitate ^[39].

D. Sodium Hydroxide

A tiny amount of the extract was dissolved in water and filtered. To give the filtrate a yellow coloration, 2ml of a 10% sodium hydroxide aqueous solution was added. When strong hydrochloric acid was added, the color changed from yellow to colorless, indicating the presence of flavonoids ^[41].

Test for Cardenolites

The extract was dissolved in ethanol in a little amount. It was then followed by the addition of concentrated tetraoxosulphate (VI) acid and 1 ml of acetic acid anhydride. The sample's color changed from pink to violet, indicating the presence of terpenoid ^[42].

i. Keller-killian's test

In 2ml of glacial acetic acid that also contained a drop of ferric chloride solution, the extract (0.5g) was dissolved. The next step was adding 1ml of concentrated tetraoxosulphate (VI) acid. A brown ring that formed during the interphase indicated the presence of the cardenolide-specific digitoxose sugar. During the acetic acid layer, a violet ring developed just below the brown ring; a greenish ring appeared just below the brown ring and slowly spread throughout this layer ^[41].

Test for Carbohydrates; general test (Molisch's test)

To the extract that had been dissolved in distilled water, a few drops of Molisch's reagent were added. Following this, 1 ml of concentrated tetraoxosulphate (VI) acid was added by the side of the test tube, forming an acid layer below the aqueous layer. After letting the mixture sit for two minutes, it was diluted with 5ml of purified water. The development of a reddish to dull violet color at the boundary between the two layers was seen as a positive test ^[41].

Test for Monosaccharide (Barfoed's test)

In distilled water, the extract (0.5g) was dissolved before being filtered. In a test tube, 1ml of the filtrate was combined with 1ml of Barfoed's reagent before being heated over a water bath for two minutes. A crimson cuprous oxide precipitate was regarded as a positive test result ^[39].

Test For free Reducing Sugars (Fehling's test)

The extract (0.2g) was filtered after being dissolved in distilled water. The presence of reducing sugars was detected by formatin, which appeared as a red cuprous oxide precipitate after heating the filtrate with 5 ml each of fehling's solutions A and B ^[41].

Test for Combined Reducing sugar's

The extract (0.2g) was boiled with 5ml of diluted hydrochloric acid to hydrolyze it, and the resultant filtrate was then neutralized with sodium hydroxide solution. It was then cooked on a water bath for two minutes with a few drops of Fehling's solution added. Because mixed sugars were present, a reddish-brown cuprous oxide precipitate formed ^[41].

Test for Ketones (Salivanoff's test)

A little amount of the extract was mixed with a few resorcinol crystals and 2 ml of strong hydrochloric acid, and the mixture was then allowed to boil for five minutes. The presence of ketones was assumed to be indicated by a red coloration ^[44].

Test for Soluble Starch

With 1ml of 5% potassium hydroxide that had been cooled and acidified with tetraoxosulphate (VI) acid, a small amount of the extract was added. It was determined that the coloring was yellowish and that soluble starch was present ^[44].

Antimicrobial Activity Assessment

Microorganism Source

Staphylococcus aureus, *Streptococcus pyogenes*, *Escherichia coli*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa*, *Corynebacterium species* and *Salmonella typhi* the only fungus utilized *Candida albicans* were used as the test organisms in this investigation. They were acquired from the university of Maiduguri Teaching Hospital's department of medical microbiology.

Anti-Microbial Test

The clinical and laboratory standards institute CLSI, ^[46] method for conducting the anti-microbial susceptibility test was used, with only minor modifications made by ^[47]. The test was performed using a stock concentration of 100 mg/ml that was created by dissolving 1 gram of crude extract into 10 ml of sterile distilled water using dilution ratios of 1: 100 and 1:500 for gram-positive bacteria and gram-negative bacteria, respectively ^[47]. The surface of sterile Petri dishes containing sterile solid nutritional agar was aseptically inoculated with around 0.5 ml of the diluted cultures. After incubating at 37°C for 24 hours, discs impregnated with the crude extract at a concentration of 5 mg were aseptically mounted on agar. The inhibitory zone was then measured in mm using a transparent meter rule. Amoxiclave (30 mg), ceflunat (30 ug), levoxine (5 ug), loxacin (5 ug), and reflatob were the standard anti-microbials utilized in the test, which was done in triplicate (5 ug)

Minimum Inhibitory Concentration (MIC)

The MIC is defined as the concentration at which there is no discernible turbidity in the test tubes. Following ^[48] earlier description, ^[47] used some magnification to obtain a concentration. For the bacteria that displayed reasonable sensitivity to the test extract, the MIC was established. The microorganism used in this test was created using broth dilution procedures. The working concentrations were made using two-fold serial dilution techniques, which varied from 0-195mg/ml to 50mg/ml using nutrient broth, and were then inoculated with 0.2ml solution of the test organism. This yielded the stocked extract concentration of 100mg/mg. After 24 hours of 37°C incubation, the tube was checked for turbidity at the lowest concentration where there was none.

Minimum Bactericidal Concentration (MBC)

MBC is the minimal bactericidal concentration was determined from Broth dilution test resulting from the MBC tubes as describe previously ^[47-48]. By inoculating the content of each test tube on a nutrient agar plate. The plate was then incubated at 37°C for 24hrs. The lowest concentration of the extract that showed no growth was noted and recorded as a minimum bactericidal concentration.

RESULTS

Qualitative phytochemical analysis

The resulting extract was evaporated to dryness in a hot air oven and then subjected to phytochemical analysis, as indicated in table 1.1, after being made from the methanolic leaf extract of dried powdered *Vernonia amygdalina*. Table 1.1 lists the different chemicals found in the crude extract.

Antibacterial sensitivity study

Table 1.2 lists the outcomes of the invitro antimicrobial screening. The results showed that the extract significantly inhibited the test organism, with *Pseudomonas aeruginosa* showing the highest activity across concentration levels (26mm at 1000mg/ml), followed by *Salmonella typhi* (9mm at 1000mg/ml) and (7mm at 800mg/ml), *Corynebacterium` species* (11mm at 600mg/ml), and *Staphylococcus* (1mm at 400mg/ml). Although not statistically comparable to that provided by the extract, the zone of inhibition created by the majority of antibiotic discs against some of the organisms was determined to be substantial in respect to those activities produced by the organisms under examination. However, inhibitory zones with widths less than 10 mm were regarded as active. This observation is consistent

with research from [47, 49]. According to the results of the minimum inhibition concentration (MIC) tests shown in Tables 1.3 and 1.4, it was found that the extract had the broadest activity against gram positive organisms (*Staphylococcus aureus*, *Staphylococcus pyogenes*, *Bacillus subtilis*, *Klebsiella pneumonia*, and *Pseudomonas aeruginosa*) at the concentrations of 200 mg/ml, 100 mg/ml, and 50mg/mg, while lowest activity was recorded against the organisms at 25mg/ml and 12.5ml. The extract exhibition considerable activity was noticed against *Candida albicans* and *E. Coli* in table 1.2. This study contradicted the conclusion of [47].

P. aeruginosa was found to have the highest sensitivity to the extract with effects at all concentrations (200-2100mg/ml), followed by *K. pneumonia*, *S. pyogenes*, *B. subtilis*, and *S. aureus*, with *Salmonella typhi* being recorded with the lowest sensitivity at only 800-1000mg/l. Table 1.2 shows zone of growth inhibition (mm) at various concentrations of the plant's (*Vernonia amygdalina*) *E. coli* and *Candida albicans* were found to be resistant over the 200–1000 mg/ml range of extract concentration.


TABLE 1.1 PHYTOCHEMICAL TEST RESULTS

S/N	TEST(S)	VALUES
1.	Test for Alkaloid	
	I. Dragendroff's reagent test	-
	II. Mayer's reagent test	-
2.	Test for cardiac glycoside	
	I. Salkowski's test	+
	II. Liebermann burchard's test	+
3.	Test for free Anthraquinone	-
4.	Test for combined Anthraquinone	-
5.	Test for Terpenoid	+
6.	Test for Cardenolites	
	I. Keller-killian's test	+
7.	Test for Saponins	
	I. Frothing test	+
8.	Test for Tannins	
	I. Ferric Chloride test	+
	II. Lead Acetate test	+
9.	Test for Flavonoids	
	I. Shinoda's test	+
	II. Ferric Chloride test	+
	III. Lead Acetate test	+
	IV. Sodium Hydroxide test	-
10.	Test for soluble Starch	+
11.	Test for phlobatannins	-
12.	Test for Carbohydrates	
	I. Molish's test	-
	II. Test for monosaccharide (barfoed's test)	+
	III. Test for free reducing sugars (fehling's test)	+
	IV. Test combined reducing sugar	+
	V. Test for ketoses	-

Key: + indicates presence, - indicates absence

According to table 1.0 above, the phytochemical analysis of *V. amygdalina* revealed the presence of cardiac glycoside, terpenoid, tannin, cardenolites, saponnins, flavonoids, and carbohydrates but the absence of alkaloid, phlobtannins, and anthraquinone.

Table 1.2 The Antimicrobial Activities of *Vernonia amygdalina* leaf Extra on micro-organism Inhibition zone



S/N	ORGANISMS	200	100	50	25	12.5
1.	<i>Staphylococcus aureus</i>	-	-	+	+	+
2.	<i>Streptococcus pyogenes</i>	-	-	-	+	+
3.	<i>Bacillus subtilis</i>	-	-	-	+	+
7.	<i>Pseudomonas aeruginosa</i>	-	-	-	-	-
8.	<i>Klebsiella pneumonia</i>	-	-	-	-	+

Key

NB. (- = Negative and + = positive)

The table above displays the antimicrobial activity of plant (*Vernonia amygdalina* leaf) extract on various organisms tested on, with the majority of results demonstrating sensitivity to the extract. *P. aeruginosa* was found to have the highest sensitivity, with effects at all concentrations (25-200mg/ml) save for the lowest (12.5), whereas *S. aureus* Only 100–200mg/ml was shown to be the sensitivity range for. *S. pyogenes*. *B. subtilis* *K. Pneumonia* exhibits moderate sensitivity to the extract, between 50 and 200 mg/ml.

Table 1.4 The Antimicrobial activities of *Vernonia amygdalina* leaf extract showing minimum Bactericidal Concentration of (mg/ml)

S/N	ORGANISMS	200	100	50	25	12.5
1.	<i>Staphylococcus aureus</i>	-	-	+	+	-
2.	<i>Streptococcus pyogenes</i>	-	-	-	+	-
3.	<i>Bacillus subtilis</i>	-	-	+	+	+
7.	<i>Pseudomonas aeruginosa</i>	-	-	-	+	+
8.	<i>Klebsiella pneumonia</i>	-	-	-	+	+

Key

NB. (- = Negative and + = positive)

Fig. 1.0

The minimum inhibitory concentration of the plant's (*Vernonia amygdalina* leaf) extract on the various organisms tested is shown in Table 1.3. The majority of these organisms exhibit sensitivity to the extract, with *P. aeruginosa* having the highest sensitivity and showing effects at all concentrations (25-200mg/ml) except for the lowest (12.5) and *S. aureus* having the lowest sensitivity. Only 100–200 mg/ml of aureus was demonstrated to be sensitive. *S. Pyogenes*, *B. subtilis*, *K. Pneumonia* exhibits moderate sensitivity to the extract, between 50 and 200 mg/ml.

DISCUSSION

The results obtained from the phytochemical analysis of methanolic leaf extract of *Vernonia amygdalina* demonstrated the presence of the following; tannins, saponins, flavonoids, soluble starch, carbohydrates, terpenoids, cardenolites, and cardiac glycosides. These findings are in accord with those of Alara and his colleagues^[49-54] (Table 1.1). The absence of Alkaloids and anthraquinones contradicts the findings of Usunomena and Ngozi^[55], that reported the presence of these bioactive metabolites. This could be as a result of solvent used for the study, geographical location, processing and extraction methods (Table 1.1). The presence of phytochemical components may be responsible for the observed antimicrobial activity of the leaf extract. These findings conform to the report of Ojmelukwe and Amaechi^[9]. When the extract was tested using comparative test methods by impregnated disc gel diffusion protocol against a number of microorganisms, such as *Staphylococcus aureus*, *Streptococcus pyogenes*, *Klebsiella pneumonia*, *Pseudomonas aeruginosa*, *Corynebacterium species* and *Salmonella typhi*. A specific pattern regarding the effect of the extract on inhibitory zone radii was discernible, except for *Escherichia coli* and *Candida albicans*. the extract was deployed in a range of concentrations from 200mg/ml to 1000mg/ml. it was discovered that 1000mg/ml, 800mg/ml and 600mg/ml demonstrated highest inhibition radii in the increasing order of concentration against the test organisms, thus indicating dose dependent effect and antimicrobial potency. This finding is in accord with Ibrahim and colleagues^[56]. Lower concentrations of the extract are 400mg/ml and below produced lower zone of inhibition radii significantly less than extract at 1000mg/ml and 800mg/ml. At lower concentration, these observations tend to imply that the extract antimicrobial activity increased as the concentration increased in a dose dependent manner. *Candida albicans* and *Escherichia coli*, on the other hand, demonstrated zero activity, this is in accord with the study by Oshim and colleagues^[57], that demonstrated the resistance of *Candida albicans* to the extract at all concentrations. This implies that the two organisms are resistance to the extract at all concentrations. The sensitivity test against the microorganisms was demonstrated in the invitro as shown in table 1.2.

Pseudomonas aeruginosa, which showed the highest sensitivity (26mm at 1000mg/ml) with effect at all concentrations, it was established to have significant levels of inhibition from the extract among the test organisms (20 mm at 600 mg/ml and 23 mm at 800 mg/ml). *Salmonella typhi* demonstrated the lowest levels of inhibition from the extract among the test organisms with inhibition radii, 9mm at 1000mg/ml and 7mm at 800mg/ml, when tested at all concentrations (200-1000 mg/ml), *S. aureus*, *S. pyogenes*, *B. subtilis*, *Corynebacterium species*, and *K. pneumonia* showed greater sensitivity than *Salmonella typhi* but less sensitivity to *P. aeruginosa*. The antibiotic disc's zone of inhibition against the tested organism was determined to be notable in respect to the study's sensitivity but not statistically in comparison to the extract's zone of inhibition. However, zones of inhibition with a diameter of 10 mm were thought to be active. This observation is consistent with research from Ngatu and colleagues^[13]; Dumas and colleagues^[58] which showed that the extracts of *V. amygdalina* exhibited inhibitory activity on all tested bacteria including *Staphylococcus aureus*, *Salmonella enterica* and *Klebsiella pneumoniae*^[59]. Habtamu and Melaku^[60], demonstrated the chloroform extract of *V. amygdalina* showed strong activity against *S. aureus* with an inhibition zone of 21 mm. This is in contrary to the findings in the current study which demonstrated higher activity of the extract to *Pseudomonas aeruginosa*, Followed by other organisms. Chukwuemeka and colleagues^[61], showed that the extract inhibited *S. aureus*, *Bacillus subtilis*, *Salmonella typhi* and *Pseudomonas*

aeruginosa activities in mice, which is in accord to this study. Based on the outcomes of the minimum inhibition concentration (MIC) and minimum bactericidal concentration (MBC) results, respectively, shown in Tables 1.3 and 1.4. *S. aureus* was found to be sensitive at only 100-200mg/ml, however *P. aeruginosa* was found to have the highest sensitivity with effect at all concentrations (25-200mg/ml) extract for the lowest (12.5). The moderate sensitivity of *S. pyogenes*, *B. subtilis*, and *K. pneumonia* ranges from 50 to 200 mg.

CONCLUSION

This study highlighted the antimicrobial effects of *V. amygdalina* leave extracts on some pathogens thereby verifying the traditional healer's claim. Also, it was concluded that the extract of *V. amygdalina* contain pharmacologically active phytochemicals which could be responsible for the numerous medicinal properties exhibited by the plant leaf extract and has a lot of potential as an antibacterial agent for the treatment of infection-causing pathogens. Further work should be carried out in order to isolate the active compounds for further antimicrobial, pharmacological and clinical testing.

RECOMMENDATION

Following the study's findings, it is advised that additional extensive research be encouraged to learn more about the plant's therapeutic characteristics. This research can involve qualitative and quantitative analysis as well as a look at the mineral makeup of the entire plant.

REFERENCES

1. Vaou, N., Stavropoulou, E., Voidarou, C., Tsigalou, C., & Bezirtzoglou, E. (2021). Towards advances in medicinal plant antimicrobial activity: A review study on challenges and future perspectives. *Microorganisms*, 9(10), 2041.

2. Veiga Junior, V. F., Pinto, A. C., & Maciel, M. A. M. (2005). Medicinal plants: safe cure?. *Quimica Nova*, 28, 519-528.
3. Alvim, N. A. T., Ferreira, M. D. A., Cabral, I. E., & Almeida Filho, A. J. D. (2006). O uso de plantas medicinais como recurso terapêutico: das influências da formação profissional às implicações éticas e legais de sua aplicabilidade como extensão da prática de cuidar realizada pela enfermeira. *Revista Latino-americana de enfermagem*, 14, 316-323.
4. Nowak, J., Kiss, A. K., Wambebe, C., Katuura, E., & Kuźma, Ł. (2022). Phytochemical Analysis of Polyphenols in Leaf Extract from Vernonia amygdalina Delile Plant Growing in Uganda. *Applied Sciences*, 12(2), 912.
5. Zhang, L., Wang, X., Guo, J., Xia, Q., Zhao, G., Zhou, H., & Xie, F. (2013). Metabolic profiling of Chinese tobacco leaf of different geographical origins by GC-MS. *Journal of agricultural and food chemistry*, 61(11), 2597-2605.
6. Ojmelukwe, P. C., & Amaechi, N. (2019). Composition of Vernonia amygdalina and its potential health benefits. *International Journal of Environment, Agriculture and Biotechnology*, 4(6), 1836-1848.
7. Bonsi, M. L. K., Osuji, P. O., Tuah, A. K., & Umunna, N. N. (1995). Vernonia amygdalina as a supplement to teff straw (Eragrostis tef) fed to Ethiopian Menz sheep. *Agroforestry systems*, 31(3), 229-241.
8. Akah, P. A., & Okafor, C. L. (1992). Blood sugar lowering effect of Vernonia amygdalina Del, in an experimental rabbit model. *Phytotherapy research*, 6(3), 171-173
9. Ojmelukwe, P. C., & Amaechi, N. (2019). Composition of Vernonia amygdalina and its potential health benefits. *International Journal of Environment, Agriculture and Biotechnology*, 4(6), 1836-1848.
10. Alara, O. R., Abdurahman, N. H., Ukaegbu, C. I., & Kabbashi, N. A. (2019). Extraction and characterization of bioactive compounds in Vernonia amygdalina leaf ethanolic extract comparing Soxhlet and microwave-assisted extraction techniques. *Journal of Taibah University for Science*, 13(1), 414-422.
11. Alara, O. R., Abdurahman, N. H., Ukaegbu, C. I., Hassan, Z., & Kabbashi, N. A. (2018). Dataset on LC-Q-TOF/MS tentative identification of phytochemicals in the extract of Vernonia amygdalina leaf through positive ionization. *Data in brief*, 21, 1686-1689.
12. Swee, K. Y., Wan, Y. H., Boon, K. B., Woon, S. L., Huynh, K. Y., Abdul, H. N. Y., & Noorjahan, B. A. (2010). Vernonia amygdalina, an ethnoveterinary and ethnomedical used green vegetable with multiple bio-activities. *Journal of medicinal plants research*, 4(25), 2787-2812.
13. Ngatu, N. R., Okajima, M. K., Yokogawa, M., Hirota, R., Takaishi, M., Eitoku, M., ... & Suganuma, N. (2012). Anti-allergic effects of Vernonia amygdalina leaf extracts in hapten-induced atopic dermatitis-like disease in mice. *Allergology International*, 61(4), 597-607.
14. Alo, M. N., Anyim, C., Igwe, J. C., Elom, M., & Uchenna, D. S. (2012). Antibacterial activity of water, ethanol and methanol extracts of Ocimum gratissimum, Vernonia amygdalina and Aframomum melegueta. *Adv. Appl. Sci. Res*, 3(2), 844-848.
15. Adetunji, C. O., Olaniyi, O. O., & Ogunkunle, A. T. J. (2013). Bacterial activity of crude extracts of Vernonia amygdalina on clinical isolates. *Journal of Microbiology and Antimicrobials*, 5(6), 60-64.
16. Akinyele, B. J., Oladejo, B. O., Akinyemi, A. I., & Ezem, L. O. (2014). Comparative study of the antibacterial effect of mouth washes and Vernonia amygdalina (del.) on some tooth decay causing bacteria. *British Microbiology Research Journal*, 4(7), 749.
17. Kupchan, S. M., Hemingway, R. J., Karim, A., & Werner, D. (1969). Tumor inhibitors. XLVII. Vernodaline and vernomygdin, two new cytotoxic sesquiterpene lactones from Vernonia amygdalina Del. *The Journal of organic chemistry*, 34(12), 3908-3911.
18. Owoeye, O., Yousuf, S., Akhtar, M. N., Qamar, K., Dar, A., Farombi, E. O., ... & Choudhary, M. I. (2010). Another anticancer elemanolide from Vernonia amygdalina Del. *International Journal of Biological and Chemical Sciences*, 4(1).

19. Aregheore, E. M., Makkar, H. P. S., & Becker, K. (1998). Feed value of some browse plants from the central zone of Delta State, Nigeria. *Tropical Science (United Kingdom)*.
20. Adama, K. K., Oberafo, A. A., & Dika, S. I. (2011). Bitterleaf as local substitute for hops in the Nigerian brewing industry. *Archives of Applied Science Research*, 3(4), 388-397.
21. Georgewill, O. A., & Georgewill, U. O. (2010). Evaluation of the anti-inflammatory activity of extract of *Vernonia amygdalina*. *Asian Pacific Journal of Tropical Medicine*, 3(2), 150-151.
22. Udochukwu, U., Omeje, F. I., Uloma, I. S., & Oseiwe, F. D. (2015). Phytochemical analysis of *Vernonia amygdalina* and *Ocimum gratissimum* extracts and their antibacterial activity on some drug resistant bacteria. *American Journal of Research Communication*, 3(5), 225-235.
23. Igile, G., Oleszek, W., Jurzysta, M., Aquino, R., de Tommasi, N., & Pizza, C. (1995). Vemoniosides D and E, two novel saponins from *Vernonia amygdalina*. *Journal of Natural Products*, 58(9), 1438-1443.
24. Okunlola, G. O., Jimoh, M. A., Olatunji, O. A., Rufai, A. B., & Omidiran, A. O. (2019). Proximate analysis, mineral composition, and antioxidant properties of bitter leaf and scent leaf. *International journal of vegetable science*, 25(4), 346-354.
25. Clement, E., Erharuyi, O., Vincent, I., Joy, A., Christopher, A., Anthony, A., ... & Abiodun, F. (2014). Significance of bitter leaf (*vernonia amagdalina*) in tropical diseases and beyond: a review. *Malar Chemoth Cont*, 3(120), 2.
26. Atangwho, I. J., Ebong, P. E., Eteng, M. U., Eyong, E. U., & Obi, A. U. (2007). Effect of *Vernonia amygdalina* Del. leaf on kidney function of diabetic rats. *International Journal of pharmacology*, 3(2), 143-148.
27. Ross, J, Mark, M. and Sinclair, B., 2000. *Phytochemistry: Saponin*. 2nd ed. Hertfordshire library. 35– 40-
28. Tedder, M., 2003. *Carbohydrate Chemistry: starch glycoside*. Annual Review of Biochemistry. 48– 50.
29. Ekam, V. S., Ebong, P. E., & Umoh, I. B. (2010). Phytochemical screening of activity directed extracts A of *vernonia amygdalina* leaves. *Global Journal of Pure and Applied Sciences*, 16(1).
30. Martindale, J., 1978. *Physiochemistry: Effects of saponins and other substances on gastrointestinal tract*. Acta of Biochemistry and Biophysics 14 . 141 – 142.
31. Igile, G. O., Oleszek, W., Burda, S., & Jurzysta, M. (1995). Nutritional assessment of *Vernonia amygdalina* leaves in growing mice. *Journal of Agricultural and Food Chemistry*, 43(8), 2162-2166.
32. Huang, I. and Ferraro, F., 1992. *Chemistry: Flavonoids chemistry*. 2nd ed. Cold spring harbor laboratory press, Cold spring harbor. NY. 77 – 78.
33. Ibrahim, T. A., Lola, A., Adetuyi, F. O., & Jude-Ojei, B. (2009). Assessment of the antibacterial activity of *Vernonia amygdalina* and *Occimum gratissimum* leaves on selected food borne pathogens. *Electronic Journal of Environmental, Agricultural and Food Chemistry*, 8(11), 1213-1217.
34. Anna, K. Q., Yakubu, J., Arnold, F. D., Isaac, A., Adwoa, A. O. K., Sussana, E. R. A., & Patience, N. (2020). Hydroethanolic stem bark extract of *Vernonia amygdalina* Del.(Asteraceae) suppresses yeast-induced pyrexia and *Plasmodium berghei* malaria in murine models. *Journal of Medicinal Plants Research*, 14(6), 258-264.
35. Tijjani, M. A., Mohammed, G. T., Alkali, Y. T., Adamu, T. B., & Abdurahaman, F. I. (2017). Phytochemical analysis, analgesic and antipyretic properties of ethanolic leaf extract of *Vernonia amygdalina* Del. *Journal of Herbmed Pharmacology*, 6(3), 95-99.
36. Mayomi, I., & Mohammed, J. A. (2014). A decade assessments of Maiduguri urban expansion (2002-2012): Geospatial approach. *Global J. Human Sci. B*, 14(2), 1-0.
37. Odebiyi, O. O., & Sofowora, E. A. (1978). Phytochemical screening of Nigerian medicinal plants II. *Lloydia*, 41(3), 234-246.

38. Evans, W. C. (2009). *Trease and Evans' pharmacognosy*. Elsevier Health Sciences.
39. Brain K.R. and Turner T.D. (1975). The practical evaluation of phytopharmaceuticals (Vol.1). Bristol: wright-scientifica. Pp. 187-189.
40. Sofowora A., (1993). Screening plants for bioactive agents. In: medicinal plants and traditional medicinal in Africa 2nd Ed. Spectrum books ltd, Sunshine house, Ibadan, Nigeria, Pp 134-156.
41. Evans, W. C. (2002). Trease and evans. *Pharmacognosy, 9th Edition published by Saunders Elsevier, 553*.
42. Silva L.G., Lee I.S., and Afkinnghorn D.A., (1998). Special problem with extraction of plant: In natural products isolation (Ed: cannell, RJD). Human press Inc. 999, review drive, suit 208, Totowa, New jersey.
43. Harborne, J. B. (1973). *A guide to modern techniques of plant analysis*. Chapman and hall.
44. Vishnoi N.R. (1979). Advanced practical chemistry. Yikas publication house, PVT ltd, Ghaziabad-india. Pp. 447-449.
45. Markham K.R. (1982). Techniques of flavonoid identification. Academic press Newyork, USA Pp 1-113.
46. Wayne, P. A. (2010). Clinical and laboratory standards Institute (CLSI); 2010. *Performance standards for antimicrobial susceptibility testing, 20*, 1-5.
47. Usman, H., Haruna, A. K., Akpulu, I. N., Ilyas, M., Ahmadu, A. A., & Musa, Y. M. (2005). Phytochemical and antimicrobial screenings of the leaf extracts of *Celtis integrifolia* Lam. *J. Trop. Biosci*, 5(2), 72-76.
48. Vollekova, A., Košťálová, D., & Sochorova, R. (2001). Isoquinoline alkaloids from *Mahonia aquifolium* stem bark are active against *Malassezia* spp. *Folia microbiologica*, 46(2), 107-111.
49. Alara, OR, Abdurahman, NH, Mudalipa, SKA, Olalere, OA., 2017. Phytochemical and pharmacological properties of *Vernonia amygdalina*: a review. *Journal of Chemical Engineering and Industrial Biotechnology* 2, 80–96. doi: 10.15282/JCEIB-V2-07.29/9/2017/2.2 .
50. Erasto, P., Grierson, D. S., & Afolayan, A. J. (2006). Bioactive sesquiterpene lactones from the leaves of *Vernonia amygdalina*. *Journal of ethnopharmacology*, 106(1), 117-120.
51. Farombi, E. O., & Owoeye, O. (2011). Antioxidative and chemopreventive properties of *Vernonia amygdalina* and *Garcinia* biflavonoid. *International journal of environmental research and public health*, 8(6), 2533-2555.
52. Toyang, N. J., & Verpoorte, R. (2013). A review of the medicinal potentials of plants of the genus *Vernonia* (Asteraceae). *Journal of Ethnopharmacology*, 146(3), 681-723.
53. Quasie, O., Zhang, Y. M., Zhang, H. J., Luo, J., & Kong, L. Y. (2016). Four new steroid saponins with highly oxidized side chains from the leaves of *Vernonia amygdalina*. *Phytochemistry Letters*, 15, 16-20.
54. Luo, X., Jiang, Y., Fronczek, F.R., Lin, C., Izevbigie, E.B., Lee, S. and Lee, K.S. 2017. Isolation and Structure Determination of a Sesquiterpene Lactone (Vernodalinol) from *Vernonia amygdalina* Extracts. *Pharm Biol* 49 (5):464–470. [https://doi: 10.3109/13880209.2010.523429](https://doi.org/10.3109/13880209.2010.523429).
55. Usunomena, U., & Ngozi, O. P. (2016). Phytochemical analysis and proximate composition of *Vernonia amygdalina*. *Int. J. Sci. World*, 4, 11-14.
56. Ibrahim, N. D. G., Abdurahman, E. M., & Ibrahim, G. (2001). Elemental analysis of the leaves of *Vernonia amygdalina* and its biological evaluation in rats. *Nigerian Journal of Natural Products and Medicine*, 5, 13.
57. Oshim, I. O., Desmond, C. O., Nwobu, R. A. U., Ezugwu, U. M., & Urama, E. U. (2016). Kinetics of minimum inhibitory concentration, minimum bactericidal concentration and minimum fungicidal concentration of *Vernonia amygdalina* (Bitter leaf) on microorganisms isolated from wound infections. *International Journal of Surgical Research*, 5(1), 8-14.

58. Dumas, N. G. E., Anderson, N. T. Y., Godswill, N. N., Thiruvengadam, M., Ana-Maria, G., Ramona, P., ... & Emmanuel, Y. (2021). Secondary metabolite contents and antimicrobial activity of leaf extracts reveal genetic variability of *Vernonia amygdalina* and *Vernonia calvoana* morphotypes. *Biotechnology and Applied Biochemistry*, 68(4), 938-947.
59. Zwadyk, P. (1972). Enteriobacteriaceae in zinsser microbiology.
60. Habtamu, A., & Melaku, Y. (2018). Antibacterial and antioxidant compounds from the flower extracts of *Vernonia amygdalina*. *Advances in Pharmacological sciences*, 2018.
61. Chukwuemeka, N. O., Oyebisi, S. O., Adebisi Oluwadare, O. J., Oladapo, R. S., & Berka, N. P. (2018). Antibacterial assay and reversion of carbon tetrachloride induced liver damage on wistar mice by *Vernonia amygdalina*. Delile. *Pakistan journal of pharmaceutical sciences*, 31(4).