

# INVESTIGATING THE EFFECT OF VITAMIN E AND C COMBINED TREATMENT ON PARAQUAT-INDUCED LIVER DAMAGE IN WISTAR RATS (*RATTUS NOVERGICUS*)

## Abstract

Paraquat has been referred to as a prominent suicide agent in a number of nations, not only because it is highly acutely poisonous, but also because it is readily available, relatively inexpensive, and has no known antidote. Vitamins ~~C-E~~ and ~~CE~~ are well-known antioxidants that react quickly with free radicals and prevent lipid peroxidation. The aim of this study was to determine the ameliorative effect of vitamin E and C in paraquat induced liver poisoning in rats. A total of 200 male albino rats were used in the study. With 50 rats each, the rats were sorted into four main groups: A, B, C, and D. Each group was divided into two subgroups (0 and VEC), each with 25 rats. The "A" group received no paraquat, while the "B," "C," and "D" groups received 0.02g, 0.04g, and 0.06g of paraquat respectively. All "0" subgroups represented those not treated with Vit E and C combination therapy and those treated with 500mg of vitamin E and 2000mg/dl of vitamin C designated VEC. The paraquat induction frequency was every two weeks for three months, followed by two months of weekly vitamin treatments. Blood was drawn for SGPT, SGOT, ALP, and GGT testing. At p-value 0.05, there was a significant difference in these liver enzymes activities among the "A0", "B0", "C0", and "D0". There was a significant difference in these liver enzymes activities among the "AVEC," "BVEC," "CVEC," and "DVEC," p-value 0.05. As a result, significant differences in intra-group comparisons of SGPT, SGOT, ALP, and GGT p-value 0.05 were observed. This study discovered that combining vitamin E and C had an ameliorative effect on the liver as a result of PQ insult.

**Comment [PM1]:** May be change numerotation to avoid confusion with VitC

**Keyword:** Paraquat, Vitamin E, Vitamin C, rat, antioxidant

## 1.0 INTRODUCTION

Paraquat (PQ) is a man-made substance. Weidel and Russo were the first to describe it in 1882. Paraquat's redox characteristics were found by Michealis and Hill in 1933, and the chemical was given the name methyl viologen. Paraquat (PQ) was first recognized as having herbicidal qualities in 1958, and it became commercially available in 1962 [1]. It has been dubbed a major suicide agent in several nations, not only because it is highly acutely poisonous, but also because it is widely available, relatively inexpensive, and has no known antidote [2]. The majority of research on paraquat toxicity has focused on reducing its absorption or improving its elimination [3]. Currently, there are no effective pharmacological antagonists for paraquat and there are no chelating agents capable of binding the poison in blood or other tissues, and so the management of paraquat poisoning has remained mostly supportive and directed toward decreasing its absorption or enhancing its elimination [3].

**Comment [PM2]:** Must be improve. It is necessary to provide more recent references to highlight the interest of the study presented here.

PQ is converted to PQ mono-cation free radical by nicotinamide adenine dinucleotide phosphate (NADPH)-cytochrome P-450 reductase once it enters the cells. The electron delivered to PQ quickly travels to oxygen, resulting in the formation of superoxide anion, and [reactive free oxygen radicals \(ROS\)](#) causes severe oxidative damage [3]. The generation of ~~reactive free oxygen radicals (ROS)~~, which induce oxidative stress and lipid peroxidation ([LPO](#)), is a common underlying mechanism of most hepatotoxicity-inducing contaminants (~~LPO~~). Damage to hepatocyte membranes results in edema, degeneration, necrosis, and fibrosis of hepatocytes as a result of the cumulative consequences of these events [4,5]. ROS and other free radicals are continuously created during normal physiologic processes and damage macromolecules such as proteins, lipids, and DNA, causing tissue harm [6]. The importance of oxidative stress in the etiology and progression of chronic liver disease has long been recognized [7,8]. Antioxidant compounds have previously been shown to protect cells from the damaging effects of a variety of environmental pollutants [9,10].

Comment [PM3]: Add scheme

Vitamins C and E are well-known antioxidants that react quickly with free radicals and prevent lipid peroxidation. As a result, both vitamins C and E are thought to be absorbed during radical reactions [11,12]. For paraquat poisoning, increasing total antioxidant status with high dosages of vitamin C is beneficial as a free radical scavenger. Vitamin C's capacity to quench radicals generated by paraquat redox cycling has been credited with these protective benefits [13]. Although many antioxidants work in concert, ascorbic acid (vitamin C) has the ability to destroy or reactivate other antioxidants [14]. Vitamin C may directly scavenge oxygen free radicals with or without enzyme catalysts, and it can also scavenge them indirectly by recycling others to their reduced state [15].

Vitamin E deficiency potentiated the development of acute paraquat toxicity in rats in multiple experiments, demonstrating its function in paraquat toxicity. One of the most well-known processes is its ability to eliminate free radicals produced by the human body's oxidation response or by foreign substances. *In vitro* and *in vivo*, it also suppresses prostaglandin synthesis and prevents platelet aggregation [16].

Formatted: Font: Italic

Formatted: Font: Italic

Antioxidant systems, in general, either prevent ROS from forming or eliminate them before they can harm essential cell components [17].

Glutamate-oxaloacetate-aminotransferase (SGOT), also known as aspartate amino transferase (AST), is an enzyme that catalyzes transamination reaction. It is found in the liver, heart, skeletal muscle, and kidney [18]. Elevations frequently predominate in cirrhosis and even in liver diseases that typically have an increased [Alanine amino transferase \(ALT\)](#) [19].

Glutarate-pyruvic-aminotransferase (SGPT), also known as Alanine amino transferase (ALT), is an enzyme found in higher concentrations in the liver than in other tissues of the body [18]. Any type of liver cell injury can cause an increase in ALT levels.

Alkaline phosphatase (ALP) is found in the small intestine's mucosal epithelia, the kidney's proximal convoluted tubule, bone, the liver, and the placenta. It is responsible for lipid transport in the intestine and bone calcification. Its activity is primarily derived from the liver, with bone contributing 50% [18].

(GGT) is a microsomal enzyme found in hepatocytes and biliary epithelial cells, as well as renal tubules, the pancreas, and the intestine. It is also found in the cell membrane, where it transports peptides into the cell and is involved in glutathione metabolism [18]. Because of the relative importance of vitamin C and E as antioxidants, this study focused on how to boost the antioxidant system in the body to counteract the harmful effects of paraquat. Because the liver is the body's energy store and source, the liver enzymes SGOT, SGPT, ALP, and GGT were chosen

Comment [PM4]: ?

## 2.0 MATERIAL AND METHODS

### 2.1 list 1 :Study Design

Table 1 name....

Total No-of Rats	Groups	Subgroups (rats)	Treatment
32 Rats	A (50 rats)	A <sub>0</sub> (25 rats)	Not induced with paraquat and no treatment with Vitamin E and C given
		A <sub>VEC</sub> (25 rats)	Not induced with paraquat but treated with Vitamin E(500mg) and C(2000mg/dl) given
		B <sub>0</sub> (25 rats)	Induced with 0.02gc of paraquat (3 months) and no Vitamin E and C treatment given
	B (50 rats)	B <sub>VEC</sub> (25 rats)	Induced with 0.02gc of paraquat (3 months) and Vitamin E(500mg) and C(2000mg/dl) treatment given(for 2 months after PQ induction)
		C <sub>0</sub> (25 rats)	Induced with 0.04gc of paraquat (3 months) with no Vitamin E and C treatment given
		C <sub>VEC</sub> (25 rats)	Induced with 0.04gc of paraquat (3 months) and Vitamin E and C treatment given

Comment [PM5]: explain please

Formatted: Centered, Space After: 0 pt

Formatted Table

Formatted: Centered, Space After: 0 pt

Formatted: Centered, Space After: 0 pt

Formatted: Centered, None, Space Before: 0 pt, Don't keep with next, Don't keep lines together

Comment [PM6]: ??

Formatted: Centered, None, Space Before: 0 pt, Don't keep with next, Don't keep lines together, Border: Top: (Single solid line, Auto, 0.5 pt Line width)

Formatted: Centered, Space After: 0 pt

Formatted: Centered, None, Space Before: 0 pt, Don't keep with next, Don't keep lines together

Formatted: Centered, None, Space Before: 0 pt, Don't keep with next, Don't keep lines together, Border: Top: (Single solid line, Auto, 0.5 pt Line width)

		(25 rats)	Vitamin E(500mg) and C(2000mg/dl) given (for 2 months after PQ induction elapsed)	Formatted: Centered, None, Space Before: 0 pt, Don't keep with next, Don't keep lines together
	C(50 rats)	D <sub>0</sub> (25 rats)	Induced with 0.06gc of paraquat (3 months) with no Vitamin E and C treatment given	Formatted: Centered, None, Space Before: 0 pt, Don't keep with next, Don't keep lines together, Border: Top: (Single solid line, Auto, 0.5 pt Line width)
		D <sub>VEC</sub> (25 rats)	Induced with 0.06gc of paraquat(3 months) and Vitamin E(500mg) and C(2000mg/dl) given (for 2 months after PQ induction elapsed)	Formatted: Centered, None, Space Before: 0 pt, Don't keep with next, Don't keep lines together
	D(50 rats)			Formatted: Centered, Space After: 0 pt
				Formatted: Centered, None, Space Before: 0 pt, Don't keep with next, Don't keep lines together

## 2.2 Animal source

The Animal House, Rivers State University of Science and Technology's Department of Biology, provided 200 rats weighing an average of 0.20±0.02kg. The rats were brought to the research site and given two weeks to adapt before the trial began. The study was conducted at the Department of Medical Laboratory Science at Rivers State University of Science and Technology.

Comment [PM7]: necessary to have carried out the study with such a sample?

### Treatment Administration

#### Procedure for Paraquat Administration

Administration of toxicant was *via* oral gavage route. The dose depended on the treatment group but in all, the treatment was performed every two weeks for three months.

Formatted: Font: Italic

The rats were held at the skin over the head and turned so that the mouth was faced upward and the body lowered towards the holder. The syringe needle bevel was then placed into the mouth of the rat a bit laterally in a way to avoid the teeth which are located centrally. The content in the syringe was then emptied into the mouth of the rat gradually [20].

#### Procedure for Vitamin Administration:

The vitamins (E and C) were given orally every week for two months at doses of 500mg and 2000mg/dl respectively [20].

### 2.3 Sample Collection method


Blood was collected *via* cardiac puncture into plain bottles (without anticoagulant) with a 2ml syringe, allowed to clot to obtain serum, and then analyzed for the liver enzymes GGT, SGOT, SGPT, and ALP. The animals were sacrificed under 70% ~~percent~~ chloroform anesthesia. To avoid pollution, the carcasses that remained were incinerated.

Formatted: Font: Italic

### 2.4 Laboratory analysis

#### 2.4.1 Serum glutarate-oxaloacetate-aminotransferase (AST/SGOT) method: by Reitman and Frankel, 1957.

Principle: AST is incubated at 37<sup>o</sup>C for exactly 60 minutes in a pH 7.5 buffered substrate containing aspartate and  $\alpha$ -ketoglutarate. AST catalyzes the transfer of the amino group from aspartate to ketoglutarate, forming oxaloacetate and glutamate. The oxaloacetate reacts with 2,4-dinitrophenylhydrazine to form 2,4-dinitrophenylhydrazone which in alkaline medium gives a red-brown colour. The absorbance of the colour produced is measured at 540nm Spectrophotometrically



Comment [PM8]: Structure ?

Procedure: In estimating the activity of this enzyme, 0.5ml of buffered-L- aspartate and  $\alpha$ -oxoglutarate solution was each added to two glass tubes labeled 'Reagent Blank' and 'Test', followed by 0.1ml each of distilled water and sample to their respective tubes, mixed and incubated for exactly 30 minutes at 37<sup>o</sup>C. After which, 0.5ml of 2,4-dinitrophenylhydrazine (2mmol/L) solution was added to each of the test tubes, mixed again and allowed to stand for exactly 20 minutes, at 20 – 25<sup>o</sup>C. At the end of the time, 5.0mls of Sodium hydroxide (0.4mol/L) was added to enhance colour development at alkaline pH. The tubes were mixed and the absorbance of 'Test' ( $A_{\text{test}}$ ) read against that of the 'Reagent blank' after 5 minutes.

Calculation: Obtain the activity of the enzyme AST in the serum from the table of values previously plotted against activities. Haemolysis interferes with the assay.

#### 2.4.2 Serum glutarate-pyruvic-aminotransferase (SGPT) by Reitman and Frankel, 1957.

Comment [PM9]: Group 2.4.2 and 2.4.1

Principle: ALT is incubated at 37<sup>o</sup>C for exactly 60 minutes in a pH 7.5 buffered substrate containing L-alanine and  $\alpha$ -ketoglutarate. ALT catalyzes the transfer of the amino group from alanine to ketoglutarate, forming pyruvate and glutamate. The pyruvate reacts with 2,4-dinitrophenylhydrazine to form 2,4-dinitrophenylhydrazone which in alkaline medium gives a red-brown colour. The absorbance of the ~~colour~~ produced is measured at 540nm Spectrophotometrically



Comment [PM10]: Structure?

Procedure: In estimating the activity of this enzyme, 0.5ml of buffered-L-alanine and  $\alpha$ -oxoglutarate solution was each added to two glass tubes labeled 'Reagent Blank' and 'Test', followed by 0.1ml each of distilled water and sample to their respective tubes, mixed and incubated for exactly 30 minutes at 37<sup>o</sup>C. After which, 0.5ml of 2,4-dinitrophenylhydrazine (2mmol/L) solution was added to each of the test tubes, mixed again and allowed to stand for exactly 20 minutes, at 20 – 25<sup>o</sup>C. At the end of the time, 5.0mls of Sodium hydroxide (0.4mol/L) was added to enhance ~~colour~~ development at alkaline pH. The tubes were mixed and the absorbance of 'Test' ( $A_{\text{test}}$ ) read against that of the 'Reagent blank' after 5 minutes.

Calculation: Obtain the activity of the enzyme ALT in the serum from the table of values previously plotted against activities. Haemolysis interferes with the assay.

### 2.4.3 Alkaline phosphatase (ALP) method by Englehardt, *et al.*, 1970.

Principle: This is an optimized standard method where ALP is measured by monitoring the concentration of *p*-nitrophenol formed with *p*-nitrophenlphosphate.

Formatted: Font: Italic

Formatted: Font: Italic

Procedure: Fresh double distilled water (ddH<sub>2</sub>O) was aspirated and used to perform a new Gain calibration in flow cell mode. This zero the equipment from previous sample run. ALP was selected in the Run Test Screen and a water blank test run was carried out, after which 0.02ml of sample and 1.0 ml of reagent (Diethanolamine buffered *p*-nitrophenylphosphate) was dispensed into a test tube and mixed for 2 minutes. The mixture was then aspirated into the Rx Monza. After about 2 minutes the result of the test sample was then printed out from a printer connected to the machine.

Formatted: Font: Italic

The advantage of this machine procedure is that up to 200 samples can be processed and results produced within one hour in S.I. unit = IU/L.

Manual calculation: To calculate the ALP activity, using the manual method, the following formula was utilized:



$$\text{IU/L} = 2760 \times \Delta A \text{ 405 nm/min.}$$

Formatted: Centered

#### 2.4.4 Gamma-Glutamyltransferase (GGT) method: (Szasz and Bergmeyer, 1974 and Teitz, 1987)

##### Principle

The substrate L-V-glutamyl-3-carboxy-4-nitroanilide, in the presence of glycylglycine is converted by GGT in the sample to 5-amino-2-nitrobenzoate, which can be measured spectrophotometrically at 405nm.



Comment [PM11]: Structure and reformulate

Procedure: 0.1ml of sample and 1.0ml of reagent (Buffered Glycylglycerine and L-gammaglutamyl-3-carboxy-4-nitrolide) were dispensed into a cuvette, mixed and the initial absorbance read at 400 – 420nm with simultaneous timer initiation. The absorbance was read again after 1, 2 and 3 minutes.

Calculation:  $\text{IU/L} = 1158 \times \Delta A \text{ (405nm/minute)}$ .

#### 2.5 Statistical analysis

The data generated from this study was analyzed using SPSS version 23.0 for descriptive and inferential statistics (ANOVA) for inter-group comparison and T-test for intra-group (sub-group) comparison at test significance, P-value<0.05.

### 3.0 RESULTS

Comment [PM12]: Must be improve. The results part cannot be a sequence of tables. Authors should present the results and explain what the tables represent.

Comment [PM13]: You must introduce results

Table 1-2: Changes in Liver enzymes biochemical data after two months treatment period.

Subgroup	SGOT (IU/L)	SGPT (IU/L)	ALP (IU/L)	GGT (IU/L)
A <sub>0</sub>	6.17 ± 1.38	4.18 ± 0.03	1.57 ± 0.05	13.33 ± 0.17

A <sub>VEC</sub>	4.12 ± 1.39	4.33 ± 0.06	1.55 ± 0.01	8.48 ± 0.47
B <sub>0</sub>	15.24 ± 1.39 <sup>a</sup>	41.15 ± 0.44 <sup>a</sup>	100.30 ± 1.00 <sup>a</sup>	63.00 ± 1.39 <sup>a</sup>
B <sub>VEC</sub>	19.72 ± 1.35 <sup>a,b</sup>	34.25 ± 0.38 <sup>a,b</sup>	66.03 ± 1.14 <sup>a,b</sup>	35.98 ± 0.48 <sup>a,b</sup>
C <sub>0</sub>	99.65 ± 1.64 <sup>a</sup>	147.00 ± 1.52 <sup>a</sup>	130.68 ± 0.84 <sup>a</sup>	76.25 ± 1.30 <sup>a</sup>
C <sub>VEC</sub>	74.60 ± 2.02 <sup>a,b</sup>	91.50 ± 0.64 <sup>a,b</sup>	80.38 ± 1.84 <sup>a,b</sup>	42.75 ± 0.46 <sup>a,b</sup>
D <sub>0</sub>	193.00 ± 2.70 <sup>a</sup>	258.75 ± 3.10 <sup>a</sup>	315.53 ± 1.63 <sup>a</sup>	99.00 ± 0.60 <sup>a</sup>
D <sub>VEC</sub>	238.75 ± 3.95 <sup>a,b</sup>	111.25 ± 2.86 <sup>a,b</sup>	164.40 ± 5.83 <sup>a,b</sup>	70.25 ± 1.86 <sup>a,b</sup>

Statistical significance: P ≤ 0.05.

➤—Index (a) = represents a statistically significant difference between the test subgroups and the control subgroups at each treatment month.

➤

➤ Index (b) = represents a statistically significant difference observed within each group (i.e. Group B: B<sub>0</sub> Vs B<sub>VE</sub>) at each month.

Formatted: Bulleted + Level: 1 + Aligned at: 0.25" + Indent at: 0.5"

A<sub>0</sub> – Not induced with paraquat and no treatment with Vitamin E and C given

A<sub>VEC</sub> – Not induced with paraquat but Vitamin E and C treatment given

B<sub>0</sub> – Induced with 0.02gc of paraquat and no Vitamin E and C treatment given

B<sub>VEC</sub> – Induced with 0.02gc of paraquat and Vitamin E and C treatment given

C<sub>0</sub> – Induced with 0.04gc of paraquat with no Vitamin E and C treatment given

C<sub>VEC</sub> – Induced with 0.04gc of paraquat with Vitamin E and C treatment given

D<sub>0</sub> – Induced with 0.06gc of paraquat with no Vitamin E and C treatment given

D<sub>VEC</sub> – Induced with 0.06gc of paraquat with Vitamin E and C treatment given

Thus, from the analysis,

Comment [PM14]: ????

**Table 32: SGOT (IU/L)** – Extrapolated table illustrating simplified statistical interpretation (Vit E + C therapy) at P ≤ 0.05.

GROUPS	DECISION
A <sub>0</sub>	6.17 ± 1.38
B <sub>0</sub>	15.24 ± 1.39 <sup>a</sup> - Significant difference
C <sub>0</sub>	99.65 ± 1.64 <sup>a</sup> - Significant difference
D <sub>0</sub>	193.00 ± 2.70 <sup>a</sup> - Significant difference
A <sub>VEC</sub>	4.12 ± 1.39
B <sub>VEC</sub>	19.72 ± 1.35 <sup>a</sup> - Significant difference
C <sub>VEC</sub>	74.60 ± 2.02 <sup>a</sup> - Significant difference

D <sub>VEC</sub>	238.75 ± 3.95 <sup>a</sup> - Significant difference
B <sub>0</sub>	15.24 ± 1.39
V <sub>s</sub>	Significant difference
B <sub>VEC</sub>	
C <sub>0</sub>	99.65 ± 1.64
V <sub>s</sub>	Significant difference
C <sub>VEC</sub>	
D <sub>0</sub>	193.00 ± 2.70
V <sub>s</sub>	Significant difference
D <sub>VEC</sub>	

**Table 43: SGPT (IU/L)**- Extrapolated table illustrating simplified statistical interpretation (Vit E + C therapy) at P ≤ 0.05.

GROUPS	DECISION
A <sub>0</sub>	4.18 ± 0.03
B <sub>0</sub>	41.15 ± 0.44 <sup>a</sup> - Significant difference
C <sub>0</sub>	147.00 ± 1.52 <sup>a</sup> - Significant difference
D <sub>0</sub>	258.75 ± 3.10 <sup>a</sup> - Significant difference
A <sub>VEC</sub>	4.33 ± 0.06
B <sub>VEC</sub>	34.25 ± 0.38 <sup>a</sup> - Significant difference
C <sub>VEC</sub>	91.50 ± 0.64 <sup>a</sup> - Significant difference
D <sub>VEC</sub>	111.25 ± 2.86 <sup>a</sup> - Significant difference
B <sub>0</sub>	41.15 ± 0.44
V <sub>s</sub>	Significant difference
B <sub>VEC</sub>	
C <sub>0</sub>	147.00 ± 1.52
V <sub>s</sub>	Significant difference
C <sub>VEC</sub>	
D <sub>0</sub>	258.75 ± 3.10
V <sub>s</sub>	Significant difference
D <sub>VEC</sub>	

**Table 54: ALP (IU/L)**- Extrapolated table illustrating simplified statistical interpretation (Vit E + C therapy) at P ≤ 0.05

GROUPS	DECISION
A <sub>0</sub>	1.57 ± 0.05
B <sub>0</sub>	100.30 ± 1.00 <sup>a</sup> - Significant difference
C <sub>0</sub>	130.68 ± 0.84 <sup>a</sup> - Significant difference
D <sub>0</sub>	315.53 ± 1.63 <sup>a</sup> - Significant difference
A <sub>VEC</sub>	1.55 ± 0.01
B <sub>VEC</sub>	66.03 ± 1.14 <sup>a</sup> - Significant difference
C <sub>VEC</sub>	80.38 ± 1.84 <sup>a</sup> - Significant difference
D <sub>VEC</sub>	164.40 ± 5.83 <sup>a</sup> - Significant difference

B <sub>0</sub> Vs B <sub>VEC</sub>	100.30 ± 1.00 Significant difference 66.03 ± 1.14 <sup>b</sup>	} Significant difference
C <sub>0</sub> Vs C <sub>VEC</sub>	130.68 ± 0.84 80.38 ± 1.84 <sup>b</sup>	
D <sub>0</sub> Vs D <sub>VEC</sub>	315.53 ± 1.63 Significant difference 164.40 ± 5.83 <sup>b</sup>	

**Table 65: GGT (IU/L)** – Extrapolated table illustrating simplified statistical interpretation (Vit E + C therapy) at P ≤ 0.05

GROUPS	DECISION			
A <sub>0</sub> B <sub>0</sub> C <sub>0</sub> D <sub>0</sub>	13.33 ± 0.17 63.00 ± 1.39 <sup>a</sup> 76.25 ± 1.30 <sup>a</sup> 99.00 ± 0.60 <sup>a</sup>	} Significant difference		
A <sub>VEC</sub> B <sub>VEC</sub> C <sub>VEC</sub> D <sub>VEC</sub>	8.48 ± 0.47 35.98 ± 0.48 <sup>a</sup> 42.75 ± 0.46 <sup>a</sup> 70.25 ± 1.86 <sup>a</sup>		} Significant difference	
B <sub>0</sub> Vs B <sub>VEC</sub>	63.00 ± 1.39 35.98 ± 0.48 <sup>b</sup>			} Significant difference
C <sub>0</sub> Vs C <sub>VEC1</sub>	76.25 ± 1.30 42.75 ± 0.46 <sup>b</sup>			
D <sub>0</sub> Vs D <sub>VEC</sub>	99.00 ± 0.60 70.25 ± 1.86 <sup>b</sup>	} Significant difference		

#### 4.0 DISCUSSION

Despite widespread international calls for a global ban on PQ use, its high weed-control effectiveness and low cost are compelling arguments for its continued use in most developing nations [21,22].

This study focused on serum hepatic enzymes like ALT, AST, ALP, and GGT for the study of hepatotoxicity and repair. When compared to the control subgroups (A<sub>0</sub> and A<sub>VEC</sub>), the enzymes activities were found to be higher in the test subgroups (B<sub>0</sub>, B<sub>VEC</sub>, C<sub>0</sub>, C<sub>VEC</sub>, D<sub>0</sub>, and D<sub>VEC</sub>), there was an interaction between the dose given and the concentration of enzyme activity; thus, the increase was found to be directly related to the dose, as seen in [23,24]. PQ studies and the toxicity effects of this chemical on the body's organs, as well as

**Comment [PM15]:** All the results of the tables must therefore be discussed and also compared with studies already published.

its mechanism of toxicity, have been reported [25]. Following a hepatocellular injury, liver enzymes are secreted into the bloodstream, resulting in an increase in their activity in serum samples. These abnormal increases in enzymes are sensitive indicators of hepatotoxicity or disease. In general, these findings may point to liver degeneration and hypofunction. Lipid peroxidation is known to disrupt the integrity of cellular membranes, resulting in cytoplasmic enzyme leakage. As a result, increased activity of these enzymes in the serum could be the result of severe hepatocellular damage. This is also consistent with several studies [24, 26, 27, 28, 29, 30] that found PQ to be a hepatotoxin.

**Comment [PM16]:** Certainly necessary to say what we see but also and above all necessary to provide an answer; Why ?

Furthermore, within-group comparisons of all enzymes revealed a well-defined reduction in the values of enzyme activities of the vitamin C and E treated subgroups (BVEC, CVEC, and DVEC) when compared to the PQ insult only subgroups (B0, C0 and D0). In this case, the combined effect of Vitamin E and C restored the above-mentioned serum parameters. Vitamin E is well known for regulating all oxidation processes in the body and thus acting as a powerful antioxidant. It works as an antioxidant by preventing chain propagation by transferring phenolic hydrogen to a peroxy free radical of a polyunsaturated fatty acid, thereby limiting the extent of lipid peroxidation. The majority of the peroxyl radical formed is reconverted to tocopherol by vitamin C (synergistic combined therapy/protective effects), while Vitamin C, to a large extent, reduced the toxic insults and acted as a substrate for the antioxidant enzyme ascorbate peroxidase, a function that is especially important in stress resistance, thereby maintaining and repairing cellular integrity and function, which resulted in the lower values.

**Comment [PM17]:** Develop please

The capacity of vitamin C to quench radicals created by PQ's redox cycling before they harm other biomolecules could potentially be linked to this protective function. This is consistent with earlier research [24, 31, 32, 33, 34, 35, 36].

Surprisingly, the findings of this study contradict those of [37, 38, 39, 40, 41], who claimed that vitamin E and C had no effect on the biochemical reaction to paraquat, implying that they had no protective effect against paraquat intoxication.

**Comment [PM18]:** If the authors' study only serves to confirm results that are already known, then there is no point. The authors must really show the innovative character of their study.

**Comment [PM19]:** Must be developp

During the PQ toxic insult, all enzyme activities in the subgroups on vitamin C and E improved in general (BVEC, CVEC, and DVEC). Despite the fact that these values were high as compared to the control subgroups (A0 and AVEC) and above the reference ranges for rat liver enzymes, they are nonetheless a sign of improved health. These findings demonstrate that if the vitamin C and E combined treatment had been continued for a longer period of

time, the liver cells may have been completely healed, and the health of the rats would have improved much better.

## 5.0 CONCLUSION

Vitamin C and E are powerful antioxidants that should be considered as a first-line treatment (emergency operation) for toxic injury. In PQ toxicity, its administration should not be overlooked, and it should be continued even after the patient has recovered to ensure enough intracellular repairs.

The findings of this study show that a combination Vitamin C and E treatment regimen can enhance the health of animals that have been exposed to PQ. PQ exposure caused a significant increase in the activities of the liver enzymes SGOT, SGPT, ALP, and GGT, which were dose-dependent. After receiving the vitamins, the activity levels were reduced throughout the study, indicating their efficacy as a supplement to the treatment.

As a result, more well-designed prospective studies are required to compare the outcomes and mortality rates of various treatment procedures. The failure of present treatments for paraquat poisoning necessitates not just extensive prospective clinical studies for paraquat poisoning treatment, but also immediate prevention actions.

## REFERENCES

1. Summers LA (1980). Paraquat and diquat. The bipyridylium herbicides, London, New York, Academic Press, p: 1-49.
2. Dinham, B. Active ingredient fact sheet, paraquat in PAN UK: Pesticide News. 1996; 32: 20: <http://www/pan-uk.org/pestnews/actives/paraquat.htm>.
3. Suntres ZE (2002). Role of antioxidants in paraquat toxicity. *Toxicology*, 180(1): 65-77. [https://doi.org/10.1016/S0300-483X\(02\)00382-7](https://doi.org/10.1016/S0300-483X(02)00382-7)
4. Selvam NT, Venkatakrisnan V, Dhamodharan R, Murugesan S, Kumar SD (2013). Hepatoprotective activity of methanolic extract of *Syzygiumjambos* (Linn.) leaf against paracetamol intoxicated Wistar albino rats. *Ayu*, 34(3): 305. <https://doi.org/10.4103/0974-8520.123133>
5. Mehmetçik G, Ozdemirler G, Koçak-Toker N, Cevikbaş U, Uysal M (2008). Effect of pretreatment with artichoke extract on carbon tetrachloride-induced liver injury and oxidative stress. *Exp. Toxicol. Pathol.*, 60(6): 475-480. <https://doi.org/10.1016/j.etp.2008.04.014>

Comment [PM20]: indicate what this study brings to the field of expertise....

Comment [PM21]: developp

Comment [PM22]: necessary to add more recent references.....

6. Halliwell B (1995). Gutteridge JM. The definition and measurement of antioxidants in biological systems. *Free Radic. Biol. Med.*, 18: 125-126. [https://doi.org/10.1016/0891-5849\(95\)91457-3](https://doi.org/10.1016/0891-5849(95)91457-3)
7. Calabrese V, Mancuso C, Sapienza M, Puleo E, Calafato S, Cornelius C, Finocchiaro M, Mangiameli A, Di Mauro M, Giuffrida Stella A, Castellino P (2007). Oxidative stress and cellular stress response in diabetic nephropathy. *Cell Stress Chaperones*, 12(4): 299. <https://doi.org/10.1379/CSC-270.1>
8. Balmus IM, Ciobica A, Trifan A, Stanciu C (2016). The implications of oxidative stress and antioxidant therapies in inflammatory bowel disease: Clinical aspects and animal models. *Saudi J. Gastroenterol. Off. J. Saudi Gastroenterol. Assoc.*, 22(1): 3. <https://doi.org/10.4103/1319-3767.173753>
9. Mohamed WA, Abd-Elhakim YM and Farouk SM (2016). Protective effects of ethanolic extract of rosemary against lead-induced hepato-renal damage in rabbits. *Exp. Toxicol. Pathol.*, 68(8): 451-461. <https://doi.org/10.1016/j.etp.2016.07.003>
10. Sarangarajan R, Meera S, Rukkumani R, Sankar P, Anuradha G (2017). Antioxidants: Friend or foe? *Asian Pac. J. Trop. Med.*, 10(12): 1111-1116. <https://doi.org/10.1016/j.apjtm.2017.10.017>
11. Carr A, Frei B. Does vitamin C act as a pro-oxidant under physiological conditions? *FASEB J.* 1999; 13:1007-1024.
12. Menzel DB. The toxicity of air pollution in experimental animals and humans: the role of oxidative stress. *ToxicolLett.* 1994; 72: 269-277.
13. Hong SY, Hwang KY, Lee EY, Eun SW, Cho SR, Han CS, Park YH, Chang SK. Effect of vitamin C on plasma total antioxidant status in patients with paraquat intoxication. *ToxicolLett.* 2002; 126: 51-59.
14. Sies H. Oxidative stress: oxidants and antioxidants. *Exp. Physiol* 1997; 82:291-5.
15. Asada K. Ascorbate peroxidase – hydrogen peroxide scavenging enzyme in plants. *Physiol Plant* 1992;85:235-41
16. Springboard (2004). Vitamin E. *The Nutrition Notebook*. SB3.com. pp:12-8. Online.
17. Rhee SG (2006). Cell signaling. H<sub>2</sub>O<sub>2</sub>, a necessary evil for cell signaling. *Sci.*, 312: 1882-3
18. Mauro P, Renze B, Wouter W. In: *Tietz text book of clinical chemistry and molecular diagnostics*. 4th edition. Carl AB, Edward R, David EB, editors. Elsevier; 2006. Enzymes; pp. 604–616.

19. AGA Technical Review on the Evaluation of Liver Chemistry Tests. *Gastroenterology*. 2002;123:1367–1384.
20. Okolonkwo, B. N., Amadi, C. F., Chikwubike, U. O and Nyenke, C. U. (2022). The comparative effects of vitamin E + C on the chronic toxicity of paraquat in albino rats (*Rattus norvegicus*). *European Journal of Medicinal Plants*, 33(6),7-13
21. Botta R, Eiamchai P, Horprathum M, Limwichean S, Chananonawathorn C, Patthanasettakul V, Maezono R, Jomphoak A, Nuntawong N (2020). 3D structured laser engraves decorated with gold nanoparticle SERS chips for paraquat herbicide detection in environments. *Sens. Actuators B: Chem.*, 304: 127327. <https://doi.org/10.1016/j.snb.2019.127327>
22. Wesseling C, van Wendel de Joode B, Ruepert C, León C, Monge P, Hermosillo H, Partanen TJ (2001). Paraquat in developing countries. *Int. J. Occup. Environ. Health*, 7(4): 275-286. <https://doi.org/10.1179/oeht.2001.7.4.275>
23. Dede EB, Okolonkwo BN, Ngokere AA. Toxicological effect of paraquat on the histology of the stomach, small Intestine and testis of male albino rats (*Rattus rattus*). *Port Harcourt Med J* 2007;2:51-5.
24. Benjamin Okolonkwo, Edna OgechiNwachukwu. The antioxidant effects of vitamin C on liver enzymes: aspartate aminotransferase, alanine aminotransferase, alkaline phosphatase and gamma-glutamyltransferase activities in rats. *Journal of Xenobiotics*.2013; 3(5):2-6. DOI: 10.4081/xeno.2013.e5
25. Bus JS, Cogen SZ, Olgaard M, Gibson JE. A mechanism of paraquat toxicity in mice and rats. *ToxicolApplPharmacol* 1976;35: 501-13
26. Clark DG, McElligott TF, Hurst EW. The toxicity of paraquat. *Br J Ind Med* 1966;23: 126-32.
27. Gibson JE, Cagen SZ. Paraquat – induced functional changes in kidney and liver. In: Wong, RC, Stevens, JB, eds. *Biochemical mechanisms of paraquat toxicity*. 2nd ed. New York: Academic Press; 1977. pp 117-136.
28. Cagen SZ, Janoff AS, Bus JS, Gibson JE. Effect of paraquat (methylviologen) on liver function in mice. *J Pharmacol Exp Ther* 1976;198:222-8.
29. Murray RE, Gibson JE. A comparative study of paraquat intoxication in rats, guinea pigs and monkeys. *Exp Mol Pathol* 1972;17
30. Akinloye, O.A., I. Adamson, O. Ademuyiwa and T.A. Arowolo, 2011. Supplementation of vitamins C, E and its combination on paraquat-intoxicated rats:

- Effects on some biochemical and markers of oxidative stress parameters. *J. Applied Pharm. Sci.*, 1: 85-91.
31. Zingg J-M (2007). Vitamin E: an overview of major research directions. *Mol. Aspects Med.*, 28(5-6): 400-422. [https:// doi.org/10.1016/j.mam.2007.05.004](https://doi.org/10.1016/j.mam.2007.05.004)
  32. Asada K. Ascorbate peroxidase – hydrogen peroxide scavenging enzyme in plants. *Physiol Plant* 1992;85:235-41.
  33. Shigeoka S, Ishikawa T, Tamoi M, Miyagawa Y, Takeda T, Yabuta Y, et al. Regulation and function of ascorbate peroxidase isoenzymes. *J Exp Bot* 2002;53: 1305-19.
  34. Wefers H, Sies H. The protection by ascorbate and glutathione against microsomal lipid peroxidation is dependent on vitamin E. *Eur J Biochem* 1988;174:353-7.
  35. Wershana KZ. The influence of vitamin C or selenium on paraquat induced toxicity in Guinea Pigs. *Pak J BiolSci* 2001;4:81-8.
  36. Tantcheva LP, Stoeva ES, Galabov AS, Braykova AA, Savov VM, Mileva MM. Effect of vitamin E and vitamin C combination on experimental influenza virus infection. *Exp ClinPharmacol* 2003;4:259-64.
  37. NastaranEizadi-Mood, Ali Mohammad Sabzghabae, Ahmad Yaraghi, Kamran Montazeri, MojganGolabi, AlirezaSharifian and ShirinsadatBadri . Effect of Antioxidants on the Outcome of Therapy in Paraquat intoxicated Patients. *Tropical Journal of Pharmaceutical Research*. 2011; 10 (1): 27-31
  38. Sullivan TM, Montgomery MR. Ascorbic acid nutritional status does not affect the biochemical response to paraquat. *Fund ApplToxicol*. 1984; 4: 754-759
  39. Eckert H, Schnorr R, Renner H, Oddoy A, Lisotschkin BG. Studies of lung damage by paraquat. 2. Effect of vitamin E on acute lung damage caused by paraquat. *Z ErkrAtmungsorgane*. 1987; 169: 147-152.
  40. Redetzki HM, Wood CD, Grafton WD. Vitamin E and paraquat poisoning. *Vet Hum Toxicol*. 1980; 22: 395-397
  41. Yasaka T, Okudaira K, Fujito H, Matsumoto J, Ohya I, Miyamoto Y. Further studies of lipid peroxidation in human paraquat poisoning. *Arch Intern Med*. 1986; 146: 681-685.