

Influence of wastes of taro leaf, sugar beet and saw dust on physiochemical parameters of produced vermicompost

Abstract:

Climatic changes and its negative impacts may ~~is represented~~ consider as one of the big global challenges, and eliminate soil and water availability time by time. ~~Thereby,~~ Adding organic fertilizers (i.e., vermicompost) as soil amelioration may consider as one of effective approaches in order to recover soil degradation and enhance water retention in soil. Through vermicomposting, agro-wastes are converted into vermicompost that rich in humus, growth promoters (i.e. amino acids, growth hormones) and nutrients.

Comment [a1]: Abstract should be more concise and not more than one paragraph.

Comment [a2]: Not appropriate here.

~~Current work-~~ This study was carried out during 2019-2020 to investigate the impact of different agro-waste types i.e., Taro leaf (TF), SawDust (SD) and Sugar Beet (SB) on vermicompost quality.

Comment [a3]: Write down in a single paragraph.

Obtained results of this study indicated that these agro-wastes resulted in varying of physiochemical parameters and vermicompost content of amino acids and growth promoters.

Comment [a4]: Not clear!! Might be parameters.

Whereas, adding SD to Cow dung during vermicomposting resulted in raising Organic matter and N content. Meanwhile, adding SD to FS resulted in increment in C:N ratio and P content in vermicompost. Besides, adding TL to cow dung during vermicomposting led to increment in amino acids.

Comment [a5]: What does it mean!

Comment [a6]: Abbreviation please!

In addition, adding SB to cow dung during vermicomposting resulted in raising ABA and GA3 content. Moreover, these different agro-wastes resulted in varying microbial activity and the highest activities produced when TL adding to FS during vermicomposting.

Comment [a7]: Need explanation!

Comment [a8]: ???

Finally, these different agro-wastes led to differing in antimicrobial activity in produced vermicompost. From the results of the study ~~se resulted~~, research team concluded that there is a great potential to produce vermicompost with specific quality that may play a crucial role in combat climatic changes particularly reinforce tolerant plant to drought stress.

Comment [a9]: Provide recommendation/suggestion using a simple sentence.

Comment [a10]: Five (5) keywords are enough!

Keywords: Vermicompost, Cow dung, Fish Sludge, Microbial activity, Antimicrobial, Nutrient contents, Growth promoters and Amino acids content

Comment [a11]: Arrange/ re-write the keywords in an alphabetical order:

Antimicrobial, Cow dung, Fish Sludge, Growth promoters and Amino acids content, Nutrient contents, Vermicompost

Introduction:

Converting agro-wastes into vermicompost is known globally as recycling organic wastes (reference!). Through this process, earthworms have a crucial role in degrading these agro-wastes through fragmentation and ingestion of organic matter (agro-wastes) and produce an efficient organic fertilizer (vermicompost) that is rich in humus and nutrients. Moreover, various microorganisms including bacteria, fungi, and actinomycetes help earthworms in its crucial role and vermicompost processing agro-wastes (reference!). Vermicomposting has been arising as an innovative eco-technology for the conversion of various types of wastes into vermicompost (Khan and Tripathi, 2020).

This vermicompost may be an effective substitute for chemical fertilizers. Whereas, Ceritoglu *et al.* (2018) reported that the vermicompost products are good sources for plant nutrient elements, various hormones, enzymes, humic substances and especially organic matter when added into the soil.

Also, Again, Rekha *et al.* (2018) emphasized that through their study, vermicompost contains a combination of macro- and micro-nutrients and the uptake of the nutrients has a positive effect on plant nutrition, growth, photosynthesis and chlorophyll content of the leaves.

Application of vermicompost may play an effective role in encourage plant growth through direct or indirect way. Vermicompost as any organic fertilizer is rich material in terms of nutrition, antioxidants, vitamins, humic and phenolic substances and various hormones (Joseph, 2019). All these substances play a significant role in promoting plant growth.

Besides, vermicompost may improve plant growth performance though its positive effect on leaf chlorophyll content, and photochemical efficiency, yield, and electron transport rate (ETR) of mature leaves, as well as increased leaf succulence, and carotenoid, protein, and amino acid content (Chenping and Mou, 2016).

In respect for indirect way, vermicompost application may improve soil conditions or works as soil amendment. Whereas several studies indicated that applying vermicompost resulted in enhancing soil properties particularly porosity, water holding capacity (WHC), cation exchange capacity (CEC) and occurrence of macronutrients (Vasanth *et al.*, 2020; Ceritoglu and Erma, 2020). Also, Additionally, vermicompost as any organic matter resulted in promoting

Comment [a12]: It is not a good idea to use same word ("ALSO") several times.

microorganisms activity in soil and that will reflect on facilitate nutrients in root zone and improving soil condition in root zone (Pathma and Sakthivel, 2012) .

In addition, Ilker and Tavalı (2014) mentioned that vermicompost appeared to more significantly increase bacterial number in soil and it has a potential to be used as an alternative to farmyard manure to improve and maintain soil biological activity. Besides, Singh *et al.* (2011) showed that applying vermicompost resulted in increasing amount of humus in soil which resulted in ~~favourable~~ favorable changes in physical, chemical and biological properties of soil, and in enhancing the water-holding capacity.

Generally, several studies reported that use of vermicompost is effective for improving soil aggregation, structure, aeration and fertility; contains most of the nutrients in plant-available form such as nitrates, phosphates, exchangeable calcium and soluble potassium; increases beneficial microbial population diversity and activity; improves soil moisture-holding capacity; contains vitamins, enzymes and hormones; and accelerates the population and activity of earthworms (Aggelides&Londra 1999, Mascolo *et al.* 1999, Albiachet *al.* 2000, Marinari *et al.* 2000, Sailajakumari&Ushakumari 2002, Arancon *et al.* 2006, Prabha *et al.* 2007, Azarmi *et al.* 2008).

Moreover, Karmakar *et al.* (2015) mentioned that the application of vermicompost showed better result in comparison to chemical fertilizers in terms of soil physical and chemical properties as well as productivity of soil.

~~Few Similar~~ studies indicated ~~the effects~~ of type agro-wastes (whether crops wastes or animal wastes) on quality of vermicompost. Ceritoglu *et al.* (2021) indicated that quality of vermicompost is ~~greatly~~ affected by ~~the~~ input of organic matter). ~~Also, A study conducted by~~ Moustafa *et al.* (2021) ~~studied reported that~~ potential of utilizing different agro-wastes (e.g., banana leaves, Rice straw and sugarcane) in feeding earthworm and effect of these different agro-wastes on quality of produced vermicompost. Their results ~~showed~~ indicated that type of agro-wastes have different impact on quality of vermicompost. ~~They Researchers~~ suggested that there is a great potential to produce vermicompost with specific quality as growers needs or requirements of plant stage via using different agro wastes.

In this context, Blouinet. *al.* (2019) showed that the best original material to be used for vermicompost production was cattle manure. Also, The maximum positive effect occurred when vermicompost represented 30 to 50% of the soil volume.

Comment [a13]: Looks like copied! Please check again.

~~Current work~~ This study investigates the impacts of different agro-wastes (i.e., taro leaf, sawdust and sugar beet) on vermicompost quality. Specially, these wastes differ among them self in ~~the~~ nutrient, amino acids contents and other composites.

MATERIALS and METHODS

This ~~work~~ study was carried out during 2019-2020 through cooperation both of National Research Center, the Central Laboratory for Aquaculture Research (CLAR) and Faculty of Science, Tanta University, [COUNTRY?]. Whereas vermicompost types were produced at the Central Laboratory for Aquaculture Research (CLAR) during (2019-2020) and all needed analysis were done at the National Research Centre (NRC), except microbiological analysis was done at the Faculty of Science, Tanta University, Country?.

Comment [a14]: Mention the "Month_Year"
Such as "from June 2019 to July 2020"

Specimen collection and processing:

Mixing three species of earthworm (*Eiseniafetida*, *lumbricusrubellus* and *Perionyxexcavatus*) were raised on cow dung and fish sludge until utilized in the experiment

Preparation of different feeding materials

Formatted: Font: Italic

- Cow dung (CD) processing:** Fresh cow dung was obtained from a cow farmer adjacent to the central lab for aquaculture research (CLAR) and applied directly to the treatments, assuming the moisture is about 50% of the wet weight.
- Fish sludge collection and preparation:** Fish sludge (FS) was collected from the concrete ponds of Nile tilapia *Oreochromis niloticus* brood stock and fry, at the Nile tilapia hatchery belonging to CLAR; during fry harvesting from the brood stock ponds as well as from fry rearing ponds. The produced FS, with a moisture content of 96.5% and dry solid content of 3.5%, was collected in barrels and then spread out in a thin layer on a cement floor for drying over fourteen days, so it can be stored safely until being used.
- Taro Leaves (TL) processing:** Taro leaves were collected from CLAR nearby farm, then sun-dried for 3 days and were crushed into small pieces.

4. **Saw dust (SD):** saw dust was bought from CLAR nearby carpentry shop. Saw dust soaked in water for three days before utilized in vermicompost producing.
5. **Sugar beet (SB):** Sugar Beetwasteswere collected from sugar beet factory and sun-dried for 3 days then minced into small pieces by machine.

Earthworm inoculation and Vermicompost production:

Both of fresh cow dung and dried fish sludge were mixed individually with agro-wastes (SD, SB or TL) at a ratio of 2:3respectively and moistened to 60-70% in Styrofoam boxes with dimensions of 60× 40× 30 cm. After 24 h, three species of earthworm (*Esienafetida*; *Perionyexcavatus* and *Lumbricusrubellus*) were added to the media at a rate of 50 g worm per 1000g media. For eight weeks the boxes were checked weekly and re-moistened and mixed until the vermicompost matured. All boxes were kept indoors and the temperature maintained between 18-25°C during the vermicompost maturation. At harvest time,vermicompost was checked manually on white plastic surface and the adult as well as pre-adult earthworms were collectedthen the vermicompost was returned to the boxes again for one more month. Later, the vermicompost was re-checked again and all hatched earthworms were collected. The harvested vermicompost was packed in plastic bags and delivered to laboratories to be analyzed.

Experimental treatments: eight setup (treatments) containing different feeding materials (Cow dung (CD) alone, Fish sludge alone (FS) and CD or FS supplemented with TL, SD and SB respectively) with three replicates of each were prepared as following

CD, FS, CD+ TL, CD+SD, CD+ SB, FS+ TL, FS+SD and FS+ SB. Samples of all these treatments delivered to different laboratories in order to be analyzed.

Analysis of chemical and microbial parameters

The following chemical and microbial parameters were measured during the study-Dry weight (g); Organic matter (%);Humidity (%); Ash; C/N ratio; Nitrogen in % (through Kjeldalh method); Available Phosphorus (through Modified Olsen's method); Available Potassium (Ammonium acetate method); Amino acids content (mg/100g dry weight); Growth promoters content (g/100g sample); Bacterial and Fungal population (through Serial dilution and plate count method) etc.

Comment [a15]: Show this explanation in a tabular form.

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Normal, No bullets or numbering

Formatted: Font: 12 pt, Bold

Formatted: Indent: Left: 0.5", No bullets or numbering

- Dry weight (g)
- Organic matter (%)
- Humidity (%)
- Ash
- C/N ratio
- Nitrogen % (Kjeldahl method)
- Available Phosphorus (Modified Olsen's method, 1958)
- Available Potassium (Ammonium acetate method)
- Amino acids content (mg/100g dry weight)
- Growth promoters content (g/100g sample)
- Bacterial and Fungal population (Serial dilution and plate count method)

Comment [a16]: Better to represent this part in a single paragraph/text.

Physiochemical analysis of vermicompost:

Vermicompost samples were dried in a ventilated oven at 70°C to constant weight for determining the dry weight and chemical analysis.

Macronutrients were extracted using the dry ashing digestion method according to Chapman and Pratt²³. Nitrogen was determined by using the Kjeldahl method, the ash was dissolved in HCl (2N) and phosphorus was photometrically determined in the digested solution using vanado-molybdate color reaction according to the method described by Jackson²⁴. Potassium was measured in the digested suspension using the Flame photometer, (Eppendorf, DR Lang). Organic matter content was determined according to Walkely and Black²⁵.

Comment [a17]: Looks like copied. Please explain more clearly.

Comment [a18]: This citation looks like copied from other sources and did not mention properly.

Analysis of free amino acids and growth promoter's analysis:

To determine the total free amino acids, the modified ninhydrine colorimetric method that described by Rosein¹⁹⁵⁷ & Selimet *et al.* (1978) was used for this purpose.

Besides, growth promoters in samples of vermicompost were estimated according to the method described by Dobrevet *et al.* (2005)

Microbiological analysis of vermicompost samples:

Sampling and sample preparation:

Five grams from vermicompost samples were placed in sterile Stomacher bags and treated by a Stomacher 400 Circulator for 60 s at middle speed after adding 45 ml sterile 0.85% NaCl. The Stomacher blending step was repeated three times and the microbial suspension was obtained.

Estimation the counts of total viable bacteria count:

Tenfold serial dilution of the microbial suspensions obtained with the protocol described above made with sterile 0.85% NaCl were plated onto plate count agar medium for the estimation of total viable counts, counts of colony forming units (CFU) were estimated after 3 days of incubation at 28°C and were calculated per gram vermicompost. The total resistant bacteria were estimated by planting the same dilution onto plate count agar medium sublimated with (20mg/l) for (penicillin, ampicillin, erythromycin and tetracycline respectively).

Antimicrobial activity assay:

This method was done by agar well diffusion test according to (Schillinger and Luck, 1989). To determine the antimicrobial activity of the vermicompost samples against the selected identified pathogenic bacteria e.g. *Citrobacterfreundii*, *Enterobacter cloacae*, *Pseudomonas aeruginosa*, and *Klebsiella pneumoniae*. The prepared nutrient agar plates were overlaid with 100 µl of overnight culture of tested pathogens (in nutrient broth), then spread well with L-shaped glass rod. After 15 min, wells of 5 mm diameter were made with a sterile cork borer. Samples of vermicomposts extract were placed into wells. Plates were then incubated at 30°C for 12 hrs. The inhibition zones were measured to assay the antimicrobial activity of vermicompost samples.

Data analysis: Vermicompost samples were analyzed using the standard procedures in the laboratory at National research Centre and Faculty of Science, Zoology department, Tanta University, [Country???](#). All data are the means of triplicates. Statistical analysis of data, analysis of variance (ANOVA) and mean separation were carried out using Duncan's multiple range test and significance were determined at the ($p \leq 0.01$) level (Duncan, 1955). Data analysis was performed using ASSISTAT version 7.7 beta (2015).

Formatted: Font: Not Bold, Italic

Results and Discussion:

Table 1 showed that utilizing Saw Dust (SD) in producing vermicompost was resulting in higher dry weight and organic matter comparing with other types of vermicompost types. In respect for ash percentage, ~~data findings was were~~ indicated that the highest ash percentage achieved with (FS) alone or (FS+TL). However low ash (%) was achieved when (CD) used alone to produce vermicompost. In respect to nutrients content utilizing SD in vermicompost production resulted in increasing N % content with CD and P% content with FS. However, higher K % content was obtained when (FS+SB) utilized in vermicompost producing.

In addition, table (2) represented ~~data~~ concern amino acids and growth promoters influenced with different carbon sources. Higher amino acids content was obtained when using TL wastes with CD in producing vermicompost. Meanwhile utilizing SB wastes with CD in vermicompost lead to increasing ABA (1.62g/100g) and GA₃ (2.23) comparing with two other carbon sources either (SD) or (TL) wastes. Besides, using wastes of SB with FS was resulted in producing the highest content of IAA (0.19g/100g).

~~There was many variation were observed in physio-chemical results,~~ physio-chemical results showed variation. The vermicompost that produced from Cow Dung organic matter represented high results in comparison with Fish Sludge organic matter in all treatments (Table 1). The organic matter results was 46.6, 50.15, 33.00 and 37.33 with Cow Dung vermicompost (Cow dung, Saw Dust, Sugar beet and Taro Leaves respectively), while, the results with Fish Sludge was 21.06, 32.14, 26.48 and 19.73 (Fish sludge, Saw Dust, Sugar beet and Taro Leaves respectively).

Similar to the aforementioned results in organic matter, Cow Dung vermicompost types represented the highest results comparing with Fish Sludge types in the case of total nitrogen.

The opposite result occurred on C/N, i.e., the most of Fish Sludge results increase comparison to Cow Dung vermicompost except with Cow Dung alone. Huiet *al.* (2007) reported that compost made from the mixture spent mushroom waste with livestock manure, and soybean cake had a C/N ratio of 30. The compost was produced through composting alone (Huiet *al.*, 2007). Based on the results of statistical analysis, composition of vermicompost materials affected the total

organic C, total N and C/N ratio of the vermicompost. It showed that vermicomposting can lead to mineralization process which reduced C by N ratio of vermicompost. Our results were in line with previous report according to the research of Huiet *al.* (2007).

Hoitink and Boehm, (1999) reported that C/N ratio is one of the parameters used for measuring compost maturity. Researchers have suggested various ideal C/N ratios ranging from >12 to <25 (Brewer and Sullivan, 2003). According to Gomez-Brandonet *al.* (2008) the stabilized compost has a C/N ratio from 10 to 15. Majlessiet *al.* (2012) showed that to produce a stable vermicompost of food waste needed 7 weeks of the process duration. The vermicomposting duration in our research was four weeks and then continued by composting for two weeks. Domiguez, (2011) reported that the decaying organic in vermicomposting system is a spatially and temporally heterogeneous matrix of organic resources with contrasting qualities that result from the different rates of degradation, occur during decomposition. This means that the vermicompost still needs the composting process. To improve the quality of generated vermicompost, the systematic study of composting process and the addition of additives material such as fish meal and egg shell flour was conducted. After the composting process, the quality of vermicompost increased which marked by a lower C/N ratio of vermicompost (average 15.3). The decrease in C/N ratio was caused by the decomposition process of microorganisms during the composting process. During decomposition process, soil microorganisms burn carbon as a source of energy, but not all of the carbon remains in its body; a certain amount is lost as carbon dioxide during respiration (Frankenberger and Abdelmagid, 1985). The decrease in C/N ratio of compost improve the compost quality. Frankenberger and Abdelmagid, (1985) states that the organic matter with a value of C/N ratio lower than 20 include high quality of organic matter and will undergo mineralization in the soil. Majlessiet *al.* (2012) stated that the vermicompost with low C/N ratio (14-30) indicate a mature and stable vermicompost.

In addition, Igbokwe *et al.* (2015) ~~showed-reported~~ that the mixing sawdust with the other organic waste in composting process ~~can~~ could increase N content of the organic fertilizer (Mtambanengwe and Kirchmann, 1995) and increase in N content lower the C/N ratio.

Formatted: Font: Not Bold

Table 1: effect of different agro-wastes (TL, Sb and SD) as carbon sources on physiochemical parameters of vermicompost

TRT	FW (g)	Dry Weight (g)		O.M (%)		Humidity (%)		Ash		N (%)		C / N ratio		P (%)		K (%)	
[#] CD (Control)	10	6.70	b	46.60	a [*]	33.00	b	20.40	d	1.33	b	20.40	a	0.35	ba	1.08	c
CD + SD	10	7.68	a	50.15	a	23.20	d	26.65	cd	2.51	a	11.48	b	0.12	b	1.65	b
CD + SB	10	6.80	b	33.00	ab	32.00	b	30.00	bcd	1.87	ab	11.58	b	0.18	ba	1.70	b
CD + TL	10	6.01	c	37.33	ab	39.90	a	22.77	d	1.87	ab	11.23	c	0.45	ba	1.58	b
FS (Control)	10	7.27	a	21.06	b	27.33	c	54.94	a	1.09	b	11.58	b	0.40	ba	1.88	a
FS + SD	10	7.62	a	32.14	ab	23.83	cd	44.03	abc	1.61	b	11.63	b	0.48	a	1.53	b
FS + SB	10	7.45	a	26.48	b	25.50	cd	46.02	ab	1.54	b	11.50	b	0.42	ba	2.13	a
FS + TL	10	7.52	a	19.73	b	24.83	cd	52.10	a	0.99	b	11.58	b	0.36	ba	1.42	b

Formatted: Superscript

Formatted: Font: Not Bold, Superscript

Means were represented as average of replicates.

^{*}Different letters are express for significant differences while the same letters are non-significant at L.S.D. $p > 0.05$ (DMRT).

[#]CD= cow dung, SD= Saw dust, SB= Sugar beet, TL= Tour leaf FS= Fish

Formatted: Font: 11 pt, Superscript

Formatted: Font: 11 pt, Superscript

Table 2: Effect of different agro-wastes (TL, SB and SD) as carbon sources on amino acids and growth promoters in produced vermicompost

	Amino acids (mg/g DW)	ABA (g/100 g)	GA3 (g/100 g)	IAA (g/100 g)
CD (Control)	0.27 c	0.33 c	1.08 c	0.04 c
CD + SD	0.41 c	0.70 b	0.37 f	0.05 c
CD + SB	1.43 b	1.62 a	2.23 a	0.15 b
CD + TL	1.82 a	0.19 d	0.94 d	0.05 c
FS (Control)	0.44 c	0.01 e	0.16 g	0.03 c
FS + SD	0.29 c	0.01 e	0.92 d	0.02 c
FS + SB	0.46 c	0.17 d	1.59 b	0.19 a
FS + TL	0.34 c	0.67 b	0.78 e	0.13 B

Means were represented as average of replicates.

Different letters are express for significant differences while the same letters are non-significant at L.S.D. $p > 0.05$ (DMRT).

CD= cow dung, SD= Saw dust, SB= Sugar beet, TL= Tour leaf FS= Fish

Effect of carbon sources Microbial activity:

Formatted: Font: Font color: Auto

Table 3: ~~show~~ Microbial activity in different vermicompost types based different agro-wastes (TL, SB and SD) as carbon source

TRT	CFU
FS + SD	11.33± 2.83
FS + SB	44.00 ± 4.71
FS + TL	56.00 ± 0.00
CD + SD	32.33 ± 4.24
CD + SB	29.17 ± 3.06
CD + TL	28.67 ± 11.79

Values are represented as average of three replicates.

Different letters within the same column express significant differences at L.S.D. $p > 0.05$.

FS= Fish sludge, CD= cow dung, SD = Saw dust SB= sugar beet, and TL= Taro leaves




Fig1: Microbial activity influenced by utilizing different agro-wastes as carbon source;

Formatted: Font: 11 pt, Font color: Auto

Formatted: Font: 11 pt, Not Bold, Font color: Auto

Formatted: Font color: Auto

Table 3 and Fig 1 represent data concern impact of different carbon sources (SD, SB and TL) on microbial activity in produced vermicompost. From first view it was observed that FS vermicompost with whether TL or SB wastes surpassed in microbial activity comparing with CD vermicompost types with the same type of carbon source (TL or SB). [Also Again](#), incorporation TL wastes with FS during vermicompost processing resulted in considerable increment in microbial activity (56 CFU) comparing with other studied treatments meanwhile adding TL to CD during vermicomposting produced lower value of microbial activity (28.67 ± 11.79). The lowest value of microbial activity (11.33 ± 2.83) was achieved with (FS+ SD).

[Also Moreover](#), *Loreset al.* (2006) stated that the vermicomposting process involved the activity of earthworms which modify wastes physically and feces excreted by worms can increase the activity of the microorganisms so that the rate of mineralization to be faster.

[Data Findings of this study represented](#) in table 4 showed that anti-pathogenic activity of vermicompost was varied with varying added agro-wastes (as carbon source i.e. SD, SB and TL) whereas CD+SD resulted in the highest anti-activity for *Bacillus anthracis* (15mm) comparing with other vermicompost types. Meanwhile, this vermicompost has no anti-pathogenic effect for

both of *Escherichia coli* and *Pseudomonas aeruginosa*. Besides, it can be noticed that FS+SD surpassed other types of vermicompost whereas it has higher anti-pathogenic effect for three pathogens *Escherichia coli*, *Klebsiella pneumoniae* and *Pseudomonas aeruginosa* in addition to *Bacillus anthracis*.

Moreover, vermicompost from (FS + TL) showed anti-pathogenic effect for only three of studied pathogenic (*Bacillus anthracis*, *Escherichia coli*, and *Pseudomonas aeruginosa*).

Obtained results-Major findings of this research indicated that anti-pathogenic effect differed in vermicompost according to input crop wastes. Also, some crop wastes may-might be resulted in multi-anti-pathogenic effect than other.

Table 4: Antimicrobial activity influenced by utilizing different agro-wastes (TL, SB and SD) as carbon source

TRT	<i>Bacillus anthracis</i>	<i>Escherichia coli</i>	<i>Klebsiella Pneumonia</i>	<i>Pseudomonas aeruginosa</i>
[#] CD+S D	15mm*	00	7mm	00
CD + SB	9mm	00	16mm	00
CD + T L	9mm	00	00	25mm
FS + SD	11mm	17mm	23mm	26mm
FS + SB	8mm	00	22mm	13mm
FS + T L	7mm	7mm	00	20mm

*Values are represented as average of three replicates.

[#]FS= Fish sludge, CD= cow dung, SD = Saw dust SB= sugar beet, and TL= Taro leaves

Numerous studies (A study carried out by Garczyńska et al. (2020) highlighted the effectiveness of vermicomposting manure of different origins to reduce pathogenic microorganisms. A reduction in pathogens occurs by passage through the intestines of an earthworm, which obviates the need to raise the temperature. Moreover, other studies indicated that earthworm influences microbial community, physical and chemical properties of soil. They breakdown large soil particles and leaf litter and thereby increase the availability of organic matter for microbial degradation and transforms organic wastes into valuable vermicompost by grinding and digesting them with the help of aerobic and anaerobic microbes (Maboeta and Van Rensburg, (2003). Earthworms' activity is found to enhance the beneficial micro flora and

Formatted: Font: 11 pt, Not Bold, Superscript

Formatted: Font: 11 pt, Not Bold

Formatted: Font: Not Bold

Formatted: Font: 11 pt, Not Bold

Formatted: Font: Not Bold

Formatted: Font: 11 pt, Not Bold

Formatted: Font: Not Bold

Formatted: Font: 11 pt, Not Bold

Formatted: Font: Not Bold

Formatted: Font: 11 pt, Not Bold

Formatted: Font: Not Bold

Formatted: Font: 11 pt, Not Bold

Formatted: Font: Not Bold

Formatted: Font: 11 pt, Superscript

Comment [a19]: Not appropriate here.

suppress harmful pathogenic microbes. Soil worm casts are rich source of micro and macronutrients, and microbial enzymes Lavelle and Martin (1992).

Other studies indicated that earthworm lives in medium rich in wide range of microbes thereby earthworm strive to cover its bodies with mucus rich in antimicrobial to protect their bodies from pathogenic. These excreted mucus lost in the medium via earthworm movement from a while to while and earthworm excretes new mucus continuously. Therebyvermicast is mixed with mucus secretion of the earthworm's gut wall (Moustafaet al., 2021). Whether the antimicrobial produced from gut wall of earthworms or from excreted mucus to protect earthworms bodies against pathogenic microbes, its affected with type of meals (type of agro-wastes) that offer to earthworms as shown in the results of current study.

Generally Gradually, these results were supported with the findings of Khwanchai, and Kanokkorn,(2018) who reported that provided new evidence that agricultural waste, especially soybean meal could be used as feeds for the high quality of vermicompost production and earthworm biomass.

Bajalet al.(2019) showed that nutrient content of vermicompost was varied significantly among the substrates(Lantana camara, Ageratum conyzoides, banana pseudo stem, garden waste, vegetable waste and cow dung). Also, They reported that Lantanawas found most effective with 2.53% N, 1.38% P and 2.28% P.

Ramnarainet al., (2019) mentioned that three agro-wastes were used in this study (study namely T1 (Rice straw), T2 (Rice straw + grass) and T3 (Grass)), and Such results showed- indicated that, the combination of rice straw and grass had the highest rate of vermicompost production of 105 kg/m² followed by grass and rice straw with 102.5 kg/m² and 87 kg/m², respectively, at the end of 120 days. Besides, the harvested vermicompost had an excellent nutrient status, confirmed by the chemical analyses, and contained all the essential macro- and micronutrients.

Conclusion

What about further research! Please explain the key findings of the research within few sentences and provide suitable recommendations based on the findings!

Formatted: Font: Not Bold, Italic

Formatted: Font: 12 pt, Not Bold, Superscript

Formatted: Font: 12 pt, Not Bold, Superscript

Formatted: Font: 12 pt, Not Bold, Superscript

Formatted: Font:

Formatted: Comment Text, Left, Line spacing: single

Author(s) can provide few suitable/ effective concluding remarks within 1-3 sentences at the end.

Formatted: Font: (Default) Times New Roman, 11 pt, Font color: Auto, Pattern: Clear

Formatted: Comment Text, Left, Line spacing: single

UNDER PEER REVIEW

References

- Khan M.A. and K. N. Tripathi.(2020).** Vermicomposting with Cow Dung, Banana Plant and Vegetable Waste. International Research Journal of Engineering and Technology (IRJET) vol.7(6): 3287- 3290.
- Ceritoğlu M., Sezerşahin and Murat Erman. (2018).** Effects of Vermicompost on Plant Growth and Soil Structure. Selcuk Journal of Agriculture and Food Sciences, ,vol 32, (2). <https://doi.org/10.15316/SJAFS.2018.14333>
- Rekha, G.S., Kaleena, P.K., Elumalai, D. M. P. Srikumaran and V. N.Maheswari. (2018).** Effects of vermicompost and plant growth enhancers on the exo-morphological features of Capsicum annum (Linn.) Hepper. Int J Recycl Org Waste Agricult 7, 83–88 (2018).<https://doi.org/10.1007/s40093-017-0191-5>
- Joseph PV, (2019)**Efficacy of Different Substrates on Vermicompost Production. In: A Biochemical Analysis, Organic Fertilizers - History, Production and Applications (Ed. by M, Larramendy, S, Soloneski) IntechOpen: <https://doi.org/10.5772/intechopen.86187>
- Chenping X. and B. Mou. (2016).** Vermicompost affects soil properties and spinach growth, physiology, and nutritional value. Hort. Sciencevol.51(7) <https://doi.org/10.21273/HORTSCI.51.7.847>
- Vasanth.P.C.; K, Balaji; Saravanan, S.; Gunasekaran, S.; Srinivasan, G.R.; P R, K.S.; G, M. (2020).** Effect of Vermicompost Application on Soil and Growth of the Plant SesamumindicumL. <https://doi.org/10.20944/preprints202002.0080.v1>
- Ceritoglu M. and M. Erman. (2020).** Effect of Vermicompost Application at Different Sowing Dates on some Phenological, Agronomic and Yield Traits. J. Int. Environmental Application & Science, Vol. 15(3): 158-166.
- Pathma, J., Sakthivel, N. (2012).** Microbial diversity of vermicompost bacteria that exhibit useful agricultural traits and waste management potential. SpringerPlus 1, 26 (2012). <https://doi.org/10.1186/2193-1801-1-26>
- Ilker U. and I. E. Tavali.(2014).**Short-Term Effect of Vermicompost Application on Biological Properties of an Alkaline Soil with High Lime Content from Mediterranean Region of Turkey.The scientific Word journal vol. 2014, Article ID 395282, 11 pages.<https://doi.org/10.1155/2014/395282>
- Singh B.K., K.A. Pathak, A.K. Verma, V.K. Vermaand B.C. Deka. (2011).** Effects of vermicompost, fertilizer and mulch on plant growth, nodulation and pod yield of French bean (Phaseolus vulgaris L.). Vegetable Crops Research Bulletin vol. 74, 153-165. <https://doi.org/10.2478/v10032-011-0013-7>

Comment [a20]:

1. Check each and every reference according to the Journal's guidelines again.
2. Rearrange the reference in an alphabetical order.
3. Check both text and reference section simultaneously.
4. Should follow the Journal style/ format.
5. Cite all the references properly.

- Aggelides S.M., Londra P.A. (1999).** Effect of compost produced from town waste and sewage sludge on the physical properties of a loamy and a clay soil. *Bioresource Technology* 71: 253-259.
- Mascolo A., Bovalo F., Gionfriddo F., Nardi S. (1999).** Earthworm humic matter produces auxin like effects on *Daucus carota* cell growth and nitrogen metabolism. *Soil Biology and Biochemistry* 31: 1303-1313.
- Albiach R., Canet R., Pomares F., Ingelmo F. (2000).** Microbial biomass content and enzymatic activities after application of organic amendments to a horticultural soil. *Bioresource Technology* 75: 43-48.
- Marinari S., Masciandaro G., Ceccanti B., Grero S. (2000).** Influence of organic and mineral fertilizers on soil biological and physical properties. *Bioresource Technology* 72: 9-17.
- Sailajakumari M.S., Ushakumari K. (2002).** Effect of vermicompost enriched with rock phosphate on the yield and uptake of nutrients in cowpea (*Vigna unguiculata* L. Walp.). *Journal of Tropical Agriculture* 40: 27- 30.
- Arancon N.Q, Edwards C.A., Bierman P. (2006).** Influences of vermicomposts on field strawberries: Part 2. Effects on soil microbial and chemical properties. *Bioresource Technology* 97: 831-840.
- Prabha K.P, Loretta Y.L., Usha R.K. (2007).** An experimental study of vermin-biowaste composting for agricultural soil improvement. *Bioresource Technology* 99: 1672-1681.
- Azarmi R., Giglou M.T. Talesmikail R.D. (2008).** Influence of vermicompost on soil chemical and physical properties in tomato field. *African Journal of Biotechnology* 7 (14): 2397-2401.
- Karmakar S., M. M. Adhikary, A Gangopadhyay and K. Brahmachari. (2015).** Impact of Vermicomposting in Agricultural Waste Management vis-à-vis Soil Health Care. *Journal of Environmental Science and Natural Resources* 8(1):99. <https://doi.org/10.3329/jesnr.v8i1.24680>
- Moustafa, Y.T. A.; E.S, El-Hady; M. F. El-Dahshour; R., Zuhair; L. Zhang and N. S. A. Mustafa. (2021).** Impact of different agro-wastes on quality of vermicompost made of either fish sludge or cow dung. *Asian Journal of Plant science*, 20 (3): 370-379. <https://doi.org/10.3923/ajps.2021.370.379>
- Blouin, M., Barrere, J., Meyer, N. et al. (2019).** Vermicompost significantly affects plant growth. A meta-analysis. *Agron. Sustain. Dev.* 39, 34. <https://doi.org/10.1007/s13593-019-0579-x>

Olsen, S .R.; C. W.Cole; F. S. Watanabe and L. A. Dean (1958). Estimation of available phosphorus in soil by extraction with sodium bicarbonate. U. S. Dept. Agric. Circular No. 930, 1 – 19.

Chapman H.D. and Pratt P.F. (1978). Methods of analysis for soils, plants and waters. Division of Agric. Sci., Univ. California, Berkeley, USA 309 p.

Formatted: Indent: Left: 0", First line: 0"

Garczyńska M. , J. Kostecka, G. Pączka, E. Hajduk, A. Mazur-Pączka and K. R. Butt. (2020). Properties of Vermicomposts Derived from Cameroon Sheep Dung. Appl. Sci. 2020, 10(15), 5048; <https://doi.org/10.3390/app10155048>

Maboeta, M.S., and L. Van Rensburg, (2003). Vermicomposting of industrially produced wood chips and sewage sludge utilizing Eisenia foetida. Ecotoxicol. Environ. Saf., 56:265–270.

Lavelle, P., and A. Martin, (1992). Small scale and large scale effects of endogeic earthworms in soil organic matter dynamics in soils of the humid tropics. Soil Biol. Biochem., 24:1491–1498.

Walkely, A. and Black, I.A. (1934). An examination of the degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. Soil Sci., 37: 29-38.

Rosein, H. (1957). A modified ninhydrine colorimetric analysis for amino acids. Acch. Biochem. Biophys. 67:10-15.

Selim, H.H.A., Fayek, M.A. and Veidan, A.M. (1978). Reproduction of Bircheraffle cultivar by layering. Annals of Agric. Sci., Moshtohor, Egypt. 9:157-166.

Dobrev, P.I., L. Havlicek, M. Vagner, J. Malbeck and M. Kaminek, (2005). Purification and determination of plant hormones auxin and abscisic acid using solid phase extraction and two-dimensional high performance liquid chromatography. J. Chromatogr. A, 1075: 159-166.

Schillinger, U. and F.K. Lucke, (1989). Antibacterial activity of Lactobacillus sake isolated from meat. Applied Environ. Microbiol., 55: 1901-1906. 30.

Duncan, D.B., (1955). Multiple range and multiple F tests. Biometrics, 11: 1-42.

Hui, Z., Jianhua, L., Jianqing, D., Meiyuan, C. and Yi, C. (2007). The alternative uses of spent mushroom compost, Fujian mushroom R&D Station, 350005, Fuzhou, P.R. China. www.tuinbouw.nl/sites/default/files/documenten/00026610. Accessed on September 21st, 2016.

- Hoitink, H. A. J. and Boehm, M. J. (1999).** Annual Rev Phytophatol, 37, 427-446.
- Brewer, L.I. and Sullivan, D.M. (2003).** Compost Sci. Util., 11, 96-112.
- Gomez-Brandon, M., Lazcano, C. and Domingueza, J. (2008).**Chemosphere, 70, 436-444.
- Majlessi, M., Eslami, A., Saleh, H.N.,Mirshafieean, S.andBabaii, S. (2012).** Iranian J Environ Health Sci Eng., 9, 1-6.
- Dominguez, J. (2011).** The Microbiology of Vermicomposting, Taylor & Francis Group, LLC.
- Frankenberger, W .T. and Abdelmagid, H. M. (1985).** Plant Soil, 87, 257-271.
- Igbokwe, P. K., Asadu, C. O. Okpe, E. C.andOkoro, S. E. (2015).** Int. J. Novel Res. Phys. Chem. Mathematics, 2 (3), 6-15.
- Mtambanengwe, F. and Kirchmann, H. (1995).** Soil Biol. Biochem, , 27, 1639-1651.
- Lores, M., Gomez-Brandon, M., Perez-Diaz, D.and Dominguez, J. (2006).** Soil Biol. Biochem, 38, 2993-2996.
- Khwanchai, K. and S. Kanokkorn.(2018).** Effect of Agricultural Waste on Vermicompost Production and Earthworm Biomass.Journal of Environmental Science and Technology 11(1):23-27. <https://doi.org/10.3923/jest.2018.23.27>.
- Bajal, S., S., Subedi and S. Bral. (2019).** Utilization of Agricultural wastes as substrates for vermicomposting. Journal of Agriculture and Veterinary vol. 12(8)pp 79-84. <https://doi.org/10.9790/2380-1208017984>
- Ramnarain, Y.I., Ansari, A.A. andOri, L.(2019).** Vermicomposting of different organic materials using the epigeic earthworm Eiseniafoetida. Int J Recycl Org Waste Agricult 8, 23–36 (2019).<https://doi.org/10.1007/s40093-018-0225-7>