

## Original Research Article

### ~~Assessment of Garlic Production Constraints and Trait Preferences in Garlic Cultivar Development in Two Woredas, Sidama Region, Ethiopia~~

Formatted: Strikethrough

#### Main traits of garlic production in Two Woredas of Ethiopia: An empirical study

Comment [A1]: I suggest modifying the title

#### Abstract:

Garlic (*Allium sativum* L.) is the major food and cash crop in the highland regions of Ethiopia. However, farmers are not integrated into the garlic breeding process. The objectives of this assessment were to identify farmers' key garlic production constraints and establish preferred traits in garlic cultivar development in the two woredas of Sidama Region (Ethiopia). A participatory rural appraisal (PRA) study was conducted through a structured survey involving 36 households in Malga and Gorche woredas. A structured survey used a questionnaire which was administered to farmers to collect information on key production constraints, varieties of garlic seed used for production, seed source etc., according to the assessment result of this study, the cultivar used for garlic production is the local variety in the two woredas. Hence, the most important garlic production constraints are lack of high yielding cultivars, garlic disease and insects. To conclude active farmer participation in early breeding stages is critical for a successful garlic breeding program. Based on the assessment it is possible to recommend that high yielding garlic variety, disease tolerance, insect resistant and cultivars having high dry matter content are the most important parameters preferred by farmers, which helps them to produce maximum products and serve as a source of income to many rural farmers.

**Keywords:** assessment, constraints, farmers, food supplement, garlic, source of income

#### 1. INTRODUCTION

Garlic (*Allium sativum* L.) belongs to the Alliaceae family and is the second most widely used Allium next to onion [19]. In Ethiopia, garlic is one of the important bulb crops produced for home consumption as spice or condiment in the preparation of soup, pickle and other

preservatives as well as a source of income to many rural farmers in many parts of the country [4]. Despite its importance, great potential for production and high market demand, the current garlic production and productivity is limited and remain seasonal. Low soil fertility is one of the factors limiting the productivity of different crops in Ethiopia mainly accounted for removal of surface soil by erosion, nutrients removal by crops from the soil, complete removal of plant residue from farm-land and lack of crop rotation system on the farm land resulting in lower crop yields [1, 6].

To mitigate these problems, Ethiopian farmers have a tradition to maintain or improve the fertility of the soil using fallowing, farmyard manure, intercropping and crop rotation systems [5]. Application of manure enhances soil fertility and increases the productivity of garlic by improving soil properties and nutrient status. Similarly, [16] reported that application of organic materials like farm yard manure, compost or green manure in combination with inorganic fertilizer improved soil physical properties and the author indicated that the uptake of nitrogen, phosphorus, potassium, zinc, manganese, copper and iron were increased significantly by crops when 50% of organic fertilizers in combination with 50% inorganic fertilizers were applied.

Several scientists have emphasized the need for active farmer participation in plant breeding is critical for successful adoption of improved varieties and their production packages [18, 11]. However, the link between research and farmers is still very weak or absent in most developing countries [13]. According to Sidama Region Agricultural Office, there is no improved garlic variety to be used by the farmers currently and it is the local garlic cultivar that the farmers use for production.

Therefore, the assessment of farmers' knowledge and preferences in cultivar development was undertaken through farmer participatory approaches. This assessment was primarily aimed at establishing farmers' knowledge on garlic production constraints and their preferred traits for future selection to guide farmer-oriented garlic breeding in the Region. Therefore, the objectives of this survey were to identify and analyze farmer's key constraints in garlic production, and establish farmers' preferred traits to be included in cultivar development and variety selection process at the study sites of Sidama Region, Ethiopia.

## 2. Material and Methods

### 2.1. Description of the study areas

The assessment of this study focused on highland areas of Sidama Region, Ethiopia and covered two garlic growing woredas namely Malga and Gorche. These areas are the most fertile and productive, and their climatic conditions are well suited for garlic production in the Region.

Malga is located at 06° 54' 31" N latitude and 38° 42' 30" E longitudes covering a total area of 206.74 square kilometers with 129,694 inhabitants [7]. Most Kebeles (smallest Ethiopian administrative unit in Ethiopia) are classified as 'rural' [7]. The rainfall distribution is bimodal with a long (June to September) and short (March to May) rainy season. The altitude ranges in elevation from 1500 to 3000 m.a.s.l and receives 1200–1600 mm rainfall annually; the average annual temperature ranges from 12.6 to 20°C. The dominant soil type in the woreda is Alisol, characterized by loam to clay-loam textural classes [10].

Gorche is one of the woredas in the Sidama Region of Ethiopia. And it is located in the Great Rift Valley, Gorche is bordered on the southwest by Wensho, on the west by Shebedino, on the north by Malga, on the east by the Oromia Region, and on the southeast by Arbegona.

### 2.2. Data collection

A questionnaire was developed and administered to farmers to collect information on farm size, land allocated to garlic, source of garlic seeds, constraints for garlic production and other parameters. Different administrative levels were considered which includes woreda and kebele leaders. A random sampling involved two woredas, three kebeles per woreda and six respondents per kebele. This resulted in a total of two woredas, 6 kebeles, and 36 respondents. Data were gathered using a structured survey questionnaire to get characteristics of the farms and production systems and different factors that reduces garlic production in the two woredas.

**Comment [A2]:** what are those other parameters, they should mention them

**Formatted:** Highlight

**Formatted:** Highlight

### 2.3. Sampling procedure

A purposive sampling procedure was used to identify the highland regions chosen for their great importance in garlic production [12]. Random sampling was employed to select farmers in each

kebele from the two woredas with the help of the kebele leaders and extension workers. During data collection, participatory rural appraisal allowed farmers to express their opinions through group discussions. A checklist was prepared in advance to guide the discussion.

The permission letter was brought from Sidama Region Agricultural Office; the Office also gave us a letter to give it to the study woredas and to the respective study kebeles.

### 2.4. Data analysis

Data were analyzed using Excel. Descriptive statistical measures like,percentage, frequency, and mean were used to summarize and categorize the assessment data.Tables and bar graphs were used to represent the data.

### 3. Results

According to the assessment results of this study from the two woredas, all household heads had greater than fifteen years of experience on garlic production (Table 1).

**Table 1.** Experience of farmer’s garlic production in the study areas

Experiences (years)	Malga		Gorche		Total	
	Frequency(n=18)	%	Frequency(n=18)	%	Frequency(n=36)	%
<5	0	0	0	0	0	0
5-10	0	0	0	0	0	0
10-15	0	0	0	0	0	0
> 15	18	100	18	100	36	100

N= number of sample household heads

#### 3.1. Gender composition and decision-making in garlic production

**Formatted Table**

**Comment [A3]:** It is not relevant to place a table showing only one result

**Formatted:** Highlight

**Formatted:** Font: Not Bold, Font color: Auto, Highlight

**Formatted:** Highlight

**Formatted:** Font: Not Bold, Font color: Auto, Highlight

**Formatted:** Highlight

**Formatted:** Font: Not Bold, Font color: Auto, Highlight

**Formatted:** Highlight

**Formatted:** Font: Not Bold, Font color: Auto, Highlight

**Formatted:** Highlight

**Formatted:** Font: Not Bold, Font color: Auto, Highlight

**Formatted:** Highlight

Results on gender involvement in decision-making on garlic production indicated that both males and females were involved in the main garlic production activities. But the number of male participants were greater than female participants in the two woredas (Table 2).


**Table 2.** The number of farmers interviewed and gender composition in the study areas

Woreda	Male	(%)	Female	(%)	Total	(%)
Malga	14	77.78	4	22.22	18	100
Gorche	16	88.89	2	11.11	18	100

Formatted Table

### 3.2. Household heads educational status in the study areas


The below figure showed that the educational level of the household head of the interviewed individuals were (50%) illiterate, (27.8%) 1-6 grade and (22.2%) 8-10 grade in Malga woreda and (66.6%) illiterate, (16.7%) 1-6 grade and (16.7%) 8-10 grade in Gorche and the percentage of literate in the two woreda was (0%) (Figure 1).



**Figure 1.** Household heads educational status in the study areas

### 3.3. Source of seed

Farmers in the study areas acquired garlic seed from their own field (33.3%) and from the open market (66.7%) in Malga woreda. And in the case of Gorche woreda(38.9%) was from their own field and(61.1%) from the open market. Research institutions and private companies play no role (0 %) as seed providers from the two woredas (Figure 2).



**Figure 2.** Source of garlic seed among 36 farmers in the study areas

### 3.4. Total and cultivated land size for garlic production

The land to be allocated for garlic production in the study areas was presented in (Table 3). Most farmers allocated farm size of (5.95%) hectare for the production of garlic in Malga and (6.58%) in Gorche woredas respectively (Table 3).

**Table 3.** Mean household farm size and cultivated land for garlic in the study areas

Woreda	mean total land holding (ha)	mean garlic production area (ha)	(%)
Malga(n=18)	0.84	0.05	5.95
Gorche(n=18)	0.76	0.05	6.58

Formatted Table

### 3.5. Major garlic production constraints

The most important identified garlic production constraints across the study areas were as follows: insects (38.9%), low yield (27.8%), lack of fertilizer (16.7%) and garlic disease (16.6%) in Malga woreda. And, lack of fertilizer (33.3%), insects (27.8%), garlic disease (22.2%) and low yield (16.7%) were constraints in Gorcheworeda (Table 4).

**Table 4.** Garlic production constraints in the study areas

Constraints	Malga		Gorche	
	Number of farmers	(%)	Number offarmers	(%)
Garlic diseases	3	16.6	4	22.2
Insects	7	38.9	5	27.8
Lack of fertilizer	3	16.7	6	33.3
Soil degradation	0	0	0	0
Low yield	5	27.8	3	16.7

Formatted Table

### 3.6. Garlic varieties grown in the study areas

Garlic varieties grown in the study areas were presented in (Table 5). Even though different improved garlic varieties were released in Ethiopia, it was the local garlic variety (100%) grown in the two woredas (Table 5).

**Table 5.** Garlic varieties grown in the study areas

Garlic varieties	Malga		Gorche	
	Number of farmers	(%)	Number of farmers	(%)
Local	18	100	18	100
Qoricho	0	0	0	0
Tsedey	0	0	0	0
BishofituNech	0	0	0	0

Comment [A4]: It is not relevant to place a table showing only one result

Formatted: Font: Not Bold, Font color: Auto, Highlight

Formatted: Highlight

Formatted: Font: Not Bold, Font color: Auto, Highlight

Formatted: Highlight

Formatted: Font: Not Bold, Font color: Auto, Highlight

Formatted: Highlight

Formatted: Font: Not Bold, Font color: Auto, Highlight

Formatted: Highlight

Formatted: Font: Not Bold, Font color: Auto, Highlight

Formatted: Highlight

Formatted Table

Concerning to the application of fertilizer farmers used compost (100%) for the growth of garlic from the two woredas (Table 6).

**Table 6.** Type of fertilizer used for garlic production in the study areas

Type of fertilizer	Malga		Gorche	
	Number of farmers	(%)	Number of farmers	(%)
Urea	0	0	0	0
DAP	0	0	0	0
Compost	18	100	18	100

Formatted Table

Comment [A5]: It is not relevant to place a table showing only one result


The number of times that garlic was grown in the two woredas was presented in (Table 7). And the participants in Malga woreda informed that (44.4%) is 1 time and (55.6%) is 2 times a year. And it was (27.8%) and (72.2%) who responded 1 time and 2 times respectively in the case of Gorche woreda (Table 7).

**Table 7.** Number of times garlic grown in the study areas

Number of times	Malga		Gorche	
	Number of farmers	(%)	Number of farmers	(%)
1 time	8	44.4	6	33.3
2 times	10	55.6	12	66.7

Formatted Table

Age of the interviewed farmers was clearly presented in (figure 3). Farmers whose ages were 34-48(33.3%), >48(66.7%) and for age ranges <18 and 19-33 were (0%) in the case of Malga woreda. Whereas, it was (0%) for age ranges <18 and 19-33 and it was (27.8%) for 34-48 and (72.2%) for age ranges >48 in the case of Gorche woreda (Figure 3).



**Figure 3.** Age of the interviewed farmers in the study areas

#### 4. Discussion

According to the assessment results, all household heads of the participants in the two woredas had greater than fifteen years of experience in garlic production (Table 1). And this indicated that farmers in the study woredas had more experience in garlic production although they didn't have an access of improved garlic varieties. These results are generally in the contrary with the results presented earlier where most of the household heads in AdetZuria kebelewere young and thus might not have much experience in garlic production [14].

Mean garlic production area to be allocated in the two woreda was 0.05 ha (Table 3). This implies that garlic is cultivated by small scale subsistent farmers. The same farming pattern was reported in Jimma area, Oromia region, Ethiopia [2].

About 100% of the interviewed sample farmers in the two woredas used compost, the organic fertilizer for garlic production (Table 6). Application of fertilizers is an important cultural practice in vegetable production as it helps to satisfy the nutrient needs of crops required for the production of high yield and farmers from AdetZuria kebele used DAP and urea fertilizers to produce garlic [14].

Formatted: Line spacing: 1.5 lines

The use of improved garlic varieties is a very important input to produce high quantity and quality of garlic yield. Although different improved garlic varieties such as Qoricho, BishoftuNetch and Tseday 92 were released from different research institutes in Ethiopia [9]; farmers from Malga and Gorche woredas used only local garlic varieties sourced from their own savings and/or from the local market (Table 5). Planting materials sourced from such informal seed systems have limitations in terms of quality as the interviewed farmers mentioned which is in agreement with the observations of [8]. According to the key informants, unavailability of planting materials of improved garlic variety is a very limiting factor for the development of garlic production in the study area.

Diseases and insect pests are a serious concern in the production of garlic in Ethiopia including the study area. The current study therefore discovered that unavailability of high yielding certified seed varieties, acute shortage of fertilizers and pesticides and prevalence of leaf rust in the on-season are the main tribulations of garlic production where their occurrence differs with the study woredas (Table 4). The results are generally in agreement with the findings of various researchers who found that white rot and rust are the common diseases of garlic in Ethiopia [17,3, 15,14].

this manuscript lacks scientific discussion, it is necessary that the authors add the following paragraphs that would improve the scientific quality of the candidate manuscript:

Many researchers have studied sociodemographic characteristics for their effect on agricultural production [20](Olivares et al. 2016). In a study carried out in Venezuela, age and education had a significant positive influence on the adoption of soil conservation measures and therefore on the production of certain crops [22, 22](Olivares et al. 2017; Olivares, 2016). Also, the study by [23](Olivares and Franco, 2015) reported a positive influence of education on the adoption of conservation agriculture technology in tropical territories. Likewise, one study reported that female-headed households were more likely to adopt land and water technologies compared to male-headed households [24](Olivares et al. 2014). By age, older farmers were less likely to adopt practices that would improve agricultural production [25](Olivares et al 2012; 2012).

Formatted: Font: Not Bold, Font color: Auto, Highlight

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: Justified

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: Font: Not Bold, Font color: Auto, English (United States)

Formatted: Font: Not Bold, English (United States)

Formatted: Font: Not Bold

In order to identify suitable adaptation practices to a certain territory, it is necessary to previously establish the groups of actors involved, defined by similar characteristics from the point of view of land use, the production system[26](Olivares y Lopez, 2019), ethnicity[27](Olivares et al. 2020), culture[28](Olivares et al. 2020), the territory where they do agriculture and the productive chains to which they belong [29, 30, 31], [Olivares y Hernandez, 2019; Olivares et al. 2019, Olivares et al. 2020]. This stage facilitates the analysis of adaptation, since in general they will have similar threats, impacts and feasibility attributes compared to the proposed actions[32, 33](Montenegro et al. 2021; Pitti et al. 2021).

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: Not Bold

## 5. Conclusion

Garlic is an important cash crop for smallholder farmers in Ethiopia in general and the study area specifically. The production and productivity of garlic in study area is facing different constraints which requires research and extension services. Therefore, information about the current garlic production practices is paramount important to design intervention strategies.

The assessment of the present study showed that garlic in the study area was mostly produced by smallholder farmers in the study area allocated about 5.95% of their cropland for the production of garlic in Malga woreda and the farmers in Gorche woreda allocated relatively high proportion 6.58% of their crop land for garlic production.

Nevertheless, the study area has also huge opportunities for the improvement of garlic production which include the increase of demand and market price of garlic and awareness of the farmers towards the economic and nutritional values of garlic crop.

I suggest adding this paragraph that summarizes one of the conclusions:

Formatted: Font color: Auto, Highlight

This type of study represents an advance of great methodological utility because it provides an approach to study this type of issues in garlic production systems, with results that increase existing knowledge with theoretical and statistical foundations. Likewise, this methodology can be applied in other institutions, communities or organizations.

This research will form an important part in establishing the baseline on the location of the areas of importance for garlic in Ethiopia,..... (complete it)

## 6. Recommendations

The findings of the assessment indicated that both Malga and Gorche woreda farmers had greater than fifteen years of experience in the production of garlic. However, there are constraints which require various intervention strategies which include among others:

Upgrading the skills, knowledge and attitudes of the smallholder farmers in the agronomic management and providing improved garlic varieties which are resistant to disease and insects, fertilizers, pesticides, insecticides.

## References

[I suggest adding recent references which address the issue in question in Latin American territories. Suggested citations are for genuine scientific reasons that emphasize the current topic of study in context:](#)

1. Abay, A. Assessment of soil fertility status of different types of soils in selected areas of Southern Ethiopia. *Journal of Natural Sciences Research*. 2016; 6(1):2224-3186.
2. Adem BE and Tadesse S.T. Evaluating the Role of Nitrogen and Phosphorous on the Growth Performance of Garlic (*Allium sativum* L.). *Asian Journal of Agricultural Research*. 2014; 8(4): 211-217.
3. Agegnehu S, Walleign Z, Bayuh B, Fikremariam A, Abebe W, Dereje B, Esmelalem M, Getachew A. Diseases, insect pests and parasitic weeds of common crops and their importance under irrigation conditions of Rib areas. *Aizen publishers*. 2013; 1:262-265.
4. Alemu, D, Nigussie, D, Fikreyohannes, G. Effects of vermicompost and inorganic NP fertilizers on growth, yield and quality of garlic (*Allium sativum* L.) in Enebse Sar

- Midir District, Northwestern Ethiopia. *Journal of Biology, Agriculture and Healthcare*.2016; 6(3):2224-3208.
5. Befekadu, T, Missiganaw, W, Endeshaw, A. The traditional practice of farmers' legume-cereal cropping system and the role of microbes for soil fertility improvement in North Shoa, Ethiopia. *Agricultural Research and Technology*.2017; 13(4):1-6.
  6. Belay, Y. Integrated soil fertility management for better crop production in Ethiopia. *International Journal of Soil Science*.2015; 10:1-16.
  7. CSA. The population estimates derived from projecting the 1994 and 2007 census, analytical report for Southern Nations, Nationalities and People's Region. Central Statistical Agency. 2018b.
  8. Emanu B, Amsalu A, Tesfaye B, Milkessa T. Scoping Study on Vegetables Seed Systems and Policy in Ethiopia. 2014.
  9. Ethiopian Institute of Agricultural Research (EIAR). Guideline of crop production technologies. Retrieved from [www.eiar.gov.et/](http://www.eiar.gov.et/). 2007.
  10. Eyasu Elias, BeyeneTekluMellisse, Getachew Agegnehu and Desalegn Ayele. Response of Food Barley (*Hordeum Vugarae* L.) To Boron Blend Fertilizer Rates on Alisols in Southern Highlands of Ethiopia. 2020.
  11. Gyawali, S, Sunwar, S, Subedi, M, Tripathi, M, Joshi, KD and Witcombe, JR."Collaborative breeding with farmers can be effective", *Field Crops Research*. 2007;101: 88-95.
  12. Munyemana, A and Von Oppen, M. "La pomme de terre au Rwanda: uneanalyse d'une filière à hautes potentialités", *Centro Internacional de la Papa*, Lima. 1999.
  13. Ortiz, O, Frias, G, Ho, R, Cisneros, H, Nelson, R, Castillo, R, Orrego, R, Pradel, W, Alcazar, J and Bazán, M. "Organizational learning through participatory research: CIP and Care in Peru", *Agriculture and Human Values*. 2008;25: 419-431.

14. ShegeGetuYayeh, Melkamu Alemayehu, Amare Hailelassie and YigzawDessaegn. Assessment of small holder farmers garlic (*Allium sativum* L.) production practices under irrigated farming system in the Highlands of Ethiopia. African Journal of Agricultural Research. 2021 17(9): 1172-1179.

15. Shibu T. *In vitro* Evaluation of Aqua Extracts of Some Botanicals against Maize Stem Borer, *Busseolafusca* F. (Lepidoptera: Noctuidae). Journal of Plant Pathology and Microbiology.2013; 4:179.

16. Shivananda, IT. Integrated nutrient management on physical conditions, soil fertility and productivity of crops in dry land vertisol. Karnataka Journal of Agricultural Sciences. 2002; 15(1):186-187.

~~17.~~Wale M, Mengistu F. Agricultural potentials, constraints and opportunities in the Megech and Ribb rivers irrigation project areas in the Lake Tana Basin of Ethiopia, (December). <https://www.researchgate.net>. 2009.

~~17.~~

~~18.~~Witcombe, JR, Joshi, KD, Gyawali, S, Musa, AM, Johansen, C, Virk, DSand Sthapit, BR. "Participatory plant breeding is better described as highly client-oriented plant breeding I. Four indicators of client-orientation in plant breeding", Experimental Agriculture. 2005; 41 :299-319.

~~18.~~

~~19.~~Yadav, RN, Bairwa, HL, Gurjar, MK. Response of garlic (*Allium sativum* L.) to organic manures and fertilizers. International Journal of Current Microbiology and Applied Sciences.2017; 6(10):4860-4867.

~~19.~~

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Line spacing: 1.5 lines, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

Formatted: Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

Formatted: Font: (Default) +Body, 11 pt

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

~~20. Olivares, B., Lobo, D., Cortez, A., Rodríguez, M.F., y Rey, J.C. 2017. Socio-economic characteristics and methods of agricultural production of indigenous community Kashaama, Anzoátegui, Venezuela. Rev. Fac. Agron. (LUZ), 2017, 34 (2): 187-215. <https://n9.cl/p2gc5>~~

20.

~~21. Olivares, B., Zingaretti, M.L., DemeyZambrano, J.A., y Demey, J.R. 2016. Tipificación de los sistemas de producción agrícola y la percepción de la variabilidad climática en Anzoátegui, Venezuela (In spanish). Revista FAVE - Ciencias Agrarias, 2016, 15 (2): 39-50. <https://doi.org/10.14409/fa.v15i2.6587> Disponible en [http://www.scielo.org.ar/scielo.php?script=sci\\_arttext&pid=S1666-77192016000200004&lng=es&nrm=iso](http://www.scielo.org.ar/scielo.php?script=sci_arttext&pid=S1666-77192016000200004&lng=es&nrm=iso)~~

21.

~~22. Olivares, B. 2016. Description of soil management in agricultural production systems of sector hammock in Anzoátegui, Venezuela. Descripción del manejo de suelos en sistemas de producción agrícola del sector Hamaca de Anzoátegui, Venezuela. La Granja: Revista de Ciencias de la Vida, 2016, 23(1): 14-24. <https://doi.org/10.17163/lgr.n23.2016.02>~~

22.

~~23. Olivares, B., y Franco, E. 2015. Agrosocial diagnostic of the indigenous community of Kashaama: An empirical study in the state of Anzoátegui, Venezuela. Diagnostico agrosocial de la comunidad indígena de Kashaama: Un estudio empírico en el estado Anzoátegui, Venezuela. Revista Científica Guillermo de Ockham, 2015, 13 (1): 87-95. <http://dx.doi.org/10.21500/22563202.1691>~~

23.

~~24. Olivares, B. 2014. Application of Principal Component Analysis (PCA) in Socio-Environmental Diagnosis. Case: The Campo Alegre Sector, Simón Rodríguez Municipality, Anzoátegui. Aplicación del Análisis de Componentes Principales (ACP) en el diagnóstico socio ambiental. Caso: sector Campo Alegre, municipio Simón Rodríguez de Anzoátegui, j. Revista Multiciencias, 2014, 14 (4): 364 - 374. <https://n9.cl/caoee>~~

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Font color: Auto, Spanish (Spain, International Sort)

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Font color: Auto, Spanish (Spain, International Sort)

Formatted: Font: Not Bold, Font color: Auto

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, Font color: Auto, Spanish (Spain, International Sort)

Formatted: Spanish (Spain, International Sort)

Formatted: Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

Formatted: Font: +Body, 11 pt, Not Bold, No underline, Font color: Auto, Portuguese (Brazil)

Formatted ... [1]

Formatted ... [2]

Formatted ... [3]

Formatted ... [4]

Formatted ... [5]

Formatted ... [6]

Formatted: Font: (Default) +Body, 11 pt

Formatted ... [7]

Formatted ... [8]

Formatted ... [9]

Formatted ... [10]

Formatted ... [11]

Formatted: Spanish (Spain, International Sort)

Formatted ... [12]

Formatted ... [13]

Formatted ... [14]

Formatted ... [15]

Formatted ... [16]

Formatted ... [17]

Formatted ... [18]

Formatted ... [19]

Formatted ... [20]

Formatted ... [21]

Formatted ... [22]

Formatted ... [23]

Formatted ... [24]

Formatted ... [25]

Formatted ... [26]

Formatted ... [27]

24. \_\_\_\_\_

25. Olivares B., Guevara E. y Demey J. ~~2012~~ The Use of Climate Biomarkers in Agricultural Production Systems, Anzoátegui, Venezuela *Utilización de bioindicadores climáticos en sistemas de producción agrícola del estado Anzoátegui, Venezuela* (In Spanish). *Revista Multiciencias*. 2012; 12 (2): 136-145. <https://n9.cl/ak22r>

25. \_\_\_\_\_

26. Olivares B., Guevara E. y Demey J. ~~2012~~ Uso y demanda de información agrometeorológica en los sistemas de producción agrícola en Anzoátegui, Venezuela. *Revista Multiciencias*. 2012; 12 (4): 372-381. <https://n9.cl/jsfpj>

26. \_\_\_\_\_

27. Olivares B., López-Beltrán M., Lobo-Luján D. Changes in land use and vegetation in the Kashaama agrarian community, Anzoátegui, Venezuela: 2001-2013 *Cambios de usos de suelo y vegetación en la comunidad agraria Kashaama, Anzoátegui, Venezuela: 2001-2013*. *Revista Geográfica De América Central*. 2019, 2(63), 269-291. <https://doi.org/10.15359/rgac.63-2.10>

27. \_\_\_\_\_

28. Olivares B., López M. Normalized Difference Vegetation Index (NDVI) applied to the agricultural indigenous territory of Kashaama, Venezuela. *UNED Research Journal*. 2019, 11(2), 112-121. <https://doi.org/10.22458/urj.v11i2.2299>

28. \_\_\_\_\_

29. Olivares B., Hernández R. ~~2019~~ Ecoterritorial sectorization for the sustainable agricultural production of potato (*Solanum tuberosum* L.) in Carabobo, Venezuela. *Agricultural Science and Technology*. 2019; 20(2): 339-354. [https://doi.org/10.21930/rcta.vol20\\_num2\\_art:1462](https://doi.org/10.21930/rcta.vol20_num2_art:1462)

29. \_\_\_\_\_

- Formatted: Spanish (Spain, International Sort)
- Formatted: Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"
- Formatted: Font: (Default) Times New Roman, 12 pt
- Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, No underline, Font color: Auto, Portuguese (Brazil)
- Formatted: Font: (Default) Times New Roman, 12 pt
- Formatted: Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"
- Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, No underline, Font color: Auto, Portuguese (Brazil)
- Formatted: No underline, Font color: Auto, Portuguese (Brazil)
- Formatted: Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto, Portuguese (Brazil)
- Formatted: Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto
- Formatted: Font: (Default) Times New Roman, 12 pt
- Formatted: Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"
- Formatted ... [28]
- Formatted ... [29]
- Formatted ... [30]
- Formatted ... [31]
- Formatted ... [32]
- Formatted ... [33]
- Formatted: Spanish (Spain, International Sort)
- Formatted: Line spacing: Multiple 1.15 li
- Formatted ... [34]
- Formatted ... [35]
- Formatted ... [36]
- Formatted ... [37]
- Formatted: English (United States)
- Formatted ... [38]
- Formatted ... [39]
- Formatted ... [40]
- Formatted ... [41]
- Formatted ... [42]

30. Olivares, B.; Hernandez, R.; Arias, A; Molina, JC., Pereira, Y. ~~2020~~ Eco-territorial adaptability of tomato crops for sustainable agricultural production in Carabobo, Venezuela. *Idesia*, 2020; 38(2):95-102. <http://dx.doi.org/10.4067/S0718-34292020000200095>

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, No underline, Font color: Auto

30.

Formatted: Font: (Default) Times New Roman, 12 pt

Formatted: Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

31. Olivares, B., Pitti, J., Montenegro, E. ~~2020~~ Socioeconomic characterization of Bocas del Toro in Panama: an application of multivariate techniques. *Revista Brasileira de Gestao e Desenvolvimento Regional*, 2020; 16(3):59-71

Formatted: Font: (Default) Times New Roman, 12 pt, Not Bold, No underline, Font color: Auto

31.

Formatted: Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto

Formatted: Font: (Default) Times New Roman, 12 pt

32. Montenegro-Gracia, Edilberto Javier, Pitti-Rodríguez, Jacob Eduardo, & Olivares-Campos, Barlin Orlando. ~~(2021)~~ Identificación de los principales cultivos de subsistencia del Teribe: un estudio de caso basado en técnicas multivariadas. *Idesia (Arica)*, ~~(2021)~~; 39(3), 83-94. <https://dx.doi.org/10.4067/S0718-34292021000300083>

Formatted: Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

Formatted: Font: (Default) Times New Roman, 12 pt, No underline, Spanish (Spain, International Sort)

32.

Formatted: Font: (Default) Times New Roman, 12 pt, No underline, Spanish (Spain, International Sort)

Pitti Rodríguez, J., Olivares, B., Montenegro, E., Miller, L., & Ñango, Y. (2021). The role of agriculture in the Cehanguinola District: a case of applied economics in Panama. *Tropical and Subtropical Agroecosystems*, 2021; 25(1). Retrieved from <https://n9.cl/quy12> <https://www.revista.ceba.uady.mx/ojs/index.php/TSA/article/view/3815>

Formatted: Font: (Default) Times New Roman, 12 pt, Not Italic, No underline

Formatted: Font: (Default) Times New Roman, 12 pt, No underline

Formatted: Font: (Default) Times New Roman, 12 pt, No underline

Formatted: Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto

Formatted: Font: (Default) Times New Roman, 12 pt

33.

Formatted ... [43]

## Appendix

Formatted ... [44]

Formatted ... [45]

Formatted ... [46]

Formatted ... [47]

Formatted ... [48]

Formatted ... [49]

Formatted ... [50]

Formatted ... [51]

Formatted ... [52]

Formatted ... [53]

Formatted: English (United States)

Formatted ... [54]

Formatted: Spanish (Spain, International Sort)

## Questionnaires for assessment of production of garlic

### Part I. Demographic information

1. Name of farmer/Code: \_\_\_\_\_
2. Sex of household head a. Female b. Male
3. Age of household in years a). <18 b). 19-33 c). 34-48 d). >48

4. Educational status of household head a). literate b). Illiterate c). 1-6 grade d). 8-10 grades

5. District \_\_\_\_\_ Kebele \_\_\_\_\_

## Part II. Farm practices

1. What is your total land holding?

a). <0.50 ha b). 0.50-1.0 ha c). 1.0-1.50 ha d). >1.50 ha

2. What size of your land is dedicated to garlic production?

a). <0.25ha b). 0.25-0.50ha c).0.50-0.75ha d).1.0ha e).>1.0ha

3. How long is your experience in garlic production?

a. <5 years b. 5-10 years c.10-15 years d. > 15 years

4.Where do you get your planting materials from?

a. from traders b. agricultural bureau c. research institutions

d. private companies e. farmers keep their own seeds from their harvests

5. Which variety of garlic do you grow?

a. BishoftuNech b. Tsedey c. Qoricho d. Local

6. How many times in a year do you produce garlic?

a. once b. twice c. trice

7. Please provide on the types of fertilizers used

**Table 1.** Type of fertilizer used in the study area

Type	Yes	No
Urea		
DAP		

Compost

Other

**Part III: Pest (Disease, Insect and Weed) Control**

1. What is the most important garlic pest in this area?
  - a. diseases
  - b. insects
  - c. weeds
  - d. others
2. What other control method (s) do you use against disease?
  - a. chemical
  - b. cultural
  - c. both

UNDER PEER REVIEW

**Page 15: [1] Formatted** **Autor**

No underline, Font color: Auto, Portuguese (Brazil)

**Page 15: [2] Formatted** **Autor**

Font: +Body, 11 pt, No underline, Font color: Auto, Portuguese (Brazil)

**Page 15: [3] Formatted** **Autor**

Font: +Body, 11 pt, No underline, Font color: Auto, Spanish (Spain, International Sort)

**Page 15: [4] Formatted** **Autor**

Font: Not Bold, No underline, Font color: Auto

**Page 15: [5] Formatted** **Autor**

Font: +Body, 11 pt, No underline, Font color: Auto, Spanish (Spain, International Sort)

**Page 15: [6] Formatted** **Autor**

Font: +Body, 11 pt, No underline, Font color: Auto, Spanish (Spain, International Sort)

**Page 15: [7] Formatted** **Autor**

Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

**Page 15: [8] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, Not Bold, No underline, Font color: Auto

**Page 15: [9] Formatted** **Autor**

No underline, Font color: Auto, English (United States)

**Page 15: [10] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto

**Page 15: [11] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto, Spanish (Spain, International Sort)

**Page 15: [12] Formatted** **Autor**

Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

**Page 15: [13] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, Not Bold, No underline, Font color: Auto

**Page 15: [14] Formatted** **Autor**

No underline, Font color: Auto, English (United States)

**Page 15: [15] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, Not Bold, No underline, Font color: Auto

**Page 15: [16] Formatted** **Autor**

No underline, Font color: Auto, English (United States)

**Page 15: [17] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto

**Page 15: [18] Formatted** **Autor**

No underline, Font color: Auto, English (United States)

**Page 15: [19] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto

**Page 15: [20] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt

**Page 15: [21] Formatted** **Autor**

Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

**Page 15: [22] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, Not Bold, No underline, Font color: Auto

**Page 15: [23] Formatted** **Autor**

No underline, Font color: Auto, English (United States)

**Page 15: [24] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto

**Page 15: [25] Formatted** **Autor**

No underline, Font color: Auto, English (United States)

**Page 15: [26] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto

**Page 15: [27] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto, Spanish (Spain, International Sort)

**Page 16: [28] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto

**Page 16: [29] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt

**Page 16: [30] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, Spanish (Spain, International Sort)

**Page 16: [31] Formatted** **Autor**

List Paragraph, Right: 0", Line spacing: 1.5 lines, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

**Page 16: [32] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt

**Page 16: [33] Formatted** **Autor**

List Paragraph, Line spacing: 1.5 lines, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

**Page 16: [34] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto

**Page 16: [35] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt

**Page 16: [36] Formatted** **Autor**

Font: (Default) Times New Roman, 12 pt, English (United States)

**Page 16: [37] Formatted** **Autor**

List Paragraph, Right: 0", Line spacing: 1.5 lines, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

**Page 16: [38] Formatted** **Autor**

Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

**Page 16: [39] Formatted** **Autor**

Page 16: [40] Formatted Autor

Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto

Page 16: [41] Formatted Autor

Font: (Default) Times New Roman, 12 pt

Page 16: [42] Formatted Autor

Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

Page 17: [43] Formatted Autor

Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

Page 17: [44] Formatted barlin olivares 2/19/2022 2:31:00 PM

Justified, Numbered + Level: 1 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.5" + Indent at: 0.75"

Page 17: [45] Formatted Autor

Font: (Default) Times New Roman, 12 pt, No underline, Spanish (Spain, International Sort)

Page 17: [46] Formatted Autor

Font: (Default) Times New Roman, 12 pt, No underline

Page 17: [47] Formatted Autor

Font: (Default) Times New Roman, 12 pt, Not Italic

Page 17: [48] Formatted Autor

Font: Not Italic, English (United States)

Page 17: [49] Formatted Autor

Font: (Default) Times New Roman, 12 pt, Not Italic

Page 17: [50] Formatted Autor

Font: Not Italic, English (United States)

Page 17: [51] Formatted Autor

Font: (Default) Times New Roman, 12 pt, Not Italic

Page 17: [52] Formatted Autor

Font: (Default) Times New Roman, 12 pt, No underline, Font color: Auto

Page 17: [53] Formatted

Autor

Font: (Default) Times New Roman, 12 pt

Page 17: [54] Formatted

Autor

No underline, Font color: Auto, Spanish (Spain, International Sort)