

Serpentine Thrombus of the Right Cavities of the Heart Extending to the Trunk of the Pulmonary Artery Complicating a Uterine Leiomyosarcoma without Pulmonary Embolism: Case Report

ABSTRACT

Serpentine thrombus or snake like floating thrombus of the right-side cavities of the heart is a rare diagnosis. Its an extreme emergency, and is often fatal due to the migration of small thrombus which can completely obstruct the pulmonary trunk. Advanced cancer disease is one of the rare causes of this diagnosis. Two-dimensional transthoracic echocardiography helps to confirm the diagnosis while Angio CT-Scan confirms the presence of pulmonary embolism in patients presenting acute or aggravative dyspnea. Treatment may involve surgical thrombectomy or intravenous thrombolysis but the use of medical treatment alone (that is oral or subcutaneous injectable anticoagulants) has never been an effective choice of therapy except in exceptional cases.

We report a rare case of a patient followed at the oncology department for locally advanced uterine leiomyosarcoma under chemotherapy who presented aggravating symptom of dyspnea at the emergency department where a serpentine thrombus was diagnosed by transthoracic echocardiography. Angio computed tomography scan (Angi-CT SCAN) showed no signs of pulmonary embolism in the patient who finally was placed on low molecular weight heparin (LMWH) due to high bleeding risk with a good outcome under strict hospital surveillance. This case reflects the difficulty of therapy strategies in some population of patients with high risk of mortality due to the presence of floating thrombus associated with high bleeding risk.

Keywords: Serpentine thrombus; uterine leiomyosarcoma; pulmonary embolism.

1. INTRODUCTION

Floating thrombus (serpentine thrombus) of the right-side cavities of the heart is a rare diagnosis and constitutes a real therapeutic emergency [1]. Serpentine thrombus seen in the right atrial is very pathognomonic of thrombus being dislodged from the deep veins of the lower limbs. It's particularly a serious clinical form of thromboembolic disease usually associated with severe pulmonary embolism [1,2]. A thrombus is the most common cardiac mass. The most common locations for cardiac thrombi are the left atrium and left atrial appendage, especially in the setting of atrial fibrillation or a dysfunctional left ventricle [3]. Uterine leiomyosarcoma (uLMS) is a rare entity among malignant gynecologic tumors with a very unfavorable prognosis and also rare cause of thrombus intracardiac [4,5].

We present the case of a patient followed at the oncology department for a uterine leiomyosarcoma at the stage of pulmonary metastases, and in whom a transthoracic

echocardiography was performed due to aggravating dyspnea symptom, which objectified a serpentine thrombus of the inferior vena cava extended to the right-side cavities of the heart up to the trunk of the pulmonary artery without pulmonary embolism who pose a difficulty in the choice of anticoagulation therapy.

2. CASE REPORT

We report the case of a female patient aged 47, with no cardiovascular risk factors. She has been followed since April 2022 for uterine leiomyosarcoma under chemotherapy, revealed clinically by metrorrhagia of which the diagnosis was retained on histological evidence of a peritoneal sample. Patient presents at the emergency department for worsening dyspnea symptom that has been evolving for 3 weeks, with no other associated clinical symptoms.

Somatic examination found a hemodynamically stable patient, polypnea at 24 cycles par minute with a left parasternal presystolic murmur

coded 3/6th at the level of the fourth intercostal space. We also noted a great abundance of ascite during abdominal exam and bilateral white soft pitting sloping and painless oedemas of the lower limbs. Pulmonary auscultation was normal. The biological assessment objectified a microcytic hypochromic anemia at 8g/dl with mild thrombocytopenia, platelet levels at 135,000U/l.

Transthoracic echocardiography (Figs. 1, 2) performed showed a "serpiginous" hyperechoic intra-cavitary mass at the right-side cavities of the heart with no adherence to the heart cavity walls, floating and extending to the trunk of the pulmonary artery. In addition, the right cavities were not dilated, the RV/LV ratio <1, with a conserved biventricular function without no other abnormalities associated.

A venous vascular doppler ultrasound studying of the deep veinous network of the lower limbs was performed, which confirmed the absents of deep vein thrombosis.

An Angio CT- scan (Fig. 3) was also performed, revealing no pulmonary embolism neither in the proximal nor the distal segmentation of the lungs.

The patient was placed on low molecular weight heparin (LMWH) at a curative dose according to her weight despite the moderate

hypochromic microcytic anemia of 8 g/dl and mild thrombocytopenia. No thrombolysis was performed due to the absence of serious pulmonary embolism and also due to the high risk of hemorrhage with a HASBLED score ≥ 3 (bleeding risk score). Oral antibiotics, amoxicillin was simultaneous given to the patient on preventive basis of pulmonary infection.

A good clinical and radiological outcome was noted. Improved respiratory state of the patient was observed one week under anticoagulation treatment with an oxygen saturated level of 97% in ambient air. Disappearance of the intra-cavitary thrombus 1 month after the installation of anticoagulation therapy was marked in the patient with a minor vaginal bleeding side effect without the need of transfusion.

3. DISCUSSION

A floating thrombus of the right-side cavities of the heart is rare, it has been the subject of publications to one or more cases. It can be an extreme emergency, and is often fatal due to the migration of small thrombus which completely obstruct the pulmonary trunk. According to studies, it concerns 7 to 11% of thromboembolic diseases, thus justifying the systematic performance of echocardiography as soon as a pulmonary embolism is suspected [6].




Fig. 1. Transthoracic Echocardiography (TTE): 4 apical chamber view: Tail of the serpentine thrombus floating in the right atrium of the heart measuring 1cm long without dilated right atrium




Fig. 2. TEE: Small axis view: Head and the middle piece of the serpentine thrombus in the right ventricle during diastole measuring 2.5cm on its long axis




Fig. 3. Angio CT-scan: Parenchymal view: Absence of pulmonary embolism

The circumstances of discovery are diverse: most often, there is a context of severe or recurrent pulmonary embolism [6,7], with clinical signs suggestive of acute cor pulmonale. Thrombosis of the lower limbs is not necessarily present in case of severe pulmonary embolism but can be orienting sign [2,8]. A tricuspid obstruction syndrome can be observed, resulting in paroxysmal syncope [7] or sudden death [9]. Although less common and of unusual location, thrombosis of the inferior vena cava is not exceptional and can be the cause of thrombus in the right cavities by extension. Basically, when it is present, it results in bilateral signs that occur in one or two successive stages (tilting venous thrombosis). On the other hand, in our case, the patient does not present either pulmonary embolism or deep vein thrombosis despite the echocardiographic appearance of a long significant serpentine thrombus of the right-side cavities of the heart.

Two-dimensional transthoracic echocardiography makes the diagnosis [9,10,11,12].

It most often shows dilated right cavities and a "serpentine" hyperechoic image of the prolapsed right atrium between the tricuspid valves towards the right ventricle [7] whereas in our patient, there was no dilated right atrium which can simply be explained by absents of total obstacle of the pulmonary trunk. In the context of pulmonary embolism, echocardiographic images of the floating thrombus and cor pulmonale are sufficient to

suggest the diagnosis, without resorting to pulmonary angiography [9,7]. In our case an angio CT-Scan was done as the patient was stable. The prognosis of floating thrombus of the right heart is reserved but mortality is very high, greater than 40% in some studies [13, 14]. This poor prognosis is linked to the combination of several factors: very high embolic risk, severe pulmonary embolism associated with detection of the thrombus, peripheral phlebothrombosis, source of recurrent embolic events.

Treatment may involve surgical thrombectomy or intravenous thrombolysis. AngioVac (AngioDynamics) large-bore catheter system can also be used for the removal of large thrombi in various clinical situations [14,15]. In our case, the patient was placed on LWMH at a curative therapy dose alone with good clinical outcome given the high risk of bleeding score in the patient.

The echocardiographic control of our patient showed the absence of intra-auricular thrombus one month after anticoagulation treatment. The literature reports only a few rare cases of favorable evolution under medical treatment alone [16].

This case report is original, as it concerns a rare case of extensive thrombosis, from the inferior vena cava to the trunk of the pulmonary artery, without echocardiographic signs of acute cor pulmonale, or pulmonary embolism at thoracic CT angio-Scan with a favorable

outcome under heparin therapy alone despite the patient's high score bleeding risk.

4. CONCLUSION

Floating thrombus “serpentine thrombus” of the right-side cavities of the heart is a rare but extremely serious event. It should be considered an immediate threat; because if left untreated, mortality can reach 90%. The association of right chamber thrombosis and pulmonary embolism has a poor prognosis. Heparin is often insufficient even when the clinical condition is stable, but in the face of a high risk of bleeding, heparin alone can be of benefit to the patient under strict surveillance.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

CONSENT

As per international standard or university standard, patient(s) written consent has been collected and preserved by the author(s).

ACKNOWLEDGEMENT

I thank the whole Department of Cardiology P37, university teaching hospital Ibn Rochd Casablanca for their outstanding collaboration in the support of this work.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Vidyashree Hulkoti, Sandeep Kamat, Sourya Acharya, Samarth Shukla, Sunil Kumar, Dhruv Talwar. Serpentine Thrombus in Right Atrium: A Tell-Tale Sign of Venous Thromboembolism. *Cureus*. 2021 Dec 10;13:e20336.
2. Bagga C, Ansari I, Garikapathi A, Irshadv, Naik S, Agrawal S, Kumar S. Status epilepticus presenting feature of pulmonary embolism: rarest of rare combination. *Med Sci*. 2020;24:3277-80.
3. Elizabeth M Johnson, Kenneth L Gage, Sebastian Feuerlein, Daniel Jeong. Cardiac Magnetic Resonance for the Evaluation of Suspected Cardiac Thrombus: Conventional and Emerging Techniques. *J Vis Exp*. 2019;11;148.
4. Ingolf Juhasz-Böss, Lena Gabriel, Rainer M Bohle, Lars C Horn, Erich-Franz Solomayer, Georg-Peter Breitbach. Uterine Leiomyosarcoma. *Oncol Res Treat*. 2018; 41(11):680-686.
5. Peng YJ, Hueng GG, Lee HS. Acute heart failure as manifestation of metastatic uterine leiomyosarcoma to the heart and lung. *Heart Lung*. 2004;33; 46-9.
6. Farfel Z, Shechier M, and al. Review of echocardiographically diagnosed right heart entrapment of pulmonary emboli-in-transit with emphasis on management. *Am Heart J*. 1987; 117:171-178.
7. Gras D, Bedossa M, Leheloco A, et al. Thrombose auriculaire droite et embolie pulmonaire grave. *Arch Mal Cœur*. 1992; 85:253-256,
8. Taupin JM, Chapoutot L, Metz D, et al. Thrombus flottant de l'oreillette droite dans l'embolie pulmonaire aigue. Aspects chroniques, écho cardiographiques et incidences thérapeutiques. *Ann Cardiol Angéiol*. 1988;37:243-247.
9. Benichou M, Bernard P J, Sarrat P, et al. Détection des caillots intracardiaques par la scintigraphie aux plaquettes marquées à l'indium 111 Confrontation avec l'échographie bidimensionnelle et la scanographie cardiaque. *Arch Mal Cœur*. 1984;77:1054-1063.
10. Agnes Mayr, Gert Klug, Simon Greulich, Bernhard Metzler, Gudrun Feuchtnr. Serpentine-like right atrial mass and fulminant bilateral pulmonary embolism during treatment with rivaroxaban. *Int J Cardiovasc Imaging*. 2016;3:1001-2.
11. D. Attias, E. Abitbol, S. Russel, J. Paziand, P. Lim, C. Roiron, J.-L. Monin, P. Guéret, J. Garot Thrombus serpentin récidivant des cavités cardiaques droites. *Presse Med*. 2005;34:1337.
12. Yastrebov K, Brunel L, Paterson HS, Williams ZA, Bannon PG. Three dimensional intracardiac echocardiography and pulmonary embolism. *Cardiovasc Ultrasound*. 2020 Aug 20;18(1):36.
13. Felner JM, Churchwell AL, Murphy DA. Right atrial thromboemboli: Clinical, echocardiographic and pathologic manifestations. *J Am Coll Cardiol*. 1984;4: 1041-1051.

14. Patnaik S, Rammohan HS, Shah M, Garg S, Figueredo V, Janzer S, Shah S. Percutaneous Embolectomy of Serpentine Thrombus from the Right Atrium in a 51-Year-Old Man. *Tex Heart Inst J.* 2016 Dec 1;43:524-527.
15. Rosenfeld BL, Bashir R, Brisco-Bacik MA, Panidis IP, Vaidya A, Minakata K, Forfia PR. Leiomyosarcoma Tumor Embolism Masquerading as Thrombus in Transit. *Am J Case Rep.* 2020;29:921124.
16. Corman C, Roudaut R, Gosse P, et al. Thromboses de l'oreillette droite: Aspect échocardiographies, conduite pratique. A propos de 8 cas. *Arch Mal Cœur.* 1986; 79:464-471.